

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

March 2002

First record of the family Roproniidae (Hymenoptera: Proctotrupoidea) from Florida

Charles C. Porter

*Florida state Collection of Arthropods, Bureau of Entomology, Division of Plant Industry, Florida
Department of Agriculture & Consumer Services, Gainesville, FL*

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

Porter, Charles C., "First record of the family Roproniidae (Hymenoptera: Proctotrupoidea) from Florida" (2002). *Insecta Mundi*. 536.

<https://digitalcommons.unl.edu/insectamundi/536>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

First record of the family Roproniidae (Hymenoptera: Proctotrupoidea) from Florida

Charles C. Porter

Florida State Collection of Arthropods
Florida Department of Agriculture and Consumer Services
P.O. Box 147100
Gainesville, Florida 32614-7100

The Holarctic genus *Ropronia* Provancher has three species in North America: *P. californica* Ashmead (California, Oregon), *P. pediculata* Provancher (New York, Ontario), and *R. garmani* Ashmead known previously in the United States from New Hampshire to northern Georgia and west to Iowa (Townes 1948).

Recently, as part of an on-going survey of Hymenoptera in Florida State Parks, the following new record of *Ropronia garmani* was obtained:

Material Examined. Two females, Florida, Suwannee Co., Suwannee River State Park, riparian hardwoods, in Malaise Trap, 2-23-V-1997, C. Porter, L. Stange. In FSCA.

Habitat. Townes (1948) found *R. garmani* in undergrowth beside streams in Temperate Deciduous Forest. Florida specimens came from similar habitats in riparian hardwood forest along the Suwannee River.

Phaenology. In the northeast, *R. garmani* flies from mid-June to August; whereas, Florida records are for May, suggesting that in Florida this species avoids the heat of summer, as do most ichneumonids and other parasitic Hymenoptera.

Hosts. *Ropronia* may parasitize tenthredinid saw-fly larvae (Hymenoptera), but evidence is inconclusive (Townes 1948).

Biogeography. *Ropronia* has five species in China (Townes 1948; He, Zhu & Tong 1988), two in

Japan (Yasumatsu 1956), one in Burma (Heqvist 1959), and two in Turkey (Madl 1991). Roproniidae thus is a Holarctic group with greatest diversity in the eastern Palaearctic.

Identification. Fore wing length ± 5.0 mm; antenna with 14 segments; fore wing with a costal cell and with a small, quadrate or hexagonal first discoidal cell; first gastric segment petiolate (3.0-4.0 as long as deep), rest of gaster strongly compressed, much higher than broad.

Acknowledgments. Drs. Lionel Stange and Mike Thomas helped with museum and fieldwork. The Florida Department of Parks and Recreation issued collecting permits.

Literature cited

- He, J., Zhu K. & Tong, K.** 1988. Description of four new species of the genus *Ropronia* from China. *Entomotaxonomica* 10: 207-214.
- Heqvist, K.** 1959. A new species of *Ropronia* from Burma. *Entomologisk Tidskrift*. 80: 137-139.
- Madl, M.** 1991. Zwei neue *Ropronia*-Arten aus Turkei. *Linzer Biol. Beitr.* 23: 387-392.
- Townes, H.K.** 1948. The serphoid Hymenoptera of the family Roproniidae. *Proc. U.S. Natl. Museum* 98: 85-89.
- Yasumatsu, K.** 1956. Two new species of Roproniidae (Hymenoptera). *Insecta Matsumurana* 19: 117-122.