

2017

Ocean changes – warming, stratification, circulation, acidification, and deoxygenation

Libby Jewett

NOAA Ocean Acidification Program

Anastasia Romanou

Columbia University, anastasia.romanou@nasa.gov

Follow this and additional works at: <https://digitalcommons.unl.edu/usdeptcommercepub>

Jewett, Libby and Romanou, Anastasia, "Ocean changes – warming, stratification, circulation, acidification, and deoxygenation" (2017). *Publications, Agencies and Staff of the U.S. Department of Commerce*. 580.
<https://digitalcommons.unl.edu/usdeptcommercepub/580>

This Article is brought to you for free and open access by the U.S. Department of Commerce at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications, Agencies and Staff of the U.S. Department of Commerce by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Ocean changes – warming, stratification, circulation, acidification, and deoxygenation

Libby Jewett, NOAA Ocean Acidification Program
Anastasia Romanou, Columbia University

Citation: In: *Climate Science Special Report: A Sustained Assessment Activity of the U.S. Global Change Research Program* [Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock (eds.)]. U.S. Global Change Research Program, Washington, DC, USA (2017), pp. 540-583.

Comments: U.S. Government work

Abstract

1. The world's oceans have absorbed about 93% of the excess heat caused by greenhouse gas warming since the mid-20th century, making them warmer and altering global and regional climate feedbacks. Ocean heat content has increased at all depths since the 1960s and surface waters have warmed by about $1.3^{\circ} \pm 0.1^{\circ}\text{F}$ ($0.7^{\circ} \pm 0.08^{\circ}\text{C}$) per century globally since 1900 to 2016. Under a high emissions scenario, a global increase in average sea surface temperature of $4.9^{\circ} \pm 1.3^{\circ}\text{F}$ ($2.7^{\circ} \pm 0.7^{\circ}\text{C}$) by 2100 is projected, with even higher changes in some U.S. coastal regions. (*Very high confidence*)
2. The potential slowing of the Atlantic Meridional Overturning Circulation (AMOC) (of which the Gulf Stream is one component)—as a result of increasing ocean heat content and freshwater driven buoyancy changes—could have dramatic climate feedbacks as the ocean absorbs less heat and CO₂ from the atmosphere. This slowing would also affect the climates of North America and Europe. Any slowing documented to date cannot be directly tied to anthropogenic forcing primarily due to lack of adequate observational data and to challenges in modeling ocean circulation changes. Under a high emissions scenario (RCP8.5) in CMIP5 simulations, it is likely that the AMOC will weaken over the 21st century by 12% to 54%. (*Low confidence*)
3. The world's oceans are currently absorbing more than a quarter of the CO₂ emitted to the atmosphere annually from human activities, making them more acidic (*very high confidence*), with potential detrimental impacts to marine ecosystems. In particular, higher-latitude systems typically have a lower buffering capacity against pH change, exhibiting seasonally corrosive conditions sooner than low-latitude systems. Acidification is regionally increasing along U.S. coastal systems as a result of upwelling (for example, in the Pacific Northwest) (*high confidence*), changes in freshwater inputs (for example, in the Gulf of Maine) (*medium confidence*), and nutrient input (for example, in urbanized estuaries) (*high confidence*). The rate of acidification is unparalleled in at least the past 66 million years (*medium confidence*). Under RCP8.5, the global average surface ocean acidity is projected to increase by 100% to 150% (*high confidence*).
4. Increasing sea surface temperatures, rising sea levels, and changing patterns of precipitation, winds, nutrients, and ocean circulation are contributing to overall declining oxygen concentrations at intermediate depths in various ocean locations and in many coastal areas. Over the last half century, major oxygen losses have occurred in inland seas, estuaries, and in the coastal and open ocean (*high confidence*). Ocean oxygen levels are projected to decrease by as much as 3.5% under the RCP8.5 scenario by 2100 relative to preindustrial values (*high confidence*).

13. Ocean Acidification and Other Ocean Changes

KEY FINDINGS

1. The world's oceans have absorbed about 93% of the excess heat caused by greenhouse gas warming since the mid-20th century, making them warmer and altering global and regional climate feedbacks. Ocean heat content has increased at all depths since the 1960s and surface waters have warmed by about $1.3^{\circ} \pm 0.1^{\circ}\text{F}$ ($0.7^{\circ} \pm 0.08^{\circ}\text{C}$) per century globally since 1900 to 2016. Under a high emissions scenario, a global increase in average sea surface temperature of $4.9^{\circ} \pm 1.3^{\circ}\text{F}$ ($2.7^{\circ} \pm 0.7^{\circ}\text{C}$) by 2100 is projected, with even higher changes in some U.S. coastal regions. (*Very high confidence*)
2. The potential slowing of the Atlantic Meridional Overturning Circulation (AMOC) (of which the Gulf Stream is one component)—as a result of increasing ocean heat content and freshwater driven buoyancy changes—could have dramatic climate feedbacks as the ocean absorbs less heat and CO_2 from the atmosphere. This slowing would also affect the climates of North America and Europe. Any slowing documented to date cannot be directly tied to anthropogenic forcing primarily due to lack of adequate observational data and to challenges in modeling ocean circulation changes. Under a high emissions scenario (RCP8.5) in CMIP5 simulations, it is likely that the AMOC will weaken over the 21st century by 12% to 54%. (*Low confidence*)
3. The world's oceans are currently absorbing more than a quarter of the CO_2 emitted to the atmosphere annually from human activities, making them more acidic (*very high confidence*), with potential detrimental impacts to marine ecosystems. In particular, higher-latitude systems typically have a lower buffering capacity against pH change, exhibiting seasonally corrosive conditions sooner than low-latitude systems. Acidification is regionally increasing along U.S. coastal systems as a result of upwelling (for example, in the Pacific Northwest) (*high confidence*), changes in freshwater inputs (for example, in the Gulf of Maine) (*medium confidence*), and nutrient input (for example, in urbanized estuaries) (*high confidence*). The rate of acidification is unparalleled in at least the past 66 million years (*medium confidence*). Under RCP8.5, the global average surface ocean acidity is projected to increase by 100% to 150% (*high confidence*).
4. Increasing sea surface temperatures, rising sea levels, and changing patterns of precipitation, winds, nutrients, and ocean circulation are contributing to overall declining oxygen concentrations at intermediate depths in various ocean locations and in many coastal areas. Over the last half century, major oxygen losses have occurred in inland seas, estuaries, and in the coastal and open ocean (*high confidence*). Ocean oxygen levels are projected to decrease by as much as 3.5% under the RCP8.5 scenario by 2100 relative to preindustrial values (*high confidence*).

13.0 A Changing Ocean

Anthropogenic perturbations to the global Earth system have included important alterations in the nutrient composition, temperature, and circulation of the oceans. Some of these changes will be distinguishable from the background natural variability in nearly half of the global open ocean within a decade, with important consequences for marine ecosystems and their services (Gattuso et al. 2015). However, the timeframe for detection will vary depending on the parameter featured (Henson et al 2010; Henson et al 2016).

13.1 Ocean Warming

13.1.1 General Background

Approximately 93% of excess heat energy trapped since the 1970s has been absorbed into the oceans, lessening atmospheric warming and leading to a variety of changes in ocean conditions, including sea level rise and ocean circulation (see Ch. 2: Physical Drivers of Climate Change, Ch. 6: Temperature Change, and Ch. 12: Sea Level Rise in this report; Rhein et al. 2013; Gattuso et al. 2015). This is the result of the high heat capacity of seawater relative to the atmosphere, the relative area of the ocean compared to the land, and the ocean circulation that enables the transport of heat into deep waters. This large heat absorption by the oceans moderates the effects of increased anthropogenic greenhouse emissions on terrestrial climates while altering the fundamental physical properties of the ocean and indirectly impacting chemical properties such as the biological pump through increased stratification (Gattuso et al. 2015; Rossby 1959). Although upper ocean temperature varies over short- and medium timescales (for example, seasonal and regional patterns), there are clear long-term increases in surface temperature and ocean heat content over the past 65 years (Cheng et al. 2017; Rhein et al. 2013; Levitus et al. 2012).

13.1.2 Ocean Heat Content

Ocean heat content (OHC) is an ideal variable to monitor changing climate as it is calculated using the entire water column, so ocean warming can be documented and compared between particular regions, ocean basins, and depths. However, for years prior to the 1970s, estimates of ocean uptake are confined to the upper ocean due to sparse spatial and temporal coverage and limited vertical capabilities of many of the instruments in use. Ocean heat content estimates are improved for time periods after 1970 with increased sampling coverage and depth (Abraham et al. 2013; Rhein et al. 2013). Estimates of OHC have been calculated going back to the 1950s using averages over longer time intervals (i.e., decadal or 5-year intervals) to compensate for sparse data distributions, allowing for clear long-term trends to emerge (e.g., Levitus et al. 2012).

From 1960 to 2015, ocean heat content (OHC) significantly increased for both 0–700 and 700–2,000 m depths, for a total ocean warming of $33.5 \pm 7.0 \times 10^{22}$ J (a net heating of 0.37 ± 0.08 W/m²), although there is some uncertainty with global ocean heat estimates (Figure 13.1; Cheng

et al. 2017). During this period, there is evidence of an acceleration of ocean warming beginning in 1998 (Lee et al. 2015), with a total heat increase of about 15.2×10^{22} J (Cheng et al. 2017). Robust ocean warming occurs in the upper 700 m and is slow to penetrate into the deep ocean. However, the 700–2,000 m depths constitute an increasing portion of the total ocean energy budget as compared to the surface ocean (Figure 13.1; Cheng et al. 2017). The role of the deep ocean (below 2,000 m [6,600 ft]) in ocean heat uptake remains uncertain, both in the magnitude but also the sign of the uptake (Purkey and Johnson 2010; Llovel et al. 2014). Penetration of surface waters to the deep ocean is a slow process, which means that while it takes only about a decade for near-surface temperatures to respond to increased heat energy, the deep ocean will continue to warm, and as a result sea levels will rise for centuries to millennia even if all further emissions cease (Rhein et al. 2013).

[INSERT FIGURE 13.1 HERE]

Several sources have documented warming in all ocean basins from 0–2,000 m depths over the past 50 years (Figure 13.2; Boyer et al. 2016; Cheng et al. 2017; Levitus et al. 2012). Annual fluctuations in surface temperatures and OHC are attributed to the combination of a long-term secular trend and decadal and smaller time scale variations, such as the Pacific Decadal Oscillation (PDO) and the Atlantic Multidecadal Oscillation (AMO) (Ch. 5: Large-Scale Circulation and Climate Variability; Ch. 12: Sea Level Rise; Trenberth et al. 2014; Steinman et al. 2015). The transport of heat to the deep ocean is likely linked to the strength of the Atlantic Meridional Overturning Circulation (see Section 13.2.1), where the Atlantic and Southern Ocean accounts for the dominant portion of total OHC at the 700–2,000 m depth (Figure 13.2; Cheng et al. 2017; Lee et al. 2015; Roemmich et al. 2015; Abraham et al. 2013). Decadal variability in ocean heat uptake is mostly attributed to ENSO phases (with El Niños warming and La Niñas cooling). For instance, La Niña conditions over the past decade have led to colder ocean temperatures in the eastern tropical Pacific (Abraham et al. 2013; Cheng et al. 2017; Lee et al. 2015; Kosaka and Xie 2013). For the Pacific and Indian Oceans, the decadal shifts are primarily observed in the upper 350 m depth, likely due to shallow subtropical circulation, leading to an abrupt increase of OHC in the Indian Ocean carried by the Indonesian throughflow from the Pacific Ocean over the last decade (Lee et al. 2015). Although there is natural variability in ocean temperature, there remain clear increasing trends due to anthropogenic influences.

[INSERT FIGURE 13.2 HERE]

13.1.3 Sea Surface Temperature and U.S. Regional Warming

In addition to OHC, sea surface temperature (SST) measurements are widely available. SST measurements are useful because 1) the measurements have been taken over 150 years (albeit using different platforms, instruments, and depths through time); 2) SST reflects the lower boundary condition of the atmosphere; and 3) SST can be used to predict specific regional impacts of global warming on terrestrial and coastal systems (Roemmich et al. 2015; Yan et al.

2016; Matthews 2013). Globally, surface ocean temperatures have increased by $1.3^{\circ} \pm 0.1^{\circ}\text{F}$ ($0.70^{\circ} \pm 0.08^{\circ}\text{C}$) per century from 1900 to 2016 for the Extended Reconstructed Sea Surface Temperature version 4 (ERSST v4) record (Huang et al. 2015). All U.S. coastal waters have warmed by more than 0.7°F (0.4°C) over this period as shown in both Table 13.1 and Chapter 6: Temperature Change, Figure 6.6. During the past century, the rates of increase of SSTs for the coastal waters of three U.S. regions were above the global average rate. These included the waters around Alaska, the Caribbean, and the Southwest (Table 13.1). Over the last decade, some regions have experienced increased high ocean temperature anomalies. For instance, due to a resilient ridge of high pressure over the North American west coast, storm activity and mixing was suppressed and heat in the upper ocean intensified in 2013 in a phenomenon known as “The Blob” (Bond et al. 2015). These anomalously warm waters persisted in the coastal waters of the Alaskan and Pacific Northwest until 2015. Under a higher emissions pathway (RCP8.5), ocean SST are projected to increase by an additional 4.9°F (2.7°C) by 2100 (Figure 13.3), whereas for a lower emissions scenario (RCP4.5) the SST increase would be 2.3°F (1.3°C ; Bopp et al. 2013). In all U.S. coastal regions, the warming since 1901 is detectable compared to natural variability and attributable to anthropogenic forcing, according to an analysis of the CMIP5 models (Ch. 6: Temperature Change, Figure 6.5).

[INSERT TABLE 13.1 AND FIGURE 13.3 HERE]

13.1.4 Ocean Heat Feedback

The residual heat not taken up by the oceans increases land surface temperatures (approximately 3%) and atmospheric temperatures (approximately 1%), and melts both land and sea ice (approximately 3%), leading to sea level rise (see Ch. 12: Sea Level Rise; Cheng et al. 2017; Rhein et al. 2013; Nieves et al. 2015). The meltwater from land and sea ice amplifies further subsurface ocean warming and ice shelf melting, primarily due to increased thermal stratification, which reduces the ocean’s efficiency in transporting heat to deep waters (Rhein et al. 2013). Surface ocean stratification has increased by about 4% during the period 1971–2010 (Ciais et al. 2013) due to thermal heating and freshening from increased freshwater inputs (precipitation and evaporation changes and land and sea ice melting). The increase of ocean stratification will contribute to further feedback of ocean warming and, indirectly, mean sea level. In addition, increases in stratification are associated with suppression of tropical cyclone intensification (Mei et al. 2015), retreat of the polar ice sheets (Straneo and Heimbach 2013), and reductions of the convective mixing at higher latitudes that transports heat to the deep ocean through the Atlantic Meridional Overturning Circulation (AMOC) (Rahmstorf et al. 2015). Ocean heat uptake therefore represents an important feedback that will have a significant influence on future shifts in climate (see Ch. 2: Physical Drivers of Climate Change).

13.2 Ocean Circulation

13.2.1 Atlantic Meridional Overturning Circulation

The Atlantic Meridional Overturning Circulation (AMOC) refers to the three-dimensional, time-dependent circulation of the Atlantic Ocean, which has been a high priority topic of study in recent decades. The AMOC plays an important role in climate through its transport of heat, freshwater, and carbon (e.g., Johns et al. 2011; McDonagh et al. 2015; Talley et al. 2016). AMOC-associated poleward heat transport substantially contributes to North American and continental European climate (see Ch. 5: Circulation and Variability). The Gulf Stream, in contrast to other western boundary currents, is expected to slow down because of the weakening of the AMOC, which would impact the European climate (Yang et al. 2016). Variability in the AMOC has been attributed to wind forcing on intra-annual time scales and to geostrophic forces on interannual to decadal timescales (Buckley and Marshall 2016). Increased freshwater fluxes from melting arctic sea and land ice can weaken open ocean convection and deep-water formation in the Labrador and Irminger Seas, which could weaken the AMOC (Ch. 11: Arctic Changes; Rahmstorf et al. 2015; Yang et al. 2016; Also see Chapter 5, Section 5.2.3: North Atlantic Oscillation and Northern Annular Mode).

While one recent study has suggested that the AMOC has slowed since preindustrial times (Rahmstorf et al. 2015) and another suggested slowing on faster time scales (Smeed et al. 2014), there is at present insufficient observational evidence to support a finding of long term slowdown of AMOC strength over the 20th century (Rhein et al. 2013) or within the last 50 years (Buckley and Marshall 2016) as decadal ocean variability can obscure long-term trends. Some studies show long-term trends (Longworth et al. 2011; Bryden et al. 2005), but the combination of sparse data and large seasonal variability may also lead to incorrect interpretations (e.g., Kanzow et al. 2010). Several recent high resolution modeling studies constrained with the limited existing observational data (Jackson et al. 2016) and/or with reconstructed freshwater fluxes (Böning et al. 2016) suggest that the recently observed AMOC slowdown at 26°N (off the Florida coast) since 2004 (e.g., as described in Smeed et al. 2014) is mainly due to natural variability, and that anthropogenic forcing has not yet caused a significant AMOC slowdown. In addition, direct observations of the AMOC in the South Atlantic fail to unambiguously demonstrate anthropogenic trends (e.g., Dong et al. 2015; Garzoli et al. 2013).

Under a high emissions future scenario (RCP8.5) in CMIP5 simulations, it is very likely that the AMOC will weaken over the 21st century. The projected decline ranges from 12% to 54% (Collins et al. 2013), with the range width reflecting substantial uncertainty in quantitative projections of AMOC behavior. In RCP4.5 scenarios, CMIP5 models predict a 20% weakening of the AMOC during the first half of the 21st century and a stabilization and slight recovery after that (Cheng et al. 2013). The projected slowdown of the AMOC will be counteracted by the warming of the deep ocean (below 700 m [2,300 ft]), which will tend to strengthen the AMOC (Patara and Böning 2014). The situation is further complicated due to the known bias in coupled

climate models related to the direction of the salinity transport in models versus observations, which is an indicator of AMOC stability (e.g., Drijhout et al. 2011; Bryden et al. 2011; Garzoli et al. 2013). Some argue that coupled climate models should be corrected for this known bias and that AMOC variations could be even larger than the gradual decrease most models predict if the AMOC were to shut down completely and “flip states” (Liu et al. 2017). Any AMOC slowdown will result in less heat and CO₂ absorbed by the ocean from the atmosphere, which is a positive feedback to climate change (also see Ch. 2: Physical Drivers of Climate Change).

13.2.2 Changes in Salinity Structure

As a response to warming, increased atmospheric moisture leads to stronger evaporation or precipitation in terrestrial and oceanic environments and melting of land and sea ice. Approximately 80% of precipitation/evaporation events occur over the ocean, leading to patterns of higher salt content or freshwater anomalies and changes in ocean circulation (see Ch. 2: Physical Drivers of Climate Change and Ch. 6: Temperature Change; Durack and Wijffels 2010). Over 1950–2010, average global amplification of the surface salinity pattern amounted to 5.3%; where fresh regions in the ocean became fresher and salty regions became saltier (Skliris et al. 2014). However, the long-term trends of these physical and chemical changes to the ocean are difficult to isolate from natural large-scale variability. In particular, ENSO displays particular salinity and precipitation/evaporation patterns that skew the trends. More research and data are necessary to better model changes to ocean salinity. Several models have shown a similar spatial structure of surface salinity changes, including general salinity increases in the subtropical gyres, a strong basin-wide salinity increase in the Atlantic Ocean, and reduced salinity in the western Pacific warm pools and the North Pacific subpolar regions (Durack and Wijffels 2010; Skliris et al. 2014). There is also a stronger distinction between the upper salty thermocline and fresh intermediate depth through the century. The regional changes in salinity to ocean basins will have an overall impact on ocean circulation and net primary production, leading to corresponding carbon export (see Ch. 2: Physical Drivers of Climate Change). In particular, the freshening of the Arctic Ocean due to melting of land and sea ice can lead to buoyancy changes which could slow down the AMOC (see Section 13.2.1).

13.2.3 Changes in Upwelling

Significant changes to ocean stratification and circulation can also be observed regionally, along the eastern ocean boundaries and at the equator. In these areas, wind-driven upwelling brings colder, nutrient- and carbon-rich water to the surface; this upwelled water is more efficient in heat and CO₂ uptake. There is some evidence that coastal upwelling in mid- to high-latitude eastern boundary regions has increased in intensity and/or frequency (García-Reyes et al. 2015), but in more tropical areas of the western Atlantic, such as in the Caribbean Sea, it has decreased between 1990 and 2010 (Taylor et al. 2012; Astor et al. 2013). This has led to a decrease in primary productivity in the southern Caribbean Sea (Taylor et al. 2012). Within the continental United States, the California Current is experiencing fewer (by about 23%–40%) but stronger

upwelling events (Hoegh-Guldberg et al. 2014; Sydeman et al. 2014; Jacox et al. 2014). Stronger offshore upwelling combined with cross-shelf advection brings nutrients from the deeper ocean but also increased offshore transport (Bakun et al. 2015). The net nutrient load in the coastal regions is responsible for increased productivity and ecosystem function.

IPCC 2013 concluded that there is low confidence in the current understanding of how eastern upwelling systems will be altered under future climate change because of the obscuring role of multidecadal climate variability (Ciais et al. 2013). However, subsequent studies show that by 2100, upwelling is predicted to start earlier, end later, and intensify in three of the four major eastern boundary upwelling systems (not in the California Current; Wang et al. 2015). Southern Ocean upwelling will intensify while the Atlantic equatorial upwelling systems will weaken (Hoegh-Guldberg et al. 2014; Wang et al. 2015). The intensification is attributed to the strengthening of regional coastal winds as observations already show (Sydeman et al. 2014), and RCP8.5 model projection scenarios estimate wind intensifying near poleward boundaries (including northern California current system [CCS]) and weakening near equatorward boundaries (including southern CCS) for the 21st century (Rykaczewski et al. 2015; Wang et al. 2015).

13.3 Ocean Acidification

13.3.1 General Background

In addition to causing changes in climate, increasing atmospheric levels of carbon dioxide (CO_2) from the burning of fossil fuels and other human activities, including changes in land use, have a direct effect on ocean chemistry (Orr et al. 2005; Feely et al. 2009). Ocean acidification refers to a change in ocean chemistry in response to the uptake of increasing CO_2 in the atmosphere.

Ocean acidification causes a variety of chemical changes in seawater: an increase in the partial pressure of CO_2 ($p\text{CO}_{2,\text{sw}}$), dissolved inorganic carbon (DIC), and an increase in the concentration of hydrogen and bicarbonate ions and a decrease in the concentration of carbonate ions (Figure 13.4). In brief, CO_2 is an acid gas that combines with water to form carbonic acid, which then dissociates to hydrogen and bicarbonate ions. Increasing concentrations of seawater hydrogen ions result in a decrease of carbonate ions through their conversion to bicarbonate ions. Ocean acidity refers to the concentration of hydrogen ions in ocean seawater regardless of ocean pH, which is fundamentally basic (e.g., $\text{pH} > 7$). Ocean surface waters have become 30% more acidic over the last 150 years as they have absorbed large amounts of CO_2 from the atmosphere (Feely et al. 2004), and anthropogenically sourced CO_2 is gradually invading into oceanic deep waters. Since the preindustrial period, the oceans have absorbed approximately 27% of all CO_2 emitted to the atmosphere. Oceans currently absorb about 26% of the human-caused CO_2 anthropogenically emitted into the atmosphere (Le Quéré et al. 2016).

[INSERT FIGURE 13.4 HERE]

13.3.2 Open Ocean Acidification

Surface waters in the open ocean experience changes in carbonate chemistry reflective of large-scale physical oceanic processes (see Ch. 2: Physical Drivers of Climate Change). These processes include both the global uptake of atmospheric CO₂ and the shoaling of naturally acidified subsurface waters due to vertical mixing and upwelling. In general, the rate of ocean acidification in open ocean surface waters at a decadal time-scale closely approximates the rate of atmospheric CO₂ increase (Bates et al. 2014). Large, multidecadal phenomena such as the Atlantic Multidecadal Oscillation and Pacific Decadal Oscillation can add variability to the observed rate of change (Bates et al. 2014).

13.3.3 Coastal Acidification

Coastal shelf and nearshore waters are influenced by the same processes as open ocean surface waters such as absorption of atmospheric CO₂ and upwelling, as well as a number of additional, local-level processes, including freshwater and nutrient input (Duarte et al. 2013). Coastal acidification generally exhibits higher-frequency variability and short-term episodic events relative to open-ocean acidification (Borges and Gypens 2010; Waldbusser and Salisbury 2014; Hendriks et al. 2015; Sutton et al. 2016). Upwelling is of particular importance in coastal waters, especially along the Pacific Coast. Deep waters that shoal with upwelling are enriched in CO₂ due to uptake of anthropogenic atmospheric CO₂ when last in contact with the atmosphere, coupled with deep water respiration processes and lack of gas exchange with the atmosphere (Feely et al. 2009; Harris et al. 2013). Freshwater inputs to coastal waters change seawater chemistry in ways that make it more susceptible to acidification, largely by freshening ocean waters and contributing varying amounts of dissolved inorganic carbon (DIC), total alkalinity (TA), dissolved and particulate organic carbon, and nutrients from riverine and estuarine sources. Coastal waters of the East Coast and mid-Atlantic are far more influenced by freshwater inputs than are Pacific Coast waters (Gledhill et al. 2015). Coastal waters can episodically experience riverine and glacial melt plumes that create conditions in which seawater can dissolve calcium carbonate structures (Evans et al. 2014; Salisbury et al. 2008). While these processes have persisted historically, climate-induced increases in glacial melt and high-intensity precipitation events can yield larger freshwater plumes than have occurred in the past. Nutrient runoff can increase coastal acidification by creating conditions that enhance biological respiration. In brief, nutrient loading typically promotes phytoplankton blooms, which, when they die, are consumed by bacteria. Bacteria respire CO₂ and thus bacterial blooms can result in acidification events whose intensity depends on local hydrographic conditions, including water column stratification and residence time (Waldbusser and Salisbury 2014). Long-term changes in nutrient loading, precipitation, and/or ice melt may also impart long-term, secular changes in the magnitude of coastal acidification.

13.3.4 Latitudinal Variation

Ocean carbon chemistry is highly influenced by water temperature, largely because the solubility of CO₂ in seawater increases as water temperature declines. Thus, cold, high-latitude waters can absorb more CO₂ than warm, lower-latitude waters (Gledhill et al. 2015; Bates and Mathis 2009). Because carbonate minerals also more readily dissolve in colder waters, these waters can more regularly become undersaturated with respect to calcium carbonate whereby mineral dissolution is energetically favored. This chemical state, often referred to as seawater being “corrosive” to calcium carbonate, is important when considering the ecological implications of ocean acidification as many species make structures such as shells and skeletons from calcium carbonate. Some high-latitude waters already experience such corrosive conditions, which are rarely documented in low-latitude systems. For example, corrosive conditions have been documented in the Arctic and northeastern Pacific Oceans (Bates and Mathis 2009; Feely et al. 2008; Qi et al. 2017; Sutton et al. 2016). It is important to note that low-latitude waters are experiencing a greater absolute rate of change in calcium carbonate saturation state than higher latitudes, though these low-latitude waters are not approaching the undersaturated state except within near-shore or some benthic habitats (Friedrich et al. 2012).

13.3.5 Paleo Evidence

Evidence suggests that the current rate of ocean acidification is the fastest in the last 66 million years (the K-Pg boundary) and possibly even the last 300 million years (when the first pelagic calcifiers evolved providing proxy information and also a strong carbonate buffer, characteristic of the modern ocean) (Hönisch et al. 2012; Zeebe et al. 2016). The Paleo-Eocene Thermal Maximum (PETM; around 56 million years ago) is often referenced as the closest analogue to the present, although the overall rate of change in CO₂ conditions during that event (estimated between 0.6 and 1.1 GtC/year) was much lower than the current increase in atmospheric CO₂ of 10 GtC/year (Wright and Schaller 2013; Zeebe et al. 2016). The relatively slower rate of atmospheric CO₂ increase at the PETM likely led to relatively small changes in carbonate ion concentration in seawater compared with the contemporary acidification rate, due to the ability of rock weathering to buffer the change over the longer time period (Zeebe et al. 2016). Some of the presumed acidification events in Earth’s history have been linked to selective extinction events suggestive of how guilds of species may respond to the current acidification event (Hönisch et al. 2012).

13.3.6 Projected Changes

Projections indicate that by the end of the century under higher emissions pathways, such as SRES A1fI or RCP8.5, open-ocean surface pH will decline from the current average level of 8.1 to a possible average of 7.8 (Figure 13.5; Gattuso et al. 2015). When the entire ocean volume is considered under the same scenario, the volume of waters undersaturated with respect to calcium carbonate could expand from 76% in the 1990s to 91% in 2100. As discussed above, for a

variety of reasons, not all ocean and coastal regions will experience acidification in the same way depending on other compounding factors. For instance, recent observational data from the Arctic Basin show that the Beaufort Sea became undersaturated, for part of the year, with respect to aragonite in 2001, while other continental shelf seas in the Arctic Basin are projected to do so closer to the middle of the century (e.g., the Chukchi Sea in about 2033 and Bering Sea in about 2062; Mathis et al. 2015). Deviation from the global average rate of acidification will be especially true in coastal and estuarine areas where the rate of acidification is influenced by other drivers than atmospheric CO₂, some of which are under the control of local management decisions (for example, nutrient pollution loads).

[INSERT FIGURE 13.5 HERE]

13.4 Ocean Deoxygenation

13.4.1 General Background

Oxygen is essential to most life in the ocean, governing a host of biogeochemical and biological processes. Oxygen influences metabolic, physiological, reproductive, behavioral, and ecological processes, ultimately shaping the composition, diversity, abundance, and distribution of organisms from microbes to whales. Increasingly, climate-induced oxygen loss (deoxygenation) associated with ocean warming and reduced ventilation to deep waters has become evident locally, regionally, and globally. Deoxygenation can also be attributed to anthropogenic nutrient input, especially in the coastal regions, where the nutrients can lead to the proliferation of primary production and, consequently, enhanced drawdown of dissolved oxygen by microbes (Altieri and Gedan 2015). In addition, acidification (Section 13.2) can co-occur with deoxygenation as a result of warming-enhanced biological respiration (Breitburg et al. 2015). As aerobic organisms respire, O₂ is consumed and CO₂ is produced. Understanding the combined effect of both low O₂ and low pH on marine ecosystems is an area of active research (Gobler et al. 2014). Warming also raises biological metabolic rates which, in combination with intensified coastal and estuarine stratification, exacerbates eutrophication-induced hypoxia. We now see earlier onset and longer periods of seasonal hypoxia in many eutrophic sites, most of which occur in areas that are also warming (Altieri and Gedan 2015).

13.4.2 Climate Drivers of Ocean Deoxygenation

Global ocean deoxygenation is a direct effect of warming. Ocean warming reduces the solubility of oxygen (that is, warmer water can hold less oxygen) and changes physical mixing (for example, upwelling and circulation) of oxygen in the oceans. The increased temperature of global oceans accounts for about 15% of current global oxygen loss (Helm et al. 2011), although changes in temperature and oxygen are not uniform throughout the ocean (Roemmich et al. 2015). Warming also exerts direct influence on thermal stratification and enhances salinity stratification through ice melt and climate change-associated precipitation effects. Intensified

1 stratification leads to reduced ventilation (mixing of oxygen into the ocean interior) and accounts
2 for up to 85% of global ocean oxygen loss (Helm et al. 2011). Effects of ocean temperature
3 change and stratification on oxygen loss are strongest in intermediate or mode waters at bathyal
4 depths (in general, 200–3,000 m) and also nearshore and in the open ocean; these changes are
5 especially evident in tropical and subtropical waters globally, in the Eastern Pacific (Stramma et
6 al. 2010), and in the Southern Ocean (Helm et al. 2011).

7 There are also other, less direct effects of global temperature increase. Warming on land reduces
8 terrestrial plant water efficiency (through effects on stomata; see Ch. 8: Drought, Floods, and
9 Wildfires, Key Message 3), leading to greater runoff, on average, into coastal zones (see Ch. 8:
10 Drought, Floods, and Wildfires for other hydrological effects of warming) and further enhancing
11 hypoxia potential because greater runoff means more nutrient transport (See Ch. 2: Physical
12 Drivers of Climate Change; Reay et al. 2008; Rabalais et al. 2009). Estuaries, especially ones
13 with minimal tidal mixing, are particularly vulnerable to oxygen-depleted dead zones from the
14 enhanced runoff and stratification. Warming can induce dissociation of frozen methane in gas
15 hydrates buried on continental margins, leading to further drawdown of oxygen through aerobic
16 methane oxidation in the water column (Boetius and Wenzhöfer 2013). On eastern ocean
17 boundaries, warming can enhance the land–sea temperature differential, causing increased
18 upwelling due to higher winds with (a) greater nutrient input leading to production, sinking,
19 decay, and biochemical drawdown of oxygen and (b) upwelling of naturally low-oxygen, high-
20 CO₂ waters onto the upper slope and shelf environments (Sydeman et al. 2014; Feely et al.
21 2009). However, in the California Current Ecosystem, upwelling intensification has occurred
22 only in the poleward regions (north of San Francisco), and the drivers may not be associated with
23 land–sea temperature differences (Rykaczewski et al. 2015). Taken together, the effects of
24 warming are manifested as low-oxygen water in open oceans are transported to and upwelled
25 along coastal regions. These low-oxygen upwelled waters are then coupled with eutrophication-
26 induced hypoxia, further reducing oxygen content in coastal areas.

27 Changes in precipitation, winds, circulation, airborne nutrients, and sea level can also contribute
28 to ocean deoxygenation. Projected increases in precipitation in some regions will intensify
29 stratification, reducing vertical mixing and ventilation, and intensify nutrient input to coastal
30 waters through excess runoff, which leads to increased algal biomass and concurrent dissolved
31 oxygen consumption via community respiration (Lee et al. 2016). Coastal wetlands that might
32 remove these nutrients before they reach the ocean may be lost through rising sea level, further
33 exacerbating hypoxia (Rabalais et al. 2009). Some observations of oxygen decline are linked to
34 regional changes in circulation involving low-oxygen water masses. Enhanced fluxes of airborne
35 iron and nitrogen are interacting with natural climate variability and contributing to fertilization,
36 enhanced respiration, and oxygen loss in the tropical Pacific (Ito et al. 2016). In contrast to the
37 many sources of climate-induced oxygen loss, the projected increase in incidence and intensity
38 of cyclones and hurricanes will induce mixing, which can ameliorate hypoxia locally (Rabalais et
39 al. 2009).

13.4.3 Biogeochemical Feedbacks of Deoxygenation to Climate and Elemental Cycles

Climate patterns and ocean circulation have a large effect on global nitrogen and oxygen cycles, which in turn affect phosphorus and trace metal availability and generate feedbacks to the atmosphere and oceanic production. Global ocean productivity may be affected by climate-driven changes below the tropical and subtropical thermocline which control the volume of suboxic waters (< 5 micromolar O_2), and consequently the loss of fixed nitrogen through denitrification (Codispoti et al. 2001; Deutsch et al. 2011). The extent of suboxia in the open ocean also regulates the production of the greenhouse gas nitrous oxide (N_2O); as oxygen declines, greater N_2O production may intensify global warming, as N_2O is about 310 times more effective at trapping heat than CO_2 (see Ch. 2: Physical Drivers of Climate Change, Section 2.3.2; Gruber 2008; EPA 2017). Production of hydrogen sulfide (H_2S , which is highly toxic) and intensified phosphorus recycling can occur at low oxygen levels (Wallmann 2003). Other feedbacks may emerge as oxygen minimum zone (OMZ) shoaling diminishes the depths of diurnal vertical migrations by fish and invertebrates, and as their huge biomass and associated oxygen consumption deplete oxygen (Bianchi et al. 2013).

13.4.4 Past Trends

Over hundreds of millions of years, oxygen has varied dramatically in the atmosphere and ocean and has been linked to biodiversity gains and losses (Knoll and Carroll 1999; McFall-Ngai et al. 2013). Variation in oxygenation in the paleo record is very sensitive to climate—with clear links to temperature and often CO_2 variation (Falkowski et al. 2011). OMZs expand and contract in synchrony with warming and cooling events, respectively (Robinson et al. 2007). Episodic climate events that involve rapid temperature increases over decades, followed by a cool period lasting a few hundred years, lead to major fluctuations in the intensity of Pacific and Indian Ocean OMZs (i.e., DO of $< 20 \mu M$). These events are associated with rapid variations in North Atlantic deep water formation (Schmittner et al. 2007). Ocean oxygen fluctuates on glacial-interglacial timescales of thousands of years in the Eastern Pacific (Galbraith et al. 2004; Moffitt et al. 2015).

13.4.5 Modern Observations (last 50+ years)

Long-term oxygen records made over the last 50 years reflect oxygen declines in inland seas (Justić et al. 1987; Zaitzev 1992; Conley et al. 2011), in estuaries (Brush 2009; Gilbert et al. 2005), and in coastal waters (Rabalais et al. 2007, 2010; Booth et al. 2012; Baden et al. 1990). The number of coastal, eutrophication-induced hypoxic sites in the United States has grown dramatically over the past 40 years (Diaz and Rosenberg 2008). Over larger scales, global syntheses show hypoxic waters have expanded by 4.5 million km^2 at a depth of 200 m (Stramma et al. 2010), with widespread loss of oxygen in the Southern Ocean (Helm et al. 2011), Western Pacific (Takatani et al. 2012), and North Atlantic (Stendardo and Gruber 2012). Overall oxygen declines have been greater in coastal ocean than in the open ocean (Gilbert et al. 2010) and often

greater inshore than offshore (Bograd et al. 2015). The emergence of a deoxygenation signal in regions with naturally high oxygen variability will unfold over longer time periods (20–50 years from now) (Long et al. 2016).

13.4.6 Projected Changes

GLOBAL MODELS

Global models generally agree that ocean deoxygenation is occurring; this finding is also reflected in in situ observations from past 50 years. Compilations of 10 Earth System models predict a global average loss of oxygen of –3.5% (RCP8.5) to –2.4% (RCP4.5) by 2100, but much stronger losses regionally, and in intermediate and mode waters (Bopp et al. 2013) (Figure 13.6). The North Pacific, North Atlantic, Southern Ocean, subtropical South Pacific, and South Indian Oceans all are expected to experience deoxygenation, with O₂ decreases of as much as 17% in the North Pacific by 2100 for the RCP8.5 pathway. However, the tropical Atlantic and tropical Indian Oceans show increasing O₂ concentrations. In the many areas where oxygen is declining, high natural variability makes it difficult to identify anthropogenically forced trends (Long et al. 2016).

[INSERT FIGURE 13.6 HERE]

REGIONAL MODELS

Regional models are critical because many oxygen drivers are local, influenced by bathymetry, winds, circulation, and fresh water and nutrient inputs. Most eastern boundary upwelling areas are predicted to experience intensified upwelling to 2100 (Wang et al. 2015), although on the West Coast projections for increasing upwelling for the northern California Current occur only north of San Francisco (see Section 13.2.3).

Particularly notable for the western United States, variation in trade winds in the eastern Pacific Ocean can affect nutrient inputs, leading to centennial periods of oxygen decline or oxygen increase distinct from global oxygen decline (Deutsch et al. 2014). Oxygen dynamics in the Eastern Tropical Pacific are highly sensitive to equatorial circulation changes (Montes et al. 2014).

Regional modeling also shows that year-to-year variability in precipitation in the central United States affects the nitrate–N flux by the Mississippi River and the extent of hypoxia in the Gulf of Mexico (Donner and Scavia 2007). A host of climate influences linked to warming and increased precipitation are predicted to lower dissolved oxygen in Chesapeake Bay (Najjar et al. 2010).

13.5 Other Coastal Changes

13.5.1 Sea Level Rise

Sea level is an important variable that affects coastal ecosystems. Global sea level rose very rapidly at the end of the last glaciation, as glaciers and the polar ice sheets thinned and melted at their fringes. On average around the globe, sea level is estimated to have risen at rates exceeding 2.5 mm/year between about 8,000 and 6,000 years before present. These rates steadily decreased to less than 2.0 mm/year through about 4,000 years ago and stabilized at less than 0.4 mm/year through the late 1800s. Global sea level rise has accelerated again within the last 100 years, and now averages about 1 to 2 mm/year (Thompson et al. 2016). See Chapter 12: Sea Level Rise for more thorough analysis of how sea level rise has already and will affect the U.S. coasts.

13.5.2 Wet and Dry Deposition

Dust transported from continental desert regions to the marine environment deposits nutrients such as iron, nitrogen, phosphorus, and trace metals that stimulate growth of phytoplankton and increase marine productivity (Jickells and Moore 2015). U.S. continental and coastal regions experience large dust deposition fluxes originating from the Saharan desert to the East and from Central Asia and China to the Northwest (Chiapello 2014). Changes in drought frequency or intensity resulting from anthropogenically forced climate change, as well as other anthropogenic activities such as agricultural practices and land-use changes may play an important role in the future viability and strength of these dust sources (e.g., Mulitza et al. 2010).

Additionally, oxidized nitrogen, released during high-temperature combustion over land, and reduced nitrogen, released from intensive agriculture, are emitted in high population areas in North America and are carried away and deposited through wet or dry deposition over coastal and open ocean ecosystems via local wind circulation. Wet deposition of pollutants produced in urban areas is known to play an important role in changes of ecosystem structure in coastal and open ocean systems through intermediate changes in the biogeochemistry, for instance in dissolved oxygen or various forms of carbon (Paerl et al. 2002).

13.5.3 Primary Productivity

Marine phytoplankton represent about half of the global net primary production (NPP) (approximately 50 ± 28 GtC/year), fixing atmospheric CO₂ into a bioavailable form for utilization by higher trophic levels (see also Ch. 2: Physical Drivers of Climate Change; Carr et al. 2006; Franz et al. 2016). As such, NPP represents a critical component in the role of the oceans in climate feedback. The effect of climate change on primary productivity varies across the coasts depending on local conditions. For instance, nutrients that stimulate phytoplankton growth are impacted by various climate conditions, such as increased stratification which limits the transport of nutrient-rich deep water to the surface, changes in circulation leading to variability in dry and wet deposition of nutrients to coasts, and altered precipitation/evaporation

1 which changes runoff of nutrients from coastal communities. The effect of the multiple physical
2 factors on NPP is complex and leads to model uncertainties (Chavez et al. 2011). There is
3 considerable variation in model projections for NPP, from estimated decreases or no changes, to
4 the potential increase by 2100 (Frölicher et al. 2016; Fu et al. 2016; Laufkötter et al. 2015).
5 Simulations from nine Earth system models projected total NPP in 2090 to decrease by 2%–16%
6 and export production (that is, particulate flux to the deep ocean) to drop by 7%–18% as
7 compared to 1990 (RCP8.5; Fu et al. 2016). More information on phytoplankton species
8 response and associated ecosystem dynamics is needed as any reduction of NPP would have a
9 strong impact on atmospheric CO₂ levels and marine ecosystems in general.

10 **13.5.4 Estuaries**

11 Estuaries are critical ecosystems of biological, economic, and social importance in the United
12 States. They are highly dynamic, influenced by the interactions of atmospheric, freshwater,
13 terrestrial, oceanic, and benthic components. Of the 28 national estuarine research reserves in the
14 United States and Puerto Rico, all are being impacted by climate change to varying levels
15 (Robinson et al. 2013). In particular, sea level rise, saltwater intrusion, and the degree of
16 freshwater discharge influence the forces and processes within these estuaries (Monbaliu et al.
17 2014). Sea level rise and subsidence are leading to drowning of existing salt marshes and/or
18 subsequent changes in the relative area of the marsh plain, if adaptive upslope movement is
19 impeded due to urbanization along shorelines. Several model scenarios indicate a decline in salt
20 marsh habitat quality and an accelerated degradation as the rate of sea level rise increases in the
21 latter half of the century (Schile et al. 2014; Swanson et al. 2015). The increase in sea level as
22 well as alterations to oceanic and atmospheric circulation can result in extreme wave conditions
23 and storm surges, impacting coastal communities (Robinson et al. 2013). Additional climate
24 change impacts to the physical and chemical estuarine processes include more extreme sea
25 surface temperatures (higher highs and lower lows compared to the open ocean due to shallower
26 depths and influence from land temperatures), changes in flow rates due to changes in
27 precipitation, and potentially greater extents of salinity intrusion.

TRACEABLE ACCOUNTS

Key Finding 1

The world's oceans have absorbed about 93% of the excess heat caused by greenhouse gas warming since the mid-20th century, making them warmer and altering global and regional climate feedbacks. Ocean heat content has increased at all depths since the 1960s and surface waters have warmed by about $1.3^{\circ} \pm 0.1^{\circ}\text{F}$ ($0.7^{\circ} \pm 0.08^{\circ}\text{C}$) per century globally since 1900 to 2016. Under a high emissions scenario, a global increase in average sea surface temperature of $4.9^{\circ} \pm 1.3^{\circ}\text{F}$ ($2.7^{\circ} \pm 0.7^{\circ}\text{C}$) by 2100 is projected, with even higher changes in some U.S. coastal regions. (*Very high confidence*)

Description of evidence base

The key finding and supporting text summarizes the evidence documented in climate science literature, including Rhein et al. 2013 and thereafter. Oceanic warming has been documented in a variety of data sources, most notably the WOCE (<http://www.nodc.noaa.gov/woce/wdiu/>), ARGO database (<https://www.nodc.noaa.gov/argo/>), and ERSSTv4 (<https://www.ncdc.noaa.gov/data-access/marineocean-data/extended-reconstructed-sea-surface-temperature-ersst-v4>). There is particular confidence in calculated warming for the time period since 1971 due to increased spatial and depth coverage and the level of agreement among independent SST observations from satellites, surface drifters and ships, and independent studies using differing analyses, bias corrections, and data sources (Cheng et al. 2017; Levitus et al. 2012; Llovel et al. 2014). Other observations such as the increase in mean sea level rise (see Ch. 12: Sea Level Rise) and reduced Arctic/Antarctic ice sheets (see Ch. 11: Arctic Changes) further confirm the increase in thermal expansion. For the purpose of extending the selected time periods back from 1900 to 2016 and analyzing U.S. regional SSTs, the Extended Reconstructed Sea Surface Temperature version 4 (ERSSTv4; Huang et al. 2015) is used. For the centennial time scale changes over 1900–2016, warming trends in all regions are statistically significant with the 95% confidence level. U.S. regional SST warming is similar between calculations using ERSSTv4 in this report and those published by Belkin (2016), suggesting confidence in these findings. The projected increase in SST is based on evidence from the latest generation of Earth System Models (CMIP5).

Major uncertainties

Uncertainties in the magnitude of ocean warming stem from the disparate measurements of ocean temperature over the last century. There is low uncertainty in warming trends of the upper ocean temperature from 0–700 m depth, whereas there is more uncertainty for deeper ocean depths of 700–2,000 m due to the short record of measurements from those areas. Data on warming trends at depths greater than 2,000 m are even more sparse. There are also uncertainties in the timing and reasons for particular decadal and interannual variations in ocean heat content and the contributions that different ocean basins play in the overall ocean heat uptake.

Summary sentence or paragraph that integrates the above information

There is *very high confidence* in measurements that show increases in the ocean heat content and warming of the ocean, based on the agreement of different methods. However, long-term data in total ocean heat uptake in the deep ocean are sparse leading to limited knowledge of the transport of heat between and within ocean basins.

Key Finding 2

The potential slowing of the Atlantic Meridional Overturning Circulation (AMOC) (of which the Gulf Stream is one component)—as a result of increasing ocean heat content and freshwater driven buoyancy changes—could have dramatic climate feedbacks as the ocean absorbs less heat and CO₂ from the atmosphere. This slowing would also affect the climates of North America and Europe. Any slowing documented to date cannot be directly tied to anthropogenic forcing primarily due to lack of adequate observational data and to challenges in modeling ocean circulation changes. Under a high emissions scenario (RCP8.5) in CMIP5 simulations, it is likely that the AMOC will weaken over the 21st century by 12% to 54%. (*Low confidence*)

Description of evidence base

Investigations both through direct observations and models since 2013 (Rhein et al. 2013) have raised significant concerns about whether there is enough evidence to determine the existence of an overall slowdown in the AMOC. As a result, more robust international observational campaigns are underway currently to measure AMOC circulation. Direct observations have determined a statistically significant slowdown at the 95% confidence level at 26°N (off Florida; see Baringer et al. 2016), but modeling studies constrained with observations cannot attribute this to anthropogenic forcing (Jackson et al. 2016). The study (Rahmstorf et al. 2015) which seemed to indicate broad-scale slowing has since been discounted due to its heavy reliance on sea surface temperature cooling as proxy for slowdown rather than actual direct observations. Since Rhein et al. 2013, more observations have led to increased statistical confidence in the measurement of the AMOC. Current observation trends indicate the AMOC slowing down at the 95% confidence level at 26°N and 41°N but a more limited in situ estimate at 35°S shows an increase in the AMOC (Smeed et al. 2014; Baringer et al. 2016). There is no one collection spot for AMOC-related data, but the U.S. Climate Variability and Predictability Program (US CLIVAR) has a U.S. AMOC priority focus area and a webpage with relevant data sites (<https://usclivar.org/amoc/amoc-time-series>).

The IPCC 2013 WG1 projections indicate a high likelihood of AMOC slowdown in the next 100 years, however overall understanding is limited by both a lack of direct observations (which is being remedied) and a lack of model skill to resolve deep ocean dynamics. As a result, this key finding was given an overall assessment of *low confidence*.

Major uncertainties

As noted, uncertainty about the overall trend of the AMOC is high given opposing trends in northern and southern ocean time series observations. Although earth system models do indicate a high likelihood of AMOC slowdown as a result of a warming, climate projections are subject to high uncertainty. This uncertainty stems from intermodel differences, internal variability that is different in each model, uncertainty in stratification changes, and most importantly uncertainty in both future freshwater input at high latitudes as well as the strength of the subpolar gyre circulation.

Summary sentence or paragraph that integrates the above information

The increased focus on direct measurements of the AMOC should lead to a better understanding of 1) how it is changing and its variability by region, and 2) whether those changes are attributable to climate drivers through both model improvements and incorporation of those expanded observations into the models.

Key Finding 3

The world's oceans are currently absorbing more than a quarter of the CO₂ emitted to the atmosphere annually from human activities, making them more acidic (*very high confidence*), with potential detrimental impacts to marine ecosystems. In particular, higher-latitude systems typically have a lower buffering capacity against pH change, exhibiting seasonally corrosive conditions sooner than low-latitude systems. Acidification is regionally increasing along U.S. coastal systems as a result of upwelling (for example, in the Pacific Northwest) (*high confidence*), changes in freshwater inputs (for example, in the Gulf of Maine) (*medium confidence*), and nutrient input (for example, in urbanized estuaries) (*high confidence*). The rate of acidification is unparalleled in at least the past 66 million years (*medium confidence*). Under RCP8.5, the global average surface ocean acidity is projected to increase by 100% to 150% (*high confidence*).

Description of evidence base

Evidence on the magnitude of the ocean sink is obtained from multiple biogeochemical and transport ocean models and two observation-based estimates from the 1990s for the uptake of the anthropogenic CO₂. Estimates of the carbonate system (DIC and alkalinity) were based on multiple survey cruises in the global ocean in the 1990s (WOCE, JGOFS). Coastal carbon and acidification surveys have been executed along the U.S. coastal large marine ecosystem since at least 2007, documenting significantly elevated pCO₂ and low pH conditions relative to oceanic waters. The data is available from the National Centers for Environmental Information (<https://www.ncei.noaa.gov/>). Other sources of biogeochemical bottle data can be found from

HOT-DOGS ALOHA (<http://hahana.soest.hawaii.edu/hot/hot-dogs>) or ERSI/GFM Data Finder (<https://www.esri.noaa.gov/gmd/dv/data>). Rates of change associated with the Palaeocene-Eocene Thermal Maximum (PETM, 56 million years ago) were derived using stable carbon and oxygen isotope records preserved in the sedimentary record from the New Jersey shelf using time series analysis and carbon cycle–climate modelling. This evidence supports a carbon release during the onset of the PETM over no less than 4,000 years, yielding a maximum sustained carbon release rate of less than 1.1 GtC per year (Zeebe et al. 2016). The projected increase in global surface ocean acidity is based on evidence from ten of the latest generation earth system models which include six distinct biogeochemical models that were included in the latest IPCC AR5 2013.

Major uncertainties

In 2014 the ocean sink was 2.6 ± 0.5 GtC (9.5 GtCO_2), equivalent to 26% of the total emissions attributed to fossil fuel use and land use changes (Le Quéré et al. 2016). Estimates of the PETM ocean acidification event evidenced in the geological record remains a matter of some debate within the community. Evidence for the 1.1 GtC per year cited by Zeebe et al. (2016), could be biased as a result of brief pulses of carbon input above average rates of emissions were they to transpire over timescales ≤ 40 years.

Summary sentence or paragraph that integrates the above information

There is *very high confidence* in evidence that the oceans absorb about a quarter of the carbon dioxide emitted in the atmosphere and hence become more acidic. The magnitude of the ocean carbon sink is known at a *high confidence* level because it is estimated using a series of disparate data sources and analysis methods, while the magnitude of the interannual variability is based only on model studies. There is medium confidence that the current rate of climate acidification is unprecedented in the past 66 million years. There is also *high confidence* that oceanic pH will continue to decrease.

Key Finding 4

Increasing sea surface temperatures, rising sea levels, and changing patterns of precipitation, winds, nutrients, and ocean circulation are contributing to overall declining oxygen concentrations at intermediate depths in various ocean locations and in many coastal areas. Over the last half century, major oxygen losses have occurred in inland seas, estuaries, and in the coastal and open ocean (*high confidence*). Ocean oxygen levels are projected to decrease by as much as 3.5% under the RCP8.5 scenario by 2100 relative to preindustrial values (*high confidence*).

1 Description of evidence base

2 The key finding and supporting text summarizes the evidence documented in climate science
3 literature including Rhein et al. 2013, Bopp et al. 2013, and Schmidtko et al. 2017. Evidence
4 arises from extensive global measurements of the World Ocean Circulation Experiment (WOCE)
5 after 1989 and individual profiles before that (Helm et al. 2011). The first basin-wide dissolved
6 oxygen surveys were performed in the 1920s (Schmidtko et al. 2017). The confidence level is
7 based on globally integrated O₂ distributions in a variety of ocean models. Although the global
8 mean exhibits low interannual variability, regional contrasts are large.

9 Major uncertainties

10 Uncertainties (as estimated from the intermodel spread) in the global mean are moderate mainly
11 because ocean oxygen content exhibits low interannual variability when globally averaged.
12 Uncertainties in long-term decreases of the global averaged oxygen concentration amount to
13 25% in the upper 1,000 m for the 1970–1992 period and 28% for the 1993–2003 period.
14 Remaining uncertainties relate to regional variability driven by mesoscale eddies and intrinsic
15 climate variability such as ENSO.

16 Summary sentence or paragraph that integrates the above information

17 Major ocean deoxygenation is taking place in bodies of water inland, at estuaries, and in the
18 coastal and the open ocean (*high confidence*). Regionally, the phenomenon is exacerbated by
19 local changes in weather, ocean circulation, and continental inputs to the oceans.

20

1 **TABLE**

2 **Table 13.1.** Historical sea surface temperature trends (°C per century) and projected trends by
3 2080 (°C) for eight U.S. coastal regions and globally. Historical temperature trends are presented
4 for the 1900–2016 and 1950–2016 periods with 95% confidence level, observed using the
5 Extended Reconstructed Sea Surface Temperature version 4 (ERSSTv4; Huang et al. 2015).
6 Global and regional predictions are calculated for RCP4.5 and RCP8.5 emission levels with 80%
7 spread of all the CMIP5 members compared to the 1976–2005 period (Scott et al. 2016). The
8 historical trends were analyzed for the latitude and longitude in the table, while the projected
9 trends were analyzed for the California current instead of the Northwest and Southwest
10 separately and for the Bering Sea in Alaska (NOAA).

Region	latitude and longitude	Historical Trend (°C/100 years)		Projected Trend by 2080 (relative to 1976- 2005 climate) (°C)	
		1900–2016	1950–2016	RCP4.5	RCP8.5
Global		0.70 ± 0.08	1.00 ± 0.11	1.3 ± 0.6	2.7 ± 0.7
Alaska	50°–66°N, 150°–170°W	0.82 ± 0.26	1.22 ± 0.59	2.5 ± 0.6	3.7 ± 1.0
Northwest (NW)	40°–50°N, 120°–132°W	0.64 ± 0.30	0.68 ± 0.70	1.7 ± 0.4	2.8 ± 0.6
Southwest (SW)	30°–40°N, 116°–126°W	0.73 ± 0.33	1.02 ± 0.79		
Hawaii (HI)	18°–24°N, 152°–162°W	0.58 ± 0.19	0.46 ± 0.39	1.6 ± 0.4	2.8 ± 0.6
Northeast (NE)	36°–46°N, 64°–76°W	0.63 ± 0.31	1.10 ± 0.71	2.0 ± 0.3	3.2 ± 0.6

Southeast (SE)	24°–34°N, 64°–80°W	0.40 ± 0.18	0.13 ± 0.34	1.6 ± 0.3	2.7 ± 0.4
Gulf of Mexico (GOM)	20°–30°N, 80°–96°W	0.52 ± 0.14	0.37 ± 0.27	1.6 ± 0.3	2.8 ± 0.3
Caribbean	10°–20°N, 66°–86°W	0.76 ± 0.15	0.77 ± 0.32	1.5 ± 0.4	2.6 ± 0.3

1

2


1 **FIGURES**

Figure 13.1: Global Ocean heat content change time series. Ocean heat content from 0 to 700 m (blue), 700 to 2,000 m (red), and 0 to 2,000 m (dark gray) from 1955 to 2015 with an uncertainty interval of ± 2 standard deviations shown in shading. All time series of the analysis performed by Cheng et al. (2017) are smoothed by a 12-month running mean filter, relative to the 1997–2005 base period. (Figure source: Cheng et al. 2017).




Figure 13.2: Ocean heat content changes from 1960 to 2015 for different ocean basins for 0 to 2,000 m depths. Time series is relative to the 1997–1999 base period and smoothed by a 12-month running filter by Cheng et al. (2017). The curves are additive, and the ocean heat content changes in different ocean basins are shaded in different colors (Figure source: Cheng et al. 2017).

CMIP5 ENSMN RCP8.5 Anomaly (2050–2099)–(1956–2005)




Figure 13.3: Projected changes in sea surface temperature (°C) for the coastal United States under emission scenario RCP8.5. Projected anomalies for the 2050–2099 period are calculated using a comparison from the average sea surface temperatures over 1956–2005. Projected changes are examined using the Coupled Model Intercomparison Project Phase 5 (CMIP5) suite of model simulations. (Figure source: NOAA).




Figure 13.4: Trends in surface (< 50 m) ocean carbonate chemistry calculated from observations obtained at the Hawai'i Ocean Time-series (HOT) Program in the North Pacific over 1988–2015. The upper panel shows the linked increase in atmospheric (red points) and seawater (blue points) CO_2 concentrations. The bottom panels shows a decline in seawater pH (black points, primary y-axis) and carbonate ion concentration (green points, secondary y-axis). Ocean chemistry data were obtained from the Hawai'i Ocean Time-series Data Organization & Graphical System (HOT-DOGS, <http://hahana.soest.hawaii.edu/hot/hot-dogs/index.html>). (Figure source: NOAA).

Surface pH in 2090s (RCP8.5, changes from 1990s)




Figure 13.5 Predicted change in sea surface pH in 2090–2099 relative to 1990–1999 under RCP8.5, based on the Community Earth System Models–Large Ensemble Experiments CMIP5 (Figure source: adapted from Bopp et al. 2013).

Projected Change in Dissolved Oxygen




Figure 13.6: Predicted change in dissolved oxygen on the $\sigma_\theta = 26.5$ (average depth of approximately 290 m) potential density surface, between the 1981–2000 and 2081–2100, based on the Community Earth System Models–Large Ensemble Experiments (Figure source: redrawn from Long et al. 2016).

1 **REFERENCES**

- 2 Abraham, J.P., M. Baringer, N.L. Bindoff, T. Boyer, L.J. Cheng, J.A. Church, J.L. Conroy, C.M.
 3 Domingues, J.T. Fasullo, J. Gilson, G. Goni, S.A. Good, J.M. Gorman, V. Gouretski, M.
 4 Ishii, G.C. Johnson, S. Kizu, J.M. Lyman, A.M. Macdonald, W.J. Minkowycz, S.E. Moffitt,
 5 M.D. Palmer, A.R. Piola, F. Reseghetti, K. Schuckmann, K.E. Trenberth, I. Velicogna, and
 6 J.K. Willis, 2013: A review of global ocean temperature observations: Implications for ocean
 7 heat content estimates and climate change. *Reviews of Geophysics*, **51**, 450–483.
 8 <http://dx.doi.org/10.1002/rog.20022>
- 9 Altieri, A.H. and K.B. Gedan, 2015: Climate change and dead zones. *Global Change Biology*,
 10 **21**, 1395–1406. <http://dx.doi.org/10.1111/gcb.12754>
- 11 Astor, Y.M., L. Lorenzoni, R. Thunell, R. Varela, F. Muller-Karger, L. Troccoli, G.T. Taylor,
 12 M.I. Scranton, E. Tappa, and D. Rueda, 2013: Interannual variability in sea surface
 13 temperature and $f\text{CO}_2$ changes in the Cariaco Basin. *Deep Sea Research Part II: Topical*
 14 *Studies in Oceanography*, **93**, 33–43. <http://dx.doi.org/10.1016/j.dsr2.2013.01.002>
- 15 Baden, S.P., L.O. Loo, L. Pihl, and R. Rosenberg, 1990: Effects of eutrophication on benthic
 16 communities including fish — Swedish west coast. *Ambio*, **19**, 113–122.
 17 www.jstor.org/stable/4313676
- 18 Bakun, A., B.A. Black, S.J. Bograd, M. García-Reyes, A.J. Miller, R.R. Rykaczewski, and W.J.
 19 Sydeman, 2015: Anticipated effects of climate change on coastal upwelling ecosystems.
 20 *Current Climate Change Reports*, **1**, 85–93. <http://dx.doi.org/10.1007/s40641-015-0008-4>
- 21 Baringer, M.O., M. Lankhorst, D. Volkov, S. Garzoli, S. Dong, U. Send, and C. Meinen, 2016:
 22 Meridional oceanic overturning circulation and heat transport in the Atlantic Ocean [in "State
 23 of the Climate in 2015"]. *Bulletin of the American Meteorological Society*, **97**, S84–S87.
 24 <http://dx.doi.org/10.1175/2015BAMSStateoftheClimate.1>
- 25 Bates, N.R. and J.T. Mathis, 2009: The Arctic Ocean marine carbon cycle: Evaluation of air-sea
 26 CO_2 exchanges, ocean acidification impacts and potential feedbacks. *Biogeosciences*, **6**,
 27 2433–2459. <http://dx.doi.org/10.5194/bg-6-2433-2009>
- 28 Bates, N.R., Y.M. Astor, M.J. Church, K. Currie, J.E. Dore, M. González-Dávila, L. Lorenzoni,
 29 F. Muller-Karger, J. Olafsson, and J.M. Santana-Casiano, 2014: A time-series view of
 30 changing ocean chemistry due to ocean uptake of anthropogenic CO_2 and ocean
 31 acidification. *Oceanography*, **27**, 126–141. <http://dx.doi.org/10.5670/oceanog.2014.16>
- 32 Belkin, I., 2016: Chapter 5.2: Sea surface temperature trends in large marine ecosystems. *Large*
 33 *Marine Ecosystems: Status and Trends*. United Nations Environment Programme, Nairobi,
 34 101–109.
 35 wedocs.unep.org/bitstream/handle/20.500.11822/13456/UNEP_DEWA_TWAP%20VOLUM

1 E%204%20REPORT_FINAL_4_MAY.pdf?sequence=1&isAllowed=y,%20English%20-
2 %20Summary

3 Bianchi, D., E.D. Galbraith, D.A. Carozza, K.A.S. Mislan, and C.A. Stock, 2013: Intensification
4 of open-ocean oxygen depletion by vertically migrating animals. *Nature Geoscience*, **6**, 545-
5 548. <http://dx.doi.org/10.1038/ngeo1837>

6 Boetius, A. and F. Wenzhofer, 2013: Seafloor oxygen consumption fuelled by methane from
7 cold seeps. *Nature Geoscience*, **6**, 725-734. <http://dx.doi.org/10.1038/ngeo1926>

8 Bograd, S.J., M.P. Buil, E.D. Lorenzo, C.G. Castro, I.D. Schroeder, R. Goericke, C.R. Anderson,
9 C. Benitez-Nelson, and F.A. Whitney, 2015: Changes in source waters to the Southern
10 California Bight. *Deep Sea Research Part II: Topical Studies in Oceanography*, **112**, 42-52.
11 <http://dx.doi.org/10.1016/j.dsr2.2014.04.009>

12 Bond, N.A., M.F. Cronin, H. Freeland, and N. Mantua, 2015: Causes and impacts of the 2014
13 warm anomaly in the NE Pacific. *Geophysical Research Letters*, **42**, 3414-3420.
14 <http://dx.doi.org/10.1002/2015GL063306>

15 Böning, C.W., E. Behrens, A. Biastoch, K. Getzlaff, and J.L. Bamber, 2016: Emerging impact of
16 Greenland meltwater on deepwater formation in the North Atlantic Ocean. *Nature*
17 *Geoscience*, **9**, 523-527. <http://dx.doi.org/10.1038/ngeo2740>

18 Booth, J.A.T., E.E. McPhee-Shaw, P. Chua, E. Kingsley, M. Denny, R. Phillips, S.J. Bograd,
19 L.D. Zeidberg, and W.F. Gilly, 2012: Natural intrusions of hypoxic, low pH water into
20 nearshore marine environments on the California coast. *Continental Shelf Research*, **45**, 108-
21 115. <http://dx.doi.org/10.1016/j.csr.2012.06.009>

22 Bopp, L., L. Resplandy, J.C. Orr, S.C. Doney, J.P. Dunne, M. Gehlen, P. Halloran, C. Heinze, T.
23 Ilyina, R. Séférian, J. Tjiputra, and M. Vichi, 2013: Multiple stressors of ocean ecosystems in
24 the 21st century: Projections with CMIP5 models. *Biogeosciences*, **10**, 6225-6245.
25 <http://dx.doi.org/10.5194/bg-10-6225-2013>

26 Borges, A.V. and N. Gypens, 2010: Carbonate chemistry in the coastal zone responds more
27 strongly to eutrophication than ocean acidification. *Limnology and Oceanography*, **55**, 346-
28 353. <http://dx.doi.org/10.4319/lo.2010.55.1.0346>

29 Boyer, T., C.M. Domingues, S.A. Good, G.C. Johnson, J.M. Lyman, M. Ishii, V. Gouretski, J.K.
30 Willis, J. Antonov, S. Wijffels, J.A. Church, R. Cowley, and N.L. Bindoff, 2016: Sensitivity
31 of global upper-ocean heat content estimates to mapping methods, XBT bias corrections, and
32 baseline climatologies. *Journal of Climate*, **29**, 4817-4842. <http://dx.doi.org/10.1175/jcli-d-15-0801.1>
33

- 1 Breitburg, D.L., J. Salisbury, J.M. Bernhard, W.-J. Cai, S. Dupont, S.C. Doney, K.J. Kroeker,
2 L.A. Levin, W.C. Long, L.M. Milke, S.H. Miller, B. Phelan, U. Passow, B.A. Seibel, A.E.
3 Todgham, and A.M. Tarrant, 2015: And on top of all that... Coping with ocean acidification
4 in the midst of many stressors. *Oceanography*, **28**, 48-61.
5 <http://dx.doi.org/10.5670/oceanog.2015.31>
- 6 Brush, G.S., 2009: Historical land use, nitrogen, and coastal eutrophication: A paleoecological
7 perspective. *Estuaries and Coasts*, **32**, 18-28. <http://dx.doi.org/10.1007/s12237-008-9106-z>
- 8 Bryden, H.L., H.R. Longworth, and S.A. Cunningham, 2005: Slowing of the Atlantic meridional
9 overturning circulation at 25[deg][thinsp]N. *Nature*, **438**, 655-657.
10 <http://dx.doi.org/10.1038/nature04385>
- 11 Bryden, H.L., B.A. King, and G.D. McCarthy, 2011: South Atlantic overturning circulation at
12 24°S. *Journal of Marine Research*, **69**, 38-55.
13 <http://dx.doi.org/10.1357/002224011798147633>
- 14 Buckley, M.W. and J. Marshall, 2016: Observations, inferences, and mechanisms of the Atlantic
15 Meridional Overturning Circulation: A review. *Reviews of Geophysics*, **54**, 5-63.
16 <http://dx.doi.org/10.1002/2015RG000493>
- 17 Carr, M.-E., M.A.M. Friedrichs, M. Schmeltz, M. Noguchi Aita, D. Antoine, K.R. Arrigo, I.
18 Asanuma, O. Aumont, R. Barber, M. Behrenfeld, R. Bidigare, E.T. Buitenhuis, J. Campbell,
19 A. Ciotti, H. Dierssen, M. Dowell, J. Dunne, W. Esaias, B. Gentili, W. Gregg, S. Groom, N.
20 Hoepffner, J. Ishizaka, T. Kameda, C. Le Quéré, S. Lohrenz, J. Marra, F. Mélin, K. Moore,
21 A. Morel, T.E. Reddy, J. Ryan, M. Scardi, T. Smyth, K. Turpie, G. Tilstone, K. Waters, and
22 Y. Yamanaka, 2006: A comparison of global estimates of marine primary production from
23 ocean color. *Deep Sea Research Part II: Topical Studies in Oceanography*, **53**, 741-770.
24 <http://dx.doi.org/10.1016/j.dsr2.2006.01.028>
- 25 Chavez, F.P., M. Messié, and J.T. Pennington, 2011: Marine primary production in relation to
26 climate variability and change. *Annual Review of Marine Science*, **3**, 227-260.
27 <http://dx.doi.org/10.1146/annurev.marine.010908.163917>
- 28 Cheng, L., K.E. Trenberth, J. Fasullo, T. Boyer, J. Abraham, and J. Zhu, 2017: Improved
29 estimates of ocean heat content from 1960 to 2015. *Science Advances*, **3**, e1601545.
30 <http://dx.doi.org/10.1126/sciadv.1601545>
- 31 Cheng, W., J.C.H. Chiang, and D. Zhang, 2013: Atlantic Meridional Overturning Circulation
32 (AMOC) in CMIP5 models: RCP and historical simulations. *Journal of Climate*, **26**, 7187-
33 7197. <http://dx.doi.org/10.1175/jcli-d-12-00496.1>

- Chiapello, I., 2014: Dust observations and climatology. *Mineral Dust: A Key Player in the Earth System*. Knippertz, P. and J.-B.W. Stuut, Eds. Springer Netherlands, Dordrecht, 149-177.
http://dx.doi.org/10.1007/978-94-017-8978-3_7
- Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, M. Heimann, C. Jones, C. Le Quéré, R.B. Myneni, S. Piao, and P. Thornton, 2013: Carbon and other biogeochemical cycles. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley, Eds. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 465–570. <http://www.climatechange2013.org/report/full-report/>
- Codispoti, L.A., J.A. Brandes, J.P. Christensen, A.H. Devol, S.W.A. Naqvi, H.W. Paerl, and T. Yoshinari, 2001: The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene? *Scientia Marina*, **65**, 85-105.
<http://dx.doi.org/10.3989/scimar.2001.65s285>
- Collins, M., R. Knutti, J. Arblaster, J.-L. Dufresne, T. Fichefet, P. Friedlingstein, X. Gao, W.J. Gutowski, T. Johns, G. Krinner, M. Shongwe, C. Tebaldi, A.J. Weaver, and M. Wehner, 2013: Long-term climate change: Projections, commitments and irreversibility. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley, Eds. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1029–1136. <http://www.climatechange2013.org/report/full-report/>
- Conley, D.J., J. Carstensen, J. Aigars, P. Axe, E. Bonsdorff, T. Eremina, B.-M. Haahti, C. Humborg, P. Jonsson, J. Kotta, C. Lännegren, U. Larsson, A. Maximov, M.R. Medina, E. Lysiak-Pastuszak, N. Remeikaitė-Nikienė, J. Walve, S. Wilhelms, and L. Zillén, 2011: Hypoxia is increasing in the coastal zone of the Baltic Sea. *Environmental Science & Technology*, **45**, 6777-6783. <http://dx.doi.org/10.1021/es201212r>
- Deutsch, C., H. Brix, T. Ito, H. Frenzel, and L. Thompson, 2011: Climate-forced variability of ocean hypoxia. *Science*, **333**, 336-339. <http://dx.doi.org/10.1126/science.1202422>
- Deutsch, C., W. Berelson, R. Thunell, T. Weber, C. Tems, J. McManus, J. Crusius, T. Ito, T. Baumgartner, V. Ferreira, J. Mey, and A. van Geen, 2014: Centennial changes in North Pacific anoxia linked to tropical trade winds. *Science*, **345**, 665-668.
<http://dx.doi.org/10.1126/science.1252332>
- Diaz, R.J. and R. Rosenberg, 2008: Spreading dead zones and consequences for marine ecosystems. *Science*, **321**, 926-929. <http://dx.doi.org/10.1126/science.1156401>

- 1 Dong, S., G. Goni, and F. Bringas, 2015: Temporal variability of the South Atlantic Meridional
2 Overturning Circulation between 20°S and 35°S. *Geophysical Research Letters*, **42**, 7655-
3 7662. <http://dx.doi.org/10.1002/2015GL065603>
- 4 Donner, S.D. and D. Scavia, 2007: How climate controls the flux of nitrogen by the Mississippi
5 River and the development of hypoxia in the Gulf of Mexico. *Limnology and Oceanography*,
6 **52**, 856-861. <http://dx.doi.org/10.4319/lo.2007.52.2.0856>
- 7 Drijfhout, S.S., S.L. Weber, and E. van der Swaluw, 2011: The stability of the MOC as
8 diagnosed from model projections for pre-industrial, present and future climates. *Climate*
9 *Dynamics*, **37**, 1575-1586. <http://dx.doi.org/10.1007/s00382-010-0930-z>
- 10 Duarte, C.M., Hendriks, I.E., Moore, T.S., Olsen, Y.S., Steckbauer, A., Ramajo, L., Carstensen,
11 J., Trotter, J.A. and McCulloch, M., 2013. Is ocean acidification an open-ocean syndrome?
12 Understanding anthropogenic impacts on seawater pH. *Estuaries and Coasts*, 36(2), pp.221-
13 236. DOI: 10.1007/s12237-013-9594-3
- 14 Durack, P.J. and S.E. Wijffels, 2010: Fifty-year trends in global ocean salinities and their
15 relationship to broad-scale warming. *Journal of Climate*, **23**, 4342-4362.
16 <http://dx.doi.org/10.1175/2010jcli3377.1>
- 17 EPA. 2017: Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2015.
18 https://www.epa.gov/sites/production/files/2017-02/documents/2017_complete_report.pdf
- 19 Evans, W., J.T. Mathis, and J.N. Cross, 2014: Calcium carbonate corrosivity in an Alaskan
20 inland sea. *Biogeosciences*, **11**, 365-379. <http://dx.doi.org/10.5194/bg-11-365-2014>
- 21 Falkowski, P.G., T. Algeo, L. Codispoti, C. Deutsch, S. Emerson, B. Hales, R.B. Huey, W.J.
22 Jenkins, L.R. Kump, L.A. Levin, T.W. Lyons, N.B. Nelson, O.S. Schofield, R. Summons,
23 L.D. Talley, E. Thomas, F. Whitney, and C.B. Pilcher, 2011: Ocean deoxygenation: Past,
24 present, and future. *Eos, Transactions, American Geophysical Union*, **92**, 409-410.
25 <http://dx.doi.org/10.1029/2011EO460001>
- 26 Fasullo, J.T., R.S. Nerem, and B. Hamlington, 2016: Is the detection of accelerated sea level rise
27 imminent? *Scientific Reports*, **6**, 31245. <http://dx.doi.org/10.1038/srep31245>
- 28 Feely, R.A., C.L. Sabine, K. Lee, W. Berelson, J. Kleypas, V.J. Fabry, and F.J. Millero, 2004:
29 Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans. *Science*, **305**, 362-366.
30 <http://dx.doi.org/10.1126/science.1097329>
- 31 Feely, R.A., C.L. Sabine, J.M. Hernandez-Ayon, D. Ianson, and B. Hales, 2008: Evidence for
32 upwelling of corrosive "acidified" water onto the continental shelf. *Science*, **320**, 1490-1492.
33 <http://dx.doi.org/10.1126/science.1155676>

- 1 Feely, R.A., S.C. Doney, and S.R. Cooley, 2009: Ocean acidification: Present conditions and
2 future changes in a high-CO₂ world. *Oceanography*, **22**, 36-47.
3 <http://dx.doi.org/10.5670/oceanog.2009.95>
- 4 Franz, B.A., M.J. Behrenfeld, D.A. Siegel, and S.R. Signorini, 2016: Global ocean
5 phytoplankton [in "State of the Climate in 2015"]. *Bulletin of the American Meteorological*
6 *Society*, **97**, S87–S89. <http://dx.doi.org/10.1175/2016BAMSStateoftheClimate.1>
- 7 Friedrich, T., A. Timmermann, A. Abe-Ouchi, N.R. Bates, M.O. Chikamoto, M.J. Church, J.E.
8 Dore, D.K. Gledhill, M. Gonzalez-Davila, M. Heinemann, T. Ilyina, J.H. Jungclaus, E.
9 McLeod, A. Mouchet, and J.M. Santana-Casiano, 2012: Detecting regional anthropogenic
10 trends in ocean acidification against natural variability. *Nature Climate Change*, **2**, 167-171.
11 <http://dx.doi.org/10.1038/nclimate1372>
- 12 Frölicher, T.L., K.B. Rodgers, C.A. Stock, and W.W.L. Cheung, 2016: Sources of uncertainties
13 in 21st century projections of potential ocean ecosystem stressors. *Global Biogeochemical*
14 *Cycles*, **30**, 1224-1243. <http://dx.doi.org/10.1002/2015GB005338>
- 15 Fu, W., J.T. Randerson, and J.K. Moore, 2016: Climate change impacts on net primary
16 production (NPP) and export production (EP) regulated by increasing stratification and
17 phytoplankton community structure in the CMIP5 models. *Biogeosciences*, **13**, 5151-5170.
18 <http://dx.doi.org/10.5194/bg-13-5151-2016>
- 19 Galbraith, E.D., M. Kienast, T.F. Pedersen, and S.E. Calvert, 2004: Glacial-interglacial
20 modulation of the marine nitrogen cycle by high-latitude O₂ supply to the global
21 thermocline. *Paleoceanography*, **19**, n/a-n/a. <http://dx.doi.org/10.1029/2003PA001000>
- 22 García-Reyes, M., W.J. Sydeman, D.S. Schoeman, R.R. Rykaczewski, B.A. Black, A.J. Smit,
23 and S.J. Bograd, 2015: Under pressure: Climate change, upwelling, and eastern boundary
24 upwelling ecosystems. *Frontiers in Marine Science*, **2**.
25 <http://dx.doi.org/10.3389/fmars.2015.00109>
- 26 Garzoli, S.L., M.O. Baringer, S. Dong, R.C. Perez, and Q. Yao, 2013: South Atlantic meridional
27 fluxes. *Deep Sea Research Part I: Oceanographic Research Papers*, **71**, 21-32.
28 <http://dx.doi.org/10.1016/j.dsr.2012.09.003>
- 29 Gattuso, J.-P., A. Magnan, R. Billé, W.W.L. Cheung, E.L. Howes, F. Joos, D. Allemand, L.
30 Bopp, S.R. Cooley, C.M. Eakin, O. Hoegh-Guldberg, R.P. Kelly, H.-O. Pörtner, A.D.
31 Rogers, J.M. Baxter, D. Laffoley, D. Osborn, A. Rankovic, J. Rochette, U.R. Sumaila, S.
32 Treyer, and C. Turley, 2015: Contrasting futures for ocean and society from different
33 anthropogenic CO₂ emissions scenarios. *Science*, **349**.
34 <http://dx.doi.org/10.1126/science.aac4722>

- 1 Gilbert, D., B. Sundby, C. Gobeil, A. Mucci, and G.-H. Tremblay, 2005: A seventy-two-year
2 record of diminishing deep-water oxygen in the St. Lawrence estuary: The northwest Atlantic
3 connection. *Limnology and Oceanography*, **50**, 1654-1666.
4 <http://dx.doi.org/10.4319/lo.2005.50.5.1654>
- 5 Gilbert, D., N.N. Rabalais, R.J. Díaz, and J. Zhang, 2010: Evidence for greater oxygen decline
6 rates in the coastal ocean than in the open ocean. *Biogeosciences*, **7**, 2283-2296.
7 <http://dx.doi.org/10.5194/bg-7-2283-2010>
- 8 Gledhill, D.K., M.M. White, J. Salisbury, H. Thomas, I. Mlsna, M. Liebman, B. Mook, J. Grear,
9 A.C. Candelmo, R.C. Chambers, C.J. Gobler, C.W. Hunt, A.L. King, N.N. Price, S.R.
10 Signorini, E. Stancioff, C. Stymiest, R.A. Wahle, J.D. Waller, N.D. Rebeck, Z.A. Wang, T.L.
11 Capson, J.R. Morrison, S.R. Cooley, and S.C. Doney, 2015: Ocean and coastal acidification
12 off New England and Nova Scotia. *Oceanography*, **28**, 182-197.
13 <http://dx.doi.org/10.5670/oceanog.2015.41>
- 14 Gobler, C.J., E.L. DePasquale, A.W. Griffith, and H. Baumann, 2014: Hypoxia and acidification
15 have additive and synergistic negative effects on the growth, survival, and metamorphosis of
16 early life stage bivalves. *PLoS ONE*, **9**, e83648.
17 <http://dx.doi.org/10.1371/journal.pone.0083648>
- 18 Gruber, N., 2008: Chapter 1 - The marine nitrogen cycle: Overview and challenges. *Nitrogen in*
19 *the Marine Environment (2nd Edition)*. Academic Press, San Diego, 1-50.
20 <http://dx.doi.org/10.1016/B978-0-12-372522-6.00001-3>
- 21 Harris, K.E., M.D. DeGrandpre, and B. Hales, 2013: Aragonite saturation state dynamics in a
22 coastal upwelling zone. *Geophysical Research Letters*, **40**, 2720-2725.
23 <http://dx.doi.org/10.1002/grl.50460>
- 24 Helm, K.P., N.L. Bindoff, and J.A. Church, 2011: Observed decreases in oxygen content of the
25 global ocean. *Geophysical Research Letters*, **38**, n/a-n/a.
26 <http://dx.doi.org/10.1029/2011GL049513>
- 27 Hendriks, I.E., C.M. Duarte, Y.S. Olsen, A. Steckbauer, L. Ramajo, T.S. Moore, J.A. Trotter,
28 and M. McCulloch, 2015: Biological mechanisms supporting adaptation to ocean
29 acidification in coastal ecosystems. *Estuarine, Coastal and Shelf Science*, **152**, A1-A8.
30 <http://dx.doi.org/10.1016/j.ecss.2014.07.019>
- 31 Henson, S.A., Sarmiento, J.L., Dunne, J.P., Bopp, L., Lima, I.D., Doney, S.C., John, J.G. and
32 Beaulieu, C., 2010. Detection of anthropogenic climate change in satellite records of ocean
33 chlorophyll and productivity. *Biogeosciences* 7: 621-640, doi:10.5194/bg-7-621-2010

- 1 Henson, S.A., Beaulieu, C. and Lampitt, R., 2016. Observing climate change trends in ocean
2 biogeochemistry: when and where. *Global change biology*, 22(4), pp.1561-1571. DOI:
3 10.1111/gcb.13152
- 4 Hoegh-Guldberg, O., R. Cai, E.S. Poloczanska, P.G. Brewer, S. Sundby, K. Hilmi, V.J. Fabry,
5 and S. Jung, 2014: The Ocean--Supplementary material. *Climate Change 2014: Impacts,*
6 *Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II*
7 *to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. Barros,
8 V.R., C.B. Field, D.J. Dokken, M.D. Mastrandrea, K.J. Mach, T.E. Bilir, M. Chatterjee, K.L.
9 Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R.
10 Mastrandrea, and L.L. White, Eds. Cambridge University Press, Cambridge, United
11 Kingdom and New York, NY, USA, 1655-1731. [ipcc.ch/pdf/assessment-](http://ipcc.ch/pdf/assessment-report/ar5/wg2/supplementary/WGIIAR5-Chap30_OLSM.pdf)
12 [report/ar5/wg2/supplementary/WGIIAR5-Chap30_OLSM.pdf](http://ipcc.ch/pdf/assessment-report/ar5/wg2/supplementary/WGIIAR5-Chap30_OLSM.pdf)
- 13 Hönisch, B., A. Ridgwell, D.N. Schmidt, E. Thomas, S.J. Gibbs, A. Sluijs, R. Zeebe, L. Kump,
14 R.C. Martindale, S.E. Greene, W. Kiessling, J. Ries, J.C. Zachos, D.L. Royer, S. Barker,
15 T.M. Marchitto, Jr., R. Moyer, C. Pelejero, P. Ziveri, G.L. Foster, and B. Williams, 2012:
16 The geological record of ocean acidification. *Science*, **335**, 1058-1063.
17 <http://dx.doi.org/10.1126/science.1208277>
- 18 Huang, B., V.F. Banzon, E. Freeman, J. Lawrimore, W. Liu, T.C. Peterson, T.M. Smith, P.W.
19 Thorne, S.D. Woodruff, and H.-M. Zhang, 2015: Extended Reconstructed Sea Surface
20 Temperature Version 4 (ERSST.v4). Part I: Upgrades and intercomparisons. *Journal of*
21 *Climate*, **28**, 911-930. <http://dx.doi.org/10.1175/JCLI-D-14-00006.1>
- 22 IPCC, 2013: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group*
23 *I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
24 Cambridge University Press, Cambridge, UK and New York, NY, 1535 pp.
25 <http://www.climatechange2013.org/report/>
- 26 Ito, T., A. Nenes, M.S. Johnson, N. Meskhidze, and C. Deutsch, 2016: Acceleration of oxygen
27 decline in the tropical Pacific over the past decades by aerosol pollutants. *Nature Geoscience*,
28 **9**, 443-447. <http://dx.doi.org/10.1038/ngeo2717>
- 29 Jackson, L.C., K.A. Peterson, C.D. Roberts, and R.A. Wood, 2016: Recent slowing of Atlantic
30 overturning circulation as a recovery from earlier strengthening. *Nature Geoscience*, **9**, 518-
31 522. <http://dx.doi.org/10.1038/ngeo2715>
- 32 Jacox, M.G., A.M. Moore, C.A. Edwards, and J. Fiechter, 2014: Spatially resolved upwelling in
33 the California Current System and its connections to climate variability. *Geophysical*
34 *Research Letters*, **41**, 3189-3196. <http://dx.doi.org/10.1002/2014GL059589>

- 1 Jickells, T. and C.M. Moore, 2015: The importance of atmospheric deposition for ocean
2 productivity. *Annual Review of Ecology, Evolution, and Systematics*, **46**, 481-501.
3 <http://dx.doi.org/10.1146/annurev-ecolsys-112414-054118>
- 4 Johns, W.E., M.O. Baringer, L.M. Beal, S.A. Cunningham, T. Kanzow, H.L. Bryden, J.J.M.
5 Hirschi, J. Marotzke, C.S. Meinen, B. Shaw, and R. Curry, 2011: Continuous, array-based
6 estimates of Atlantic ocean heat transport at 26.5°N. *Journal of Climate*, **24**, 2429-2449.
7 <http://dx.doi.org/10.1175/2010jcli3997.1>
- 8 Justić, D., T. Legović, and L. Rottini-Sandrini, 1987: Trends in oxygen content 1911–1984 and
9 occurrence of benthic mortality in the northern Adriatic Sea. *Estuarine, Coastal and Shelf*
10 *Science*, **25**, 435-445. [http://dx.doi.org/10.1016/0272-7714\(87\)90035-7](http://dx.doi.org/10.1016/0272-7714(87)90035-7)
- 11 Kanzow, T., S.A. Cunningham, W.E. Johns, J.J.-M. Hirschi, J. Marotzke, M.O. Baringer, C.S.
12 Meinen, M.P. Chidichimo, C. Atkinson, L.M. Beal, H.L. Bryden, and J. Collins, 2010:
13 Seasonal variability of the Atlantic meridional overturning circulation at 26.5°N. *Journal of*
14 *Climate*, **23**, 5678-5698. <http://dx.doi.org/10.1175/2010jcli3389.1>
- 15 Knoll, A.H. and S.B. Carroll, 1999: Early animal evolution: Emerging views from comparative
16 biology and geology. *Science*, **284**, 2129-2137.
17 <http://dx.doi.org/10.1126/science.284.5423.2129>
- 18 Kosaka, Y. and Xie, S.P., 2013. Recent global-warming hiatus tied to equatorial Pacific surface cooling.
19 *Nature*, 501(7467), pp.403-407. doi:10.1038/nature12534
- 20 Laufkötter, C., M. Vogt, N. Gruber, M. Aita-Noguchi, O. Aumont, L. Bopp, E. Buitenhuis, S.C.
21 Doney, J. Dunne, T. Hashioka, J. Hauck, T. Hirata, J. John, C. Le Quéré, I.D. Lima, H.
22 Nakano, R. Seferian, I. Totterdell, M. Vichi, and C. Völker, 2015: Drivers and uncertainties
23 of future global marine primary production in marine ecosystem models. *Biogeosciences*, **12**,
24 6955-6984. <http://dx.doi.org/10.5194/bg-12-6955-2015>
- 25 Le Quéré, C., R.M. Andrew, J.G. Canadell, S. Sitch, J.I. Korsbakken, G.P. Peters, A.C. Manning,
26 T.A. Boden, P.P. Tans, R.A. Houghton, R.F. Keeling, S. Alin, O.D. Andrews, P. Anthoni, L.
27 Barbero, L. Bopp, F. Chevallier, L.P. Chini, P. Ciais, K. Currie, C. Delire, S.C. Doney, P.
28 Friedlingstein, T. Gkritzalis, I. Harris, J. Hauck, V. Haverd, M. Hoppema, K. Klein
29 Goldewijk, A.K. Jain, E. Kato, A. Körtzinger, P. Landschützer, N. Lefèvre, A. Lenton, S.
30 Lienert, D. Lombardozzi, J.R. Melton, N. Metzl, F. Millero, P.M.S. Monteiro, D.R. Munro,
31 J.E.M.S. Nabel, S.I. Nakaoka, K. O'Brien, A. Olsen, A.M. Omar, T. Ono, D. Pierrot, B.
32 Poulter, C. Rödenbeck, J. Salisbury, U. Schuster, J. Schwinger, R. Séférian, I. Skjelvan, B.D.
33 Stocker, A.J. Sutton, T. Takahashi, H. Tian, B. Tilbrook, I.T. van der Laan-Luijkx, G.R. van
34 der Werf, N. Viovy, A.P. Walker, A.J. Wiltshire, and S. Zaehle, 2016: Global carbon budget
35 2016. *Earth System Science Data*, **8**, 605-649. <http://dx.doi.org/10.5194/essd-8-605-2016>

- 1 Lee, M., E. Shevliakova, S. Malyshev, P.C.D. Milly, and P.R. Jaffé, 2016: Climate variability
2 and extremes, interacting with nitrogen storage, amplify eutrophication risk. *Geophysical*
3 *Research Letters*, **43**, 7520-7528. <http://dx.doi.org/10.1002/2016GL069254>
- 4 Lee, S.-K., W. Park, M.O. Baringer, A.L. Gordon, B. Huber, and Y. Liu, 2015: Pacific origin of
5 the abrupt increase in Indian Ocean heat content during the warming hiatus. *Nature*
6 *Geoscience*, **8**, 445-449. <http://dx.doi.org/10.1038/ngeo2438>
- 7 Levitus, S., J.I. Antonov, T.P. Boyer, O.K. Baranova, H.E. Garcia, R.A. Locarnini, A.V.
8 Mishonov, J.R. Reagan, D. Seidov, E.S. Yarosh, and M.M. Zweng, 2012: World ocean heat
9 content and thermosteric sea level change (0–2000 m), 1955–2010. *Geophysical Research*
10 *Letters*, **39**, L10603. <http://dx.doi.org/10.1029/2012GL051106>
- 11 Liu, W., S.-P. Xie, Z. Liu, and J. Zhu, 2017: Overlooked possibility of a collapsed Atlantic
12 Meridional Overturning Circulation in warming climate. *Science Advances*, **3**, e1601666.
13 <http://dx.doi.org/10.1126/sciadv.1601666>
- 14 Llovel, W., J.K. Willis, F.W. Landerer, and I. Fukumori, 2014: Deep-ocean contribution to sea
15 level and energy budget not detectable over the past decade. *Nature Climate Change*, **4**,
16 1031-1035. <http://dx.doi.org/10.1038/nclimate2387>
- 17 Long, M.C., C. Deutsch, and T. Ito, 2016: Finding forced trends in oceanic oxygen. *Global*
18 *Biogeochemical Cycles*, **30**, 381-397. <http://dx.doi.org/10.1002/2015GB005310>
- 19 Longworth, H.R., H.L. Bryden, and M.O. Baringer, 2011: Historical variability in Atlantic
20 meridional baroclinic transport at 26.5°N from boundary dynamic height observations. *Deep*
21 *Sea Research Part II: Topical Studies in Oceanography*, **58**, 1754-1767.
22 <http://dx.doi.org/10.1016/j.dsr2.2010.10.057>
- 23 Mathis, J.T., S.R. Cooley, N. Lucey, S. Colt, J. Ekstrom, T. Hurst, C. Hauri, W. Evans, J.N.
24 Cross, and R.A. Feely, 2015: Ocean acidification risk assessment for Alaska's fishery sector.
25 *Progress in Oceanography*, **136**, 71-91. <http://dx.doi.org/10.1016/j.pocean.2014.07.001>
- 26 Matthews, J. B. R. "Comparing historical and modern methods of sea surface temperature
27 measurement—Part 1: Review of methods, field comparisons and dataset adjustments." *Ocean*
28 *Science* 9.4 (2013): 683-694. doi:10.5194/os-9-683-2013
- 29 McDonagh, E.L., B.A. King, H.L. Bryden, P. Courtois, Z. Szuts, M. Baringer, S.A. Cunningham,
30 C. Atkinson, and G. McCarthy, 2015: Continuous estimate of Atlantic oceanic freshwater
31 flux at 26.5°N. *Journal of Climate*, **28**, 8888-8906. <http://dx.doi.org/10.1175/jcli-d-14->
32 00519.1
- 33 McFall-Ngai, M., M.G. Hadfield, T.C.G. Bosch, H.V. Carey, T. Domazet-Lošo, A.E. Douglas,
34 N. Dubilier, G. Eberl, T. Fukami, S.F. Gilbert, U. Hentschel, N. King, S. Kjelleberg, A.H.

- Knoll, N. Kremer, S.K. Mazmanian, J.L. Metcalf, K. Nealson, N.E. Pierce, J.F. Rawls, A. Reid, E.G. Ruby, M. Rumpho, J.G. Sanders, D. Tautz, and J.J. Wernegreen, 2013: Animals in a bacterial world, a new imperative for the life sciences. *Proceedings of the National Academy of Sciences*, **110**, 3229-3236. <http://dx.doi.org/10.1073/pnas.1218525110>
- Mei, W., S.-P. Xie, F. Primeau, J.C. McWilliams, and C. Pasquero, 2015: Northwestern Pacific typhoon intensity controlled by changes in ocean temperatures. *Science Advances*, **1**. <http://dx.doi.org/10.1126/sciadv.1500014>
- Moffitt, S.E., R.A. Moffitt, W. Sauthoff, C.V. Davis, K. Hewett, and T.M. Hill, 2015: Paleoceanographic insights on recent oxygen minimum zone expansion: Lessons for modern oceanography. *PLoS ONE*, **10**, e0115246. <http://dx.doi.org/10.1371/journal.pone.0115246>
- Monbaliu, J., Z. Chen, D. Felts, J. Ge, F. Hissel, J. Kappenberg, S. Narayan, R.J. Nicholls, N. Ohle, D. Schuster, J. Sothmann, and P. Willems, 2014: Risk assessment of estuaries under climate change: Lessons from Western Europe. *Coastal Engineering*, **87**, 32-49. <http://dx.doi.org/10.1016/j.coastaleng.2014.01.001>
- Montes, I., B. Dewitte, E. Gutknecht, A. Paulmier, I. Dadou, A. Oschlies, and V. Garçon, 2014: High-resolution modeling of the eastern tropical Pacific oxygen minimum zone: Sensitivity to the tropical oceanic circulation. *Journal of Geophysical Research: Oceans*, **119**, 5515-5532. <http://dx.doi.org/10.1002/2014JC009858>
- Mulitza, S., D. Heslop, D. Pittauerova, H.W. Fischer, I. Meyer, J.-B. Stuut, M. Zabel, G. Mollenhauer, J.A. Collins, H. Kuhnert, and M. Schulz, 2010: Increase in African dust flux at the onset of commercial agriculture in the Sahel region. *Nature*, **466**, 226-228. <http://dx.doi.org/10.1038/nature09213>
- Najjar, R.G., C.R. Pyke, M.B. Adams, D. Breitburg, C. Hershner, M. Kemp, R. Howarth, M.R. Mulholland, M. Paolisso, D. Secor, K. Sellner, D. Wardrop, and R. Wood, 2010: Potential climate-change impacts on the Chesapeake Bay. *Estuarine, Coastal and Shelf Science*, **86**, 1-20. <http://dx.doi.org/10.1016/j.ecss.2009.09.026>
- Nieves, V., J.K. Willis, and W.C. Patzert, 2015: Recent hiatus caused by decadal shift in Indo-Pacific heating. *Science*, **349**, 532-535. <http://dx.doi.org/10.1126/science.aaa4521>
- Orr, J.C., V.J. Fabry, O. Aumont, L. Bopp, S.C. Doney, R.A. Feely, A. Gnanadesikan, N. Gruber, A. Ishida, F. Joos, R.M. Key, K. Lindsay, E. Maier-Reimer, R. Matear, P. Monfray, A. Mouchet, R.G. Najjar, G.-K. Plattner, K.B. Rodgers, C.L. Sabine, J.L. Sarmiento, R. Schlitzer, R.D. Slater, I.J. Totterdell, M.-F. Weirig, Y. Yamanaka, and A. Yool, 2005: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**, 681-686. <http://dx.doi.org/10.1038/nature04095>

- 1 Paerl, H.W., R.L. Dennis, and D.R. Whitall, 2002: Atmospheric deposition of nitrogen:
2 Implications for nutrient over-enrichment of coastal waters. *Estuaries*, **25**, 677-693.
3 <http://dx.doi.org/10.1007/bf02804899>
- 4 Patara, L. and C.W. Böning, 2014: Abyssal ocean warming around Antarctica strengthens the
5 Atlantic overturning circulation. *Geophysical Research Letters*, **41**, 3972-3978.
6 <http://dx.doi.org/10.1002/2014GL059923>
- 7 Purkey, S.G. and G.C. Johnson, 2010: Warming of global abyssal and deep Southern Ocean
8 waters between the 1990s and 2000s: Contributions to global heat and sea level rise budgets.
9 *Journal of Climate*, **23**, 6336-6351. <http://dx.doi.org/10.1175/2010JCLI3682.1>
- 10 Qi, D., L. Chen, B. Chen, Z. Gao, W. Zhong, R.A. Feely, L.G. Anderson, H. Sun, J. Chen, M.
11 Chen, L. Zhan, Y. Zhang, and W.-J. Cai, 2017: Increase in acidifying water in the western
12 Arctic Ocean. *Nature Climate Change*, **7**, 195-199. <http://dx.doi.org/10.1038/nclimate3228>
- 13 Rabalais, N.N., R.E. Turner, B.K. Sen Gupta, D.F. Boesch, P. Chapman, and M.C. Murrell,
14 2007: Hypoxia in the northern Gulf of Mexico: Does the science support the plan to reduce,
15 mitigate, and control hypoxia? *Estuaries and Coasts*, **30**, 753-772.
16 <http://dx.doi.org/10.1007/bf02841332>
- 17 Rabalais, N.N., R.E. Turner, R.J. Díaz, and D. Justić, 2009: Global change and eutrophication of
18 coastal waters. *ICES Journal of Marine Science: Journal du Conseil*, **66**, 1528-1537.
19 <http://dx.doi.org/10.1093/icesjms/fsp047>
- 20 Rabalais, N.N., R.J. Díaz, L.A. Levin, R.E. Turner, D. Gilbert, and J. Zhang, 2010: Dynamics
21 and distribution of natural and human-caused hypoxia. *Biogeosciences*, **7**, 585-619.
22 <http://dx.doi.org/10.5194/bg-7-585-2010>
- 23 Rahmstorf, S., J.E. Box, G. Feulner, M.E. Mann, A. Robinson, S. Rutherford, and E.J.
24 Schaffernicht, 2015: Exceptional twentieth-century slowdown in Atlantic Ocean overturning
25 circulation. *Nature Climate Change*, **5**, 475-480. <http://dx.doi.org/10.1038/nclimate2554>
- 26 Reay, D.S., F. Dentener, P. Smith, J. Grace, and R.A. Feely, 2008: Global nitrogen deposition
27 and carbon sinks. *Nature Geoscience*, **1**, 430-437. <http://dx.doi.org/10.1038/ngeo230>
- 28 Rhein, M., S.R. Rintoul, S. Aoki, E. Campos, D. Chambers, R.A. Feely, S. Gulev, G.C. Johnson,
29 S.A. Josey, A. Kostianoy, C. Mauritzen, D. Roemmich, L.D. Talley, and F. Wang, 2013:
30 Observations: Ocean. *Climate Change 2013: The Physical Science Basis. Contribution of*
31 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
32 *Change*. Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.
33 Nauels, Y. Xia, V. Bex, and P.M. Midgley, Eds. Cambridge University Press, Cambridge,
34 United Kingdom and New York, NY, USA, 255–316.
35 <http://www.climatechange2013.org/report/full-report/>

- 1 Robinson, P., A.K. Leight, D.D. Trueblood, and B. Wood, 2013: Climate sensitivity of the
2 National Estuarine Research Reserve System. NERRS, NOAA National Ocean Service,
3 Silver Spring, Maryland. 79 pp.
4 https://coast.noaa.gov/data/docs/nerrs/Research_DataSyntheses_130725_climate%20sensitivity%20of%20nerrs_Final-Rpt-in-Layout_FINAL.pdf
5
- 6 Robinson, R.S., A. Mix, and P. Martinez, 2007: Southern Ocean control on the extent of
7 denitrification in the southeast Pacific over the last 70 ka. *Quaternary Science Reviews*, **26**,
8 201-212. <http://dx.doi.org/10.1016/j.quascirev.2006.08.005>
- 9 Roemmich, D., J. Church, J. Gilson, D. Monselesan, P. Sutton, and S. Wijffels, 2015: Unabated
10 planetary warming and its ocean structure since 2006. *Nature Climate Change*, **5**, 240-245.
11 <http://dx.doi.org/10.1038/nclimate2513>
- 12 Rossby, C.-G., 1959: Current problems in meteorology. *The Atmosphere and the Sea in Motion*.
13 Bolin, B., Ed. Rockefeller Institute Press, New York, 9-50.
- 14 Rykaczewski, R.R., J.P. Dunne, W.J. Sydeman, M. García-Reyes, B.A. Black, and S.J. Bograd,
15 2015: Poleward displacement of coastal upwelling-favorable winds in the ocean's eastern
16 boundary currents through the 21st century. *Geophysical Research Letters*, **42**, 6424-6431.
17 <http://dx.doi.org/10.1002/2015GL064694>
- 18 Salisbury, J., M. Green, C. Hunt, and J. Campbell, 2008: Coastal acidification by rivers: A threat
19 to shellfish? *Eos, Transactions, American Geophysical Union*, **89**, 513-513.
20 <http://dx.doi.org/10.1029/2008EO500001>
- 21 Schile, L.M., J.C. Callaway, J.T. Morris, D. Stralberg, V.T. Parker, and M. Kelly, 2014:
22 Modeling tidal marsh distribution with sea-level rise: Evaluating the role of vegetation,
23 sediment, and upland habitat in marsh resiliency. *PLOS ONE*, **9**, e88760.
24 <http://dx.doi.org/10.1371/journal.pone.0088760>
- 25 Schmidtko, S., L. Stramma, and M. Visbeck, 2017: Decline in global oceanic oxygen content
26 during the past five decades. *Nature*, **542**, 335-339. <http://dx.doi.org/10.1038/nature21399>
- 27 Schmittner, A., E.D. Galbraith, S.W. Hostetler, T.F. Pedersen, and R. Zhang, 2007: Large
28 fluctuations of dissolved oxygen in the Indian and Pacific oceans during Dansgaard–
29 Oeschger oscillations caused by variations of North Atlantic Deep Water subduction.
30 *Paleoceanography*, **22**, n/a-n/a. <http://dx.doi.org/10.1029/2006PA001384>
- 31 Scott, J.D., Alexander, M.A., Murray, D.R., Swales, D. and Eischeid, J., 2016. The Climate
32 Change Web Portal: a system to access and display climate and Earth system model output
33 from the CMIP5 archive. *Bulletin of the American Meteorological Society*, **97**(4), 523-530.
34 <https://doi.org/10.1175/BAMS-D-15-00035.1>

- 1 Skliris, N., R. Marsh, S.A. Josey, S.A. Good, C. Liu, and R.P. Allan, 2014: Salinity changes in
2 the World Ocean since 1950 in relation to changing surface freshwater fluxes. *Climate*
3 *Dynamics*, **43**, 709-736. <http://dx.doi.org/10.1007/s00382-014-2131-7>
- 4 Smeed, D.A., G.D. McCarthy, S.A. Cunningham, E. Frajka-Williams, D. Rayner, W.E. Johns,
5 C.S. Meinen, M.O. Baringer, B.I. Moat, A. Ducheze, and H.L. Bryden, 2014: Observed
6 decline of the Atlantic meridional overturning circulation 2004–2012. *Ocean Science*, **10**, 29-
7 38. <http://dx.doi.org/10.5194/os-10-29-2014>
- 8 Steinman, B.A., M.E. Mann, and S.K. Miller, 2015: Atlantic and Pacific multidecadal
9 oscillations and Northern Hemisphere temperatures. *Science*, **347**, 988-991.
10 <http://dx.doi.org/10.1126/science.1257856>
- 11 Stendardo, I. and N. Gruber, 2012: Oxygen trends over five decades in the North Atlantic.
12 *Journal of Geophysical Research*, **117**, C11004. <http://dx.doi.org/10.1029/2012JC007909>
- 13 Stramma, L., S. Schmidtke, L.A. Levin, and G.C. Johnson, 2010: Ocean oxygen minima
14 expansions and their biological impacts. *Deep Sea Research Part I: Oceanographic Research*
15 *Papers*, **57**, 587-595. <http://dx.doi.org/10.1016/j.dsr.2010.01.005>
- 16 Straneo, F. and P. Heimbach, 2013: North Atlantic warming and the retreat of Greenland's outlet
17 glaciers. *Nature*, **504**, 36-43. <http://dx.doi.org/10.1038/nature12854>
- 18 Sutton, A.J., C.L. Sabine, R.A. Feely, W.J. Cai, M.F. Cronin, M.J. McPhaden, J.M. Morell, J.A.
19 Newton, J.H. Noh, S.R. Ólafsdóttir, J.E. Salisbury, U. Send, D.C. Vandemark, and R.A.
20 Weller, 2016: Using present-day observations to detect when anthropogenic change forces
21 surface ocean carbonate chemistry outside preindustrial bounds. *Biogeosciences*, **13**, 5065-
22 5083. <http://dx.doi.org/10.5194/bg-13-5065-2016>
- 23 Swanson, K.M., J.Z. Drexler, C.C. Fuller, and D.H. Schoellhamer, 2015: Modeling tidal
24 freshwater marsh sustainability in the Sacramento–San Joaquin delta under a broad suite of
25 potential future scenarios. *San Francisco Estuary and Watershed Science*, **13**, 21.
26 <http://dx.doi.org/10.15447/sfews.2015v13iss1art3>
- 27 Sydeman, W.J., M. García-Reyes, D.S. Schoeman, R.R. Rykaczewski, S.A. Thompson, B.A.
28 Black, and S.J. Bograd, 2014: Climate change and wind intensification in coastal upwelling
29 ecosystems. *Science*, **345**, 77-80. <http://dx.doi.org/10.1126/science.1251635>
- 30 Takatani, Y., D. Sasano, T. Nakano, T. Midorikawa, and M. Ishii, 2012: Decrease of dissolved
31 oxygen after the mid-1980s in the western North Pacific subtropical gyre along the 137°E
32 repeat section. *Global Biogeochemical Cycles*, **26**, n/a-n/a.
33 <http://dx.doi.org/10.1029/2011GB004227>

- 1 Talley, L.D., R.A. Feely, B.M. Sloyan, R. Wanninkhof, M.O. Baringer, J.L. Bullister, C.A.
2 Carlson, S.C. Doney, R.A. Fine, E. Firing, N. Gruber, D.A. Hansell, M. Ishii, G.C. Johnson,
3 K. Katsumata, R.M. Key, M. Kramp, C. Langdon, A.M. Macdonald, J.T. Mathis, E.L.
4 McDonagh, S. Mecking, F.J. Millero, C.W. Mordy, T. Nakano, C.L. Sabine, W.M. Smethie,
5 J.H. Swift, T. Tanhua, A.M. Thurnherr, M.J. Warner, and J.-Z. Zhang, 2016: Changes in
6 ocean heat, carbon content, and ventilation: A review of the first decade of GO-SHIP global
7 repeat hydrography. *Annual Review of Marine Science*, **8**, 185-215.
8 <http://dx.doi.org/10.1146/annurev-marine-052915-100829>
- 9 Taylor, G.T., F.E. Muller-Karger, R.C. Thunell, M.I. Scranton, Y. Astor, R. Varela, L.T.
10 Ghinaglia, L. Lorenzoni, K.A. Fanning, S. Hameed, and O. Doherty, 2012: Ecosystem
11 responses in the southern Caribbean Sea to global climate change. *Proceedings of the*
12 *National Academy of Sciences*, **109**, 19315-19320.
13 <http://dx.doi.org/10.1073/pnas.1207514109>
- 14 Thompson, P.R., B.D. Hamlington, F.W. Landerer, and S. Adhikari, 2016: Are long tide gauge
15 records in the wrong place to measure global mean sea level rise? *Geophysical Research*
16 *Letters*, **43**, 10,403-10,411. <http://dx.doi.org/10.1002/2016GL070552>
- 17 Trenberth, K.E., J.T. Fasullo, and M.A. Balmaseda, 2014: Earth's energy imbalance. *Journal of*
18 *Climate*, **27**, 3129-3144. <http://dx.doi.org/10.1175/jcli-d-13-00294.1>
- 19 Waldbusser, G.G. and J.E. Salisbury, 2014: Ocean acidification in the coastal zone from an
20 organism's perspective: Multiple system parameters, frequency domains, and habitats.
21 *Annual Review of Marine Science*, **6**, 221-247. [http://dx.doi.org/10.1146/annurev-marine-](http://dx.doi.org/10.1146/annurev-marine-121211-172238)
22 [121211-172238](http://dx.doi.org/10.1146/annurev-marine-121211-172238)
- 23 Wallmann, K., 2003: Feedbacks between oceanic redox states and marine productivity: A model
24 perspective focused on benthic phosphorus cycling. *Global Biogeochemical Cycles*, **17**, n/a-
25 n/a. <http://dx.doi.org/10.1029/2002GB001968>
- 26 Wang, D., T.C. Gouhier, B.A. Menge, and A.R. Ganguly, 2015: Intensification and spatial
27 homogenization of coastal upwelling under climate change. *Nature*, **518**, 390-394.
28 <http://dx.doi.org/10.1038/nature14235>
- 29 Wright, J.D. and M.F. Schaller, 2013: Evidence for a rapid release of carbon at the Paleocene-
30 Eocene thermal maximum. *Proceedings of the National Academy of Sciences*, **110**, 15908-
31 15913. <http://dx.doi.org/10.1073/pnas.1309188110>
- 32 Yan, X.-H., T. Boyer, K. Trenberth, T.R. Karl, S.-P. Xie, V. Nieves, K.-K. Tung, and D.
33 Roemmich, 2016: The global warming hiatus: Slowdown or redistribution? *Earth's Future*, **4**,
34 472-482. <http://dx.doi.org/10.1002/2016EF000417>

- 1 Yang, H., G. Lohmann, W. Wei, M. Dima, M. Ionita, and J. Liu, 2016: Intensification and
2 poleward shift of subtropical western boundary currents in a warming climate. *Journal of*
3 *Geophysical Research: Oceans*, **121**, 4928-4945. <http://dx.doi.org/10.1002/2015JC011513>
- 4 Zaitsev, Y.P., 1992: Recent changes in the trophic structure of the Black Sea. *Fisheries*
5 *Oceanography*, **1**, 180-189. <http://dx.doi.org/10.1111/j.1365-2419.1992.tb00036.x>
- 6 Zeebe, R.E., A. Ridgwell, and J.C. Zachos, 2016: Anthropogenic carbon release rate
7 unprecedented during the past 66 million years. *Nature Geoscience*, **9**, 325-329.
8 <http://dx.doi.org/10.1038/ngeo2681>