University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Biological Systems Engineering: Papers and Publications

Biological Systems Engineering

2016

Direct and indirect effects of climatic variations on the interannual variability in net ecosystem exchange across terrestrial ecosystems

Junjiong Shao East China Normal University

Xuhui Zhou East China Normal University, xhzhou@des.ecnu.edu.cn

Yiqi Luo University of Oklahoma

Bo Li Fudan University

Mika Aurela *Einnish Meteorological Institute* Follow this and additional works at: https://digitalcommons.unl.edu/biosysengfacpub

Part of the Bioresource and Agricultural Engineering Commons, Environmental Engineering Commons, See the bage for additional authorial Engineering Commons

Shao, Junjiong; Zhou, Xuhui; Luo, Yiqi; Li, Bo; Aurela, Mika; Billesbach, David; Blanken, Peter D.; Bracho, Rosvel; Chen, Jiquan; Fischer, Marc; Fu, Yuling; Gu, Lianhong; Han, Shijie; He, Yongtao; Kolb, Thomas; Li, Yingnian; Nagy, Zoltan; Niu, Shuli; Oechel, Walter C.; Pinter, Krisztina; Shi, Peili; Suyker, Andrew; Torn, Margaret; Varlagin, Andrej; Wang, Huimin; Yan, Junhua; Yu, Guirui; and Zhang, Junhui, "Direct and indirect effects of climatic variations on the interannual variability in net ecosystem exchange across terrestrial ecosystems" (2016). *Biological Systems Engineering: Papers and Publications*. 610. https://digitalcommons.unl.edu/biosysengfacpub/610

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biological Systems Engineering: Papers and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Junjiong Shao, Xuhui Zhou, Yiqi Luo, Bo Li, Mika Aurela, David Billesbach, Peter D. Blanken, Rosvel Bracho, Jiquan Chen, Marc Fischer, Yuling Fu, Lianhong Gu, Shijie Han, Yongtao He, Thomas Kolb, Yingnian Li, Zoltan Nagy, Shuli Niu, Walter C. Oechel, Krisztina Pinter, Peili Shi, Andrew Suyker, Margaret Torn, Andrej Varlagin, Huimin Wang, Junhua Yan, Guirui Yu, and Junhui Zhang





Tellus B: Chemical and Physical Meteorology

ISSN: (Print) 1600-0889 (Online) Journal homepage: https://www.tandfonline.com/loi/zelb20

Direct and indirect effects of climatic variations on the interannual variability in net ecosystem exchange across terrestrial ecosystems

Junjiong Shao, Xuhui Zhou, Yiqi Luo, Bo Li, Mika Aurela, David Billesbach, Peter D. Blanken, Rosvel Bracho, Jiquan Chen, Marc Fischer, Yuling Fu, Lianhong Gu, Shijie Han, Yongtao He, Thomas Kolb, Yingnian Li, Zoltan Nagy, Shuli Niu, Walter C. Oechel, Krisztina Pinter, Peili Shi, Andrew Suyker, Margaret Torn, Andrej Varlagin, Huimin Wang, Junhua Yan, Guirui Yu & Junhui Zhang

To cite this article: Junjiong Shao, Xuhui Zhou, Yiqi Luo, Bo Li, Mika Aurela, David Billesbach, Peter D. Blanken, Rosvel Bracho, Jiquan Chen, Marc Fischer, Yuling Fu, Lianhong Gu, Shijie Han, Yongtao He, Thomas Kolb, Yingnian Li, Zoltan Nagy, Shuli Niu, Walter C. Oechel, Krisztina Pinter, Peili Shi, Andrew Suyker, Margaret Torn, Andrej Varlagin, Huimin Wang, Junhua Yan, Guirui Yu & Junhui Zhang (2016) Direct and indirect effects of climatic variations on the interannual variability in net ecosystem exchange across terrestrial ecosystems, Tellus B: Chemical and Physical Meteorology, 68:1, 30575, DOI: <u>10.3402/tellusb.v68.30575</u>

To link to this article: https://doi.org/10.3402/tellusb.v68.30575

9	© 2016 J. Shao et al.	+	View supplementary material 🖸
	Published online: 02 Aug 2016.		Submit your article to this journal $arsigma$
111	Article views: 496	CrossMark	View Crossmark data 🗹
	Citing articles: 5 View citing articles 🖸		



Direct and indirect effects of climatic variations on the interannual variability in net ecosystem exchange across terrestrial ecosystems

By JUNJIONG SHAO^{1,2,3}, XUHUI ZHOU^{1,2}*, YIQI LUO⁴, BO LI³, MIKA AURELA⁵, DAVID BILLESBACH⁶, PETER D. BLANKEN⁷, ROSVEL BRACHO⁸, JIQUAN CHEN^{9,10} MARC FISCHER⁶, YULING FU¹¹, LIANHONG GU¹², SHIJIE HAN¹³, YONGTAO HE¹⁴, THOMAS KOLB¹⁵, YINGNIAN LI¹⁶, ZOLTAN NAGY¹⁷, SHULI NIU¹⁴, WALTER C. OECHEL¹⁸, KRISZTINA PINTER¹⁷, PEILI SHI¹⁴, ANDREW SUYKER¹⁹, MARGARET TORN⁶, ANDREJ VARLAGIN²⁰, HUIMIN WANG¹⁴, JUNHUA YAN²¹, GUIRUI YU¹⁴ and JUNHUI ZHANG¹³, ¹School of Ecological and Environmental Sciences, East China Normal University, Shanghai, 200062, China; ²Center for Global Change and Ecological Forecasting, East China Normal University, Shanghai, 200062, China; ³Coastal Ecosystems Research Station of Yangtze River Estuary, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Institute of Biodiversity Science, School of Life Sciences, Fudan University, 2005 Songhu Road, Shanghai, 200438, China; ⁴Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK 73019, USA; ⁵Finnish Meteorological Institute, P.O. Box 503, FI-00101, Helsinki, Finland; ⁶Department of Biological Systems Engineering, University of Nebraska-Lincoln, Lincoln, NE 68583-0726, USA; ⁷Department of Geography, University of Colorado at Boulder, Boulder, CO 80309-0260, USA; ⁸School of Forest Resources and Conservation, University of Florida, Gainesville, Florida, 32611, USA; ⁹Landscape Ecology & Ecosystem Science (LEES) Lab, CGCEO, Michigan State University, East Lansing, MI 48823, USA; ¹⁰Department of Geography, Michigan State University, East Lansing, MI 48823, USA; ¹¹The Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, East China Normal University, Shanghai, 2000241, China; ¹²Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA; ¹³State Key Laboratory of Forest and Soil Ecology, Institute of Applied Ecology, Chinese Academy of Sciences, Liaoning 110016, China; ¹⁴Institute of Geographical Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China; ¹⁵School of Forestry, Northern Arizona University, Flagstaff, AZ 86011-5018, USA; ¹⁶Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Qinghai, 810001, China; ¹⁷MTA-SZIE Plant Ecology Research Group, Szent Istvan University, Pater 1. 2103 Godollo, Hungary; ¹⁸Change Research Group, San Diego State University, San Diego, CA 92182, USA; ¹⁹School of Natural Resources, University of Nebraska–Lincoln, Lincoln, NE 68583, USA; ²⁰A. N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, 119071, Russia; ²¹South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 51065, China

(Manuscript received 30 November 2015; in final form 10 July 2016)

ABSTRACT

Climatic variables not only directly affect the interannual variability (IAV) in net ecosystem exchange of CO_2 (NEE) but also indirectly drive it by changing the physiological parameters. Identifying these direct and indirect paths can reveal the underlying mechanisms of carbon (C) dynamics. In this study, we applied a path analysis using flux data from 65 sites to quantify the direct and indirect climatic effects on IAV in NEE and to

*Corresponding author.

Tellus B 2016. \bigcirc 2016 J. Shao et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material for any purpose, even commercially, provided the original work is properly cited and states its license.

Citation: Tellus B 2016, 68, 30575, http://dx.doi.org/10.3402/tellusb.v68.30575

email: xhzhou@des.ecnu.edu.cn

Responsible Editor: Anders Lindroth, Lund University, Sweden.

evaluate the potential relationships among the climatic variables and physiological parameters that represent physiology and phenology of ecosystems. We found that the maximum photosynthetic rate was the most important factor for the IAV in gross primary productivity (GPP), which was mainly induced by the variation in vapour pressure deficit. For ecosystem respiration (RE), the most important drivers were GPP and the reference respiratory rate. The biome type regulated the direct and indirect paths, with distinctive differences between forests and non-forests, evergreen needleleaf forests and deciduous broadleaf forests, and between grasslands and croplands. Different paths were also found among wet, moist and dry ecosystems. However, the climatic variables can only partly explain the IAV in physiological parameters, suggesting that the latter may also result from other biotic and disturbance factors. In addition, the climatic variables related to NEE were not necessarily the same as those related to GPP and RE, indicating the emerging difficulty encountered when studying the IAV in NEE. Overall, our results highlight the contribution of certain physiological parameters to the IAV in C fluxes and the importance of biome type and multi-year water conditions, which should receive more attention in future experimental and modelling research.

Keywords: net ecosystem exchange, interannual variability, climatic variations, physiological parameters, direct and indirect effects, relative importance

To access the supplementary material to this article, please see <u>Supplementary files</u> under 'Article Tools'.

1. Introduction

During the past century, increased human activities have caused large changes in the global climate system, such as an increase in mean air temperature, a decrease in diurnal temperature range, a redistribution of precipitation and intensification of extreme climate events (IPCC, 2013). Terrestrial ecosystems can not only respond to the climate change, but they also regulate the changing rate of the climate because they absorb 30 % of the anthropogenic CO₂ emissions and determine the interannual fluctuation in the atmospheric CO₂ concentration growth rate (Canadell et al., 2007; Le Quéré et al., 2014; Ahlström et al., 2015). The interannual variability (IAV) in the net ecosystem exchange of CO₂ (NEE) is widely observed in eddy-flux networks, and its drivers have been attributed to a series of climatic and biotic factors (Barr et al., 2007; Pintér et al., 2008; Richardson et al., 2009; Yuan et al., 2009; Dragoni et al., 2011; Humphreys and Lafleur, 2011; Wu et al., 2013). However, model simulations of the IAV in NEE is one of the most urgent and challenging scientific tasks because of the lack of substantial understanding of the intrinsic predictability of IAV in the terrestrial ecosystem carbon (C) cycle (Luo et al., 2015).

The difficulty in predicting the IAV in NEE might result partially from the failure of models to represent the temporal evolution of ecophysiological (e.g. physiological and phenological) factors, which outweigh the climatic factors in most eddy-flux sites at the interannual scale (Shao et al., 2015). Although empirical and semi-empirical research suggests that the ecophysiological properties of ecosystems should be taken into consideration for better simulation of the IAV in C fluxes (Kuzyakov and Gavrichkova, 2010; Migliavacca et al., 2011; Xia et al., 2015), the results from modelling studies indicate that it is still troublesome to predict the temporal changes in the biotic factors or ecophysiological properties of ecosystems (Richardson et al., 2012). Therefore, appropriately representing the dynamics of ecosystem ecophysiology embraces the opportunity to address the issues raised by the IAV in NEE.

The ecophysiological factors that are directly related to the IAV in NEE include photosynthetic and respiratory parameters (hereafter referred as the physiological parameters) that characterize the responses of photosynthesis and respiration to the climatic variations (Shao et al., 2015). Other biotic factors (e.g. biotic disturbance) can only affect the IAV in NEE by changing these parameters. The changes in the physiological parameters can be induced by climatic variations, internal changes in the ecosystem and natural or human disturbances (Richardson et al., 2007; Shao et al., 2015). Among these, the effects of climatic factors on physiological parameters reflect the ecophysiological regulation of ecosystem C cycling by the climatic variations, which occur from the leaf to the ecosystem level (Niu et al., 2012; Wythers et al., 2013; Way and Yamori, 2014; Yamori et al., 2014). At the ecosystem level, a handful of studies have shown that the seasonality of physiological parameters, such as the maximum carboxylation rate (V_{cmax} , maximum velocity of the carboxylase), maximum photosynthetic rate $(A_m, \text{light-saturated photosynthetic rate at the})$ ambient CO₂ concentration), light use efficiency, water use efficiency (WUE), reference respiratory rate at 10 °C (R_{10}) and temperature sensitivity (Q_{10}) were related to radiation, temperature and water conditions in some ecosystems (Aber et al., 1996; Barr et al., 2007; Ricciuto et al., 2008; Yu et al., 2008; Ju et al., 2010; Shao et al., 2014; Shi et al., 2014; Peichl et al., 2015). At a larger temporal scale (interannual), the optimum temperature of NEE, the Michaelis constant of gross primary productivity (GPP), WUE and R_{10} have also been related to the climatic variations (Bahn et al., 2008; Niu et al., 2011, 2012; Shao et al., 2014).

The effects of climate on the IAV in C fluxes mediated by physiological parameters can be considered to be the indirect climatic effects, the strength of which depends on both the correlations between climatic variables and physiological parameters, and the correlations between physiological parameters and C fluxes. If the indirect climatic effects are important contributors to the IAV in C fluxes, the performance of a model can be improved by expressing the physiological parameters as functions of the climatic variables (Ricciuto et al., 2008; Shao et al., 2014). Whether this approach will work or not largely depends on the explicit paths through which the climate directly and indirectly impacts the C fluxes. However, these paths have not been synthetically investigated within eddy-flux networks. To address this issue, we applied a path analysis to 481 site-years of flux data from 65 sites in order to examine the direct and indirect effects of climatic variations on the IAV in NEE and its components (GPP and ecosystem respiration (RE)). Path analysis is a statistical method that is able to quantify the direct and indirect dependencies between or among many variables under consideration (i.e. climatic variables, physiological parameters and C fluxes in this study). The main questions we focused on were: (1) how and to what extent does the climate directly and indirectly affect IAV in C fluxes; and (2) how are these relationships affected by ecosystem characteristics, such as biome types and multi-year water conditions?

2. Materials and methods

2.1. Data sources

The original half-hourly C flux and auxiliary climate data were obtained from AmeriFlux (www.ameriflux.lbl.gov/), CarboEurope (www.europe-fluxdata.eu) and ChinaFLUX (www.chinaflux.org). Only the sites with data for ≥ 5 yr were selected. In total, there were 481 site-years of data belonging to 65 eddy covariance measurement sites from 1992 to 2010 (Table 1, Supplementary Fig. 1). These sites covered a large range of geographical positions and climatic conditions, which can be broadly classified into evergreen needleleaf forests, deciduous broadleaf forests, evergreen broadleaf forests, mixed forests, shrublands, grasslands and croplands (Table 1). The original data included CO2 flux (Fc), friction velocity, photosynthetically active radiation (PAR) or global radiation, air temperature (Ta), soil temperature (Ts), annual precipitation (PPT), relative humidity (RH), vapour pressure deficit (VPD) and latent heat flux (LE).

2.2. Data processing

To obtain the annual values for the C fluxes, climatic variables and physiological parameters, several data processing procedures were applied. The original half-hourly Fc data were first spike-screened by using the median of absolute deviation as a spike estimator based on the double differenced time series (Papale et al., 2006). Next, the nighttime Fc was rejected if the friction velocity was lower than a threshold, which was determined by a 99 % threshold criterion method based on each year's night-time data (Reichstein et al., 2005; Papale et al., 2006). After that, the artificial neural network model with a feed-forward backpropagation algorithm and sigmoid transfer function was applied to partition the NEE into GPP and RE, and to fill the gaps (Shao et al., 2015). The artificial neural network model was trained and validated separately for each year's daytime and night-time data. The input variables of the night-time model were Ts, RH and four seasonal indices, which represented the four seasons (Papale and Valentini, 2003), and the output variable was the night-time RE. The hidden layer had six nodes. The dataset was randomly divided into a training dataset (70 %) and a validation dataset (30 %). This process was repeated 20 times, and the median of 20 simulated outputs was used to fill the gaps in night-time RE and to simulate the daytime RE. The input variables of the daytime model were PAR, Ta, RH and the four seasonal indices, and the output variable was GPP (daytime RE - NEE). Six nodes were set in the hidden layer. The training and validation procedures were repeated 20 times and the median of the simulated GPP was used to fill the GPP gaps. The NEE gaps were filled by the gapfilled RE – GPP. Finally, the gap-filled NEE, GPP and RE, as well as the corresponding climatic variables, were integrated or averaged on an annual basis.

The physiological (photosynthetic and respiratory) parameters used in this study were the maximum photosynthetic rate (A_m), apparent quantum yield (α), WUE, R_{10} and Q_{10} . A_m , α , R_{10} and Q_{10} were derived from the Michaelis–Menten and Q_{10} equations (Falge et al., 2001; Richardson and Hollinger, 2005):

$$GPP = \frac{\alpha \times A_m \times PAR}{A_m + PAR}$$
(1)

$$RE = R_{10} Q_{10}^{\frac{T_5 - 10}{10}}$$
(2)

These equations were parameterised on the basis of 30-d moving windows with a step of 1 d, and only the GPP and RE partitioned from the NEE data were used. The estimated parameters with coefficients of variation larger than 0.5 were excluded (Reichstein et al., 2005) and the median of a certain parameter in a year was considered to be the annual value of the parameter. The WUE is the ratio

Table 1. Information of study sites

Site	Lat (°)	Long (°)	Alt (m)	Eco	n (years)	Ta (°C)	PPT (mm)	References
US-Blo	38.90	-120.63	1315	ENF	9	11.9	1073	Goldstein et al. (2000)
US-Dk3	35.98	-79.09	163	ENF	8	15.0	1050	Stoy et al. (2008)
US-Fmf	35.14	-111.73	2160	ENF	5	9.7	521	Dore et al. (2008)
US-Fuf	35.09	-111.76	2180	ENF	5	8.9	497	Dore et al. (2008)
US-Ho1	45.20	-68.74	60	ENF	13	6.7	840	Richardson et al. (2009)
US-Ho2	45.21	-68.75	91	ENF	11	6.6	842	Xiao et al. (2005)
US-Ho3	45.21	-68.73	61	ENF	5	6.5	877	Davidson et al. (2006)
US-Me2	44.45	-121.56	1253	ENF	9	7.4	441	Thomas et al. (2009)
US-NC2	35.80	-76.67	12	ENF	5	15.8	1212	Noormets et al. (2010)
US-NR1	40.03	-105.55	3050	ENF	10	2.4	663	Blanken et al. (2009)
US-SP3	29.75	-82.16	36	ENF	11	20.1	1034	Powell et al. (2008)
DE-Tha	50.96	13.57	380	ENF	12	7.6	820	Prescher et al. (2010)
DE-Wet	50.45	11.46	703	ENF	7	6.6	967	Rebmann et al. (2010)
ES-ES1	39.35	-0.32	1	ENF	5	17.5	495	Reichstein et al. (2005)
FI-Hvv	61.85	24 29	185	ENF	11	4.8	702	Suni et al. (2003)
FI-Sod	67.36	26.64	189	ENF	7	1.1	425	Aurela (2005)
FR-Pue	43 74	3 60	211	ENF	8	13.6	904	Allard et al. (2008)
IT-Cpz	41 71	12.38	9	ENF	8	15.0	729	Garbulsky et al. (2008)
NI -L oo	52 17	5 74	25	ENF	10	10.3	843	Dolman et al. (2002)
RU-Evo	56.46	32.92	265	ENF	9	5.2	762	Milyukova et al. (2002)
SE-Nor	60.09	17.48	205	ENE	5	6.4	486	Lagergren et al. (2002)
CN-Oia	26.74	115.06	100	ENE	7	18.3	1130	Thang et al. (2003)
US-Bar	44.06	_71.29	272	DRE	5	74	1330	Lenkins et al. (2017)
US-Dk2	35.07	-79.10	168	DBF	7	15.2	1097	Stov et al. (2007)
US-Hal	42 54	-72.17	340	DBF	15	8.5	1153	Urbanski et al. (2007)
US MMS	30.32	86.41	275	DBE	5	12.7	1195	Dragoni et al. (2007)
US MO7	39.32	- 30.41	219 4	DBF	5	12.7	1035	Vang et al. (2010)
	30.01	- 92.20	30	DBF	5	12.0	1033	Hollinger et al. (2010)
US Svy	16.24	- 74.00	540	DBF	5	12.4	696	Desai et al. (2010)
US-Syv	40.24	- 89.55	224		5	4.5	412	Hardiman at al. (2003)
US-UMB	45.50	- 84.71	234 515		0	5.7	412	Cook at al. (2004)
DE U	43.91	- 90.08	515	DBF	/	3.7	/44	
DE-Hai	51.08	10.45	433	DBF	8	8.3	803	Kutsch et al. (2008)
FR-Hes	48.67	/.0/	293	DBF	12	10.4	918	Granier et al. (2000)
IT-Rol	42.41	11.93	153	DBF	1	15.5	769	Rey et al. (2002)
CN-Din	23.17	112.53	240	EBF	6	20.1	1481	Zhang et al. (2010)
BE-Bra	51.31	4.52	15	MF	9	11.4	792	Carrara et al. (2003)
BE-Vie	50.31	6.00	491	MF	11	8.3	719	Aubinet et al. (2001)
CH-Lae	47.48	8.37	645	MF	5	8.3	605	Eugster et al. (2007)
DK-Sor	55.49	11.64	35	MF	13	8.7	656	Pilegaard et al. (2001)
CN-Cha	42.40	128.10	738	MF	6	3.9	498	Guan et al. (2006)
US-Los	46.08	- 89.98	480	SHR	6	3.7	716	Sulman et al. (2009)
MX-Lpa	24.13	-110.44	21	SHR	5	23.4	137	Bell et al. (2012)
US-Aud	31.59	-110.51	1469	GRA	6	15.9	366	Krishnan et al. (2012)
US-Bkg	44.35	-96.84	510	GRA	5	6.3	540	Gilmanov et al. (2005)
US-Dk1	35.97	-79.09	168	GRA	7	15.1	1022	Stoy et al. (2008)
US-FPe	48.31	-105.10	634	GRA	8	5.7	391	Gilmanov et al. (2005)
US-Fwf	35.45	-111.77	2270	GRA	5	8.4	562	Dore et al. (2008)
US-SRM	31.82	-110.87	1116	GRA	7	19.0	324	Scott et al. (2009)
US-Ton	38.43	-120.97	177	GRA	9	16.9	504	Ma et al. (2007)
US-Var	38.41	-120.95	129	GRA	9	15.9	493	Xu and Baldocchi (2004)
US-Wkg	31.74	-109.94	1531	GRA	6	17.2	274	Scott et al. (2010)
AT-Neu	47.12	11.32	1095	GRA	7	6.9	731	Wohlfahrt et al. (2008)
DE-Gri	50.95	13.51	385	GRA	5	7.8	901	Prescher et al. (2010)
DK-Lva	55.68	12.08	15	GRA	5	9.0	655	Gilmanov et al. (2007)

Site	Lat (°)	Long (°)	Alt (m)	Eco	n (years)	Ta (°C)	PPT (mm)	References
HU-Bug	46.69	19.60	106	GRA	6	10.3	499	Nagy et al. (2007)
NL-Cal	51.97	4.93	1	GRA	6	11.0	759	Gilmanov et al. (2007)
CN-Dan	29.67	91.33	4333	GRA	5	2.47	470	Xu et al. (2007)
CN-Hab	37.62	101.28	3220	GRA	6	-0.8	484	Kato et al. (2006)
US-ARM	36.61	-97.49	314	CRO	7	15.3	658	Fischer et al. (2007)
US-Bol	40.01	-88.29	219	CRO	11	11.6	748	Hollinger et al. (2005)
US-Ne1	41.17	-96.48	361	CRO	8	10.7	846	Suyker et al. (2004)
US-Ne2	41.16	-96.47	362	CRO	8	10.2	872	Suyker et al. (2004)
BE-Lon	50.55	4.74	165	CRO	5	11.2	611	Moureaux et al. (2006)
DE-Geb	51.10	10.91	157	CRO	6	9.9	503	Anthoni et al. (2004)
DE-Kli	50.89	13.52	478	CRO	6	7.7	951	Prescher et al. (2010)

Table 1 (Continued)

Lat, latitude; Long, longitude; Alt, altitude; Eco, ecosystem type; n, data record length; Ta, mean annual air temperature; PPT, annual precipitation; ENF, evergreen needleleaf forest; DBF, deciduous broadleaf forest; EBF, evergreen broadleaf forest; MF, mixed forest; SHR, shrubland; GRA, grassland; CRO, cropland.

of GPP to evapotranspiration (ET, calculated from LE [ET $(mm/day) = LE (W/m^2)/28.94$]) (Yuan et al., 2009).

2.3. Path analysis

The dataset used in this study had previously been analysed by Shao et al. (2015) but with different methods and aims. The additive model in Shao et al. (2015) aimed to quantify the relative contribution of climatic and biotic factors to IAV in C fluxes, but it could not identify the paths through which the climate directly and indirectly affected the C fluxes. Therefore, in order to elucidate the direct and indirect effects of climatic variables on the IAV in C fluxes, we applied path analysis, which is an extension of regression model and can deal with multiple casual relationships between or among correlated variables (Shipley, 2004), to the dataset in Shao et al. (2015). Theoretically, path analysis can incorporate nonlinear interactions between climatic variables and C fluxes by using nonlinear functional forms. However, in our study, no nonlinear relationships between variables were found at the interannual scale, though they might exist at shorter temporal scales. The standardised path coefficient (ρ , the path coefficient when all the variables were standardised before the analysis), which is an analogy of the correlation coefficient, was used to quantify the effect size of one variable on another.

Shao et al. (2015) suggested that the correlations between climatic and biotic factors might depend on biome types and multi-year water conditions. Therefore, the study sites were grouped into evergreen needleleaf forests, deciduous broadleaf forests, grasslands and croplands according to their plant function type. In addition, we used the water balance index (WBI), which is the difference between evapotranspiration and precipitation (Law et al., 2002), to group the sites into wet (WBI below – 500 mm), moist

(-500 mm < WBI < 0 mm) and dry (WBI > 0 mm) ecosystems according to their multi-year water conditions. In order to exclude the influence of different magnitudes of the variables among ecosystems, the annual climatic variables, physiological parameters and C fluxes in each site were first standardised by subtracting the mean and then dividing it by the standard deviation of the multi-year data. Path analysis was then applied to each group separately.

For each ecosystem group, two path models were conducted: one for NEE and another for GPP and RE. The use of two models instead of one is because even though some variables are critical to GPP and RE, they may not necessarily be important to NEE. The potential paths in the models mainly included three aspects: the direct effects of climatic variables on C fluxes, the direct effects of climatic variables on physiological parameters and the direct effects of physiological parameters on C fluxes. The first of these three groups were the direct climatic effects; the latter two together characterised the indirect climatic effects on the IAV in C fluxes. In addition, we also considered the regulation of GPP on RE (Supplementary Tables 1 and 2).

The potential paths were based on the following considerations (Supplementary Tables 1 and 2). (1) For the causal paths from climate to C fluxes, five climatic variables (PAR, *Ta*, *Ts*, VPD and WBI) were considered to affect NEE and GPP directly, whereas *Ta*, *Ts*, VPD and WBI affect RE (precipitation was not considered as an explanatory variable because its effects were actually mediated by water availability, which was the opposite of WBI). (2) The IAV in photosynthetic parameters (A_m , α and WUE) may be induced by changes in radiation (PAR), temperature (*Ta*, *Ts*) and water conditions (VPD, WBI), whereas the IAV in respiratory parameters (R_{10} and Q_{10}) might be driven by temperature and water conditions. (3) The photosynthetic and respiratory parameters may influence GPP and RE, respectively, and both kinds of physiological parameter may be important to NEE. (4) The correlation between GPP and RE was found within and across ecosystems (Chen et al., 2015), and may be caused by the substrate supply of photosynthesis to both autotrophic and heterotrophic respiration (Schimel et al., 1994). Therefore, we also assumed that GPP might influence R_{10} . However, autotrophic and heterotrophic respiration might be driven by different factors and correlated differently with GPP, which might result in relationships other than that between GPP and R_{10} . Thus, the control of GPP on RE was also included in the path model. (5) Water availability in autumn and winter might influence the plants' phenology in the next spring and thus was important to the C fluxes in the next year (Gordo and Sanz, 2005). Therefore, the lag effects of autumn-winter WBI on next year's C fluxes and physiological parameters were considered.

The full path models contained too many paths compared to the sample size, which might have caused an overparameterisation problem or may have even failed to derive any statistical tests. Therefore, we simplified the model structure by excluding the paths where the corresponding bivariate correlation coefficients (r) were larger than -0.1 and smaller than 0.1. A path model with the remaining paths was conducted, and the standardised path coefficients (ρ) and the corresponding *p*-values were calculated via maximum likelihood estimation (Beaujean, 2014). To obtain the most parsimonious path model for each ecosystem group, the non-significant paths were excluded recursively by dropping the path with the largest non-significant *p*-value each time and then re-parameterising the new model. This procedure was repeated until all the paths were significant (p < 0.05). In order to make the results among ecosystem groups comparable, we constructed final path models that contained all the significant paths in the most parsimonious models of all the ecosystem groups to be compared. This treatment helped to eliminate the uncertainty caused by variable selection. The path analysis was achieved by using the package lavaan in R software (Rosseel, 2012; R Core Team, 2014).

3. Results

3.1. Direct and indirect climatic effects in different biomes

In all the biomes being studied (evergreen needleleaf forests, deciduous broadleaf forests, grasslands and croplands), the maximum photosynthetic rate (A_m) was the most important factor for the IAV in C fluxes (the standardised path coefficient (ρ) as an analogy of the correlation coefficient fell between -0.32 and -0.68 for NEE and 0.72-0.85 for

GPP), which was mainly induced by the variation in VPD (Figs. 1 and 2). For the IAV in RE, the most important drivers were GPP ($\rho = 0.47-0.67$) and reference respiratory rate (R_{10} , $\rho = 0.24-0.29$ except for croplands; Fig. 2). Though we also considered the lag effects of autumn–winter WBI on the physiological parameters and C fluxes in the next year (Supplementary Table 3), they were not selected in the final path models.

There were also different direct and indirect paths from climate to C fluxes among biomes. For example, a positive correlation between temperature and apparent quantum yield (α) , as well as a negative correlation between temperature sensitivity (Q_{10}) and RE, was only found in forests, while the controls of GPP on reference respiratory rate (R_{10}) only appeared in non-forests (Fig. 2). Compared with deciduous broadleaf forests, the evergreen needleleaf forests showed direct climatic effects on C fluxes (PAR on GPP, and Ts on RE), while there were more significant indirect climatic effects in deciduous broadleaf forests than in evergreen needleleaf forests (Fig. 2a and b). Although PAR influenced the maximum photosynthetic rate in both grasslands $(\rho = -0.39)$ and croplands $(\rho = 0.34)$ within non-forests, it did so in opposite directions. In addition, the indirect climatic effects on RE in grasslands were not found in croplands (Fig. 2c and d).

3.2. Direct and indirect climatic effects in ecosystems with different aridity

The direct and indirect effects of climate on IAV in C fluxes were also different among ecosystems with different multiyear water conditions (Fig. 3). Overall, the moist ecosystems had more complicated direct and indirect paths than wet and dry ecosystems. Compared with wet ecosystems, the IAV in GPP of moist and dry ecosystems can be induced by apparent quantum yield (α , an indicator of light use efficiency, $\rho = 0.08$ and 0.17, respectively) and WUE ($\rho = 0.13$ and 0.24, respectively), and the IAV in RE can be induced by reference respiratory rate (R_{10} , $\rho = 0.31$ and 0.25, respectively). On the other hand, direct effects of WBI on RE were found in moist ($\rho = -0.10$) and dry ($\rho = -0.18$), but not wet ecosystems.

3.3. Model performance

The results derived from the NEE model were generally consistent with those from the GPP and RE model, but with some discrepancies (Figs. 1–3). For example, the complicated indirect climatic effects in grasslands seen in the GPP and RE model (Fig. 2c) were largely simplified in the NEE model (Fig. 1c). In the dry ecosystems, Ts emerged to directly influence NEE, though it did not affect



Fig. 1. The direct and indirect effects of climatic variation on the interannual variability (IAV) in net ecosystem exchange (NEE) for evergreen needleleaf forests (a), deciduous broadleaf forests (b), grasslands (c) and croplands (d). Green and red lines represent positive and negative effects, respectively. Solid and dashed lines represent significant (p < 0.05) and non-significant ($p \ge 0.05$) effects, respectively. The values beside the paths are the standardised (0–1) path coefficients (ρ), which are only shown for significant effects. PAR, photosynthetically active radiation; *T*a, air temperature; *Ts*, soil temperature; VPD, vapour pressure deficit; WBI, water balance index; A_m , maximum photosynthetic rate; α , apparent quantum yield; WUE, water use efficiency; R_{10} , reference respiratory rate at 10 °C; Q_{10} , temperature sensitivity.

GPP or RE (Fig. 3e and f). Overall, the path models were better for explaining the IAV in GPP ($r^2 = 0.54-0.81$) and RE ($r^2 = 0.36-0.60$) than that in NEE ($r^2 = 0.17-0.61$; Figs. 1–3). Among ecosystem groups, the model performed best in grasslands and croplands (the r^2 values of C fluxes were all above 0.5) and worst in evergreen needleleaf forests (the r^2 values of NEE were below 0.2). The climatic variables explained the variations in the physiological parameters best in grasslands, with the r^2 of maximum photosynthetic rate being 0.31. In other ecosystems, climatic variations could only induce very small levels of the IAV in the physiological parameters (Figs. 1–3).

Because the physiological parameters were not directly measured but derived from equations, the potential correlations among them might cause some uncertainty. Although significant correlations were found in many pairs of parameters, the coefficients were generally very small, except for those between maximum photosynthetic rate and WUE in grasslands and croplands and dry ecosystems (r > 0.4, Table 2). In addition, there was no significant correlation between reference respiratory rate and temperature sensitivity (Table 2).

4. Discussion

4.1. Regulation of biome type

The climatic variations can drive the IAV in C fluxes directly and indirectly by changing the physiological parameters, with both shared and unique paths among ecosystem types. In all the biome types being studied (i.e. evergreen needleleaf forests, deciduous broadleaf forests, grasslands and croplands), the importance of maximum photosynthetic rate (A_m) was highlighted (Figs. 1 and 2), a result that has also been found in a tundra and a fen (Humphreys and Lafleur, 2011). Xia et al. (2015) also showed that the seasonal maximal photosynthetic capacity was the best indicator of IAV in GPP for all the biomes. The control of VPD on A_m at



Fig. 2. The direct and indirect effects of climatic variation on in the interannual variability (IAV) in gross primary productivity (GPP) and ecosystem respiration (RE) for evergreen needleleaf forests (a), deciduous broadleaf forests (b), grasslands (c) and croplands (d). Green and red lines represent positive and negative effects, respectively. Solid and dashed lines represent significant (p < 0.05) and non-significant ($p \ge 0.05$) effects, respectively. The values beside the paths are the standardised (0–1) path coefficients (ρ), which are only shown for significant effects. PAR, photosynthetically active radiation; *Ta*, air temperature; *Ts*, soil temperature; VPD, vapour pressure deficit; WBI, water balance index; $A_{\rm m}$, maximum photosynthetic rate; α , apparent quantum yield; WUE, water use efficiency; R_{10} , reference respiratory rate at 10 °C; Q_{10} , temperature sensitivity.

the interannual scale (Figs. 1 and 2) reflected the control of water stress on stomatal conductance (Medlyn et al., 2001). For the IAV in RE, there was no consistent driver across all the biome types except for the strong control of GPP (Fig. 2). The tight correlation between GPP and RE might be derived from the influence of GPP on the components of RE: autotrophic and heterotrophic respiration. The relationship between GPP and autotrophic respiration was mainly caused by the proportional changes in growth respiration to GPP (Amthor, 2000). The heterotrophic respiration can be correlated with GPP because the microbes prefer to decompose the younger organic matter derived from more recent photosynthate (Schimel et al., 1994).

However, there were some distinctive differences between forests and non-forests (Fig. 2). For example, in forests, the warmer environment stimulated GPP by promoting the apparent quantum yield (α), which is physiologically defined as the ratio of photosynthesis to the absorbed PAR and is equivalent to the concept of light use efficiency, which has been applied in light use efficiency models (Johnson and Barber, 2003). This might be because the enzyme activity or leaf area index was enhanced in warmer environments (Shi et al., 2014; Gitelson and Gamon, 2015). The negative correlation between temperature sensitivity (Q_{10}) and RE found in forests seemed to contradict common sense. However, previous studies confirmed that Q_{10} generally declined with the increase in temperature (Tjoelker et al., 2001), which, in turn, indicated a negative relationship between Q_{10} and RE because both autotrophic and heterotrophic respiration were positively correlated with temperature.

Compared with forests, the non-forest ecosystems showed the control of GPP on reference respiratory rate (R_{10}). This might be caused by the different transportation rates of photosynthetic production between forests and non-forests. The carbohydrate synthesised in leaves usually takes a couple of days to arrive at the roots in trees but the duration is shorter than 1 d in herbs (Högberg et al., 2001; Högberg and Read, 2006; Kuzyakov and Gavrichkova, 2010). In addition, the maintenance respiration might account for the



Fig. 3. The direct and indirect effects of climatic variation on the interannual variability (IAV) in net ecosystem exchange (NEE, a–c), gross primary productivity (GPP) and ecosystem respiration (RE, d–f) for wet (water balance index, or WBI, below – 500 mm; a, d), moist (-500 mm < WBI < 0 mm; b, e), and dry (WBI > 0 mm; c, f) ecosystems. Note that a larger WBI indicates a drier environment. Green and red lines represent positive and negative effects, respectively. Solid and dashed lines represent significant (p < 0.05) and non-significant ($p \ge 0.05$) effects, respectively. The values beside the paths are the standardised (0–1) path coefficients (ρ), which are only shown for significant effects. PAR, photosynthetically active radiation; *T*a, air temperature; *Ts*, soil temperature; VPD, vapour pressure deficit; WBI, water balance index; A_m , maximum photosynthetic rate; α , apparent quantum yield; WUE, water use efficiency; R_{10} , reference respiratory rate at 10 °C; Q_{10} , temperature sensitivity.

larger proportion of autotrophic respiration in trees compared with herbs, which will weaken the relationships between photosynthesis and respiration (DeLucia et al., 2007).

Previous studies have suggested that the deciduous broadleaf forests are more sensitive to climatic variations than evergreen needleleaf forests (Welp et al., 2007; Yuan et al., 2009). Our results showed that the C fluxes in deciduous broadleaf forests were more sensitive to indirect climatic effects than those in evergreen needleleaf forests, while there were direct climatic effects on GPP and RE in evergreen needleleaf forests but not in deciduous broadleaf forests (Fig. 2a and b). These differences might be due to the different leaf life spans, phenology and the environment

Table 2. Correlation coefficients (*r*) of the physiological parameters in different ecosystem types

Ecosystem group	$r(A_m, \alpha)$	$r(A_m, WUE)$	$r(\alpha, WUE)$	$r(R_{10}, Q_{10})$
ENF	0.21*	0.10	0.11	0.14
DBF	0.10	0.11	0.28*	0.07
GRA	0.29*	0.41*	0.27*	0.07
CRO	0.02	0.56*	0.10	0.07
Wet	0.05	0.11	0.02	0.10
Moist	0.15*	0.29*	0.19*	0.10
Dry	0.09	0.49*	0.09	0.14

ENF, evergreen needleleaf forests; DBF, deciduous broadleaf forests; GRA, grasslands; CRO, croplands; Wet, ecosystems with a water balance index (WBI) below -500 mm; Moist, ecosystems with -500 mm < WBI < 0 mm; Dry, ecosystems with WBI > 0 mm; A_{m} , maximum photosynthetic rate; α , apparent quantum yield; WUE, water use efficiency; R_{10} , reference respiratory rate at 10 °C; Q_{10} , temperature sensitivity.

*p < 0.05.

being adapted to. Compared to deciduous broadleaf forests, evergreen needleleaf forests are usually with longer foliage longevity, lower specific leaf area and are adapted to unfavourable environments, which make the nutrient (e.g. nitrogen and phosphorus) concentrations and thus the photosynthetic parameters more constant in a changing environment (Bloom et al., 1985; Givnish, 2002). On the other hand, the foliage nutrient concentrations would be more variable and correlated with climatic variables in deciduous broadleaf forests, as the leaf development is largely controlled by climate (Polgar and Primack, 2011). Likewise, the annual litter fall with varying litter quality (e.g. C: N) in deciduous broadleaf forests might contribute to the indirect climatic effects on RE.

The most significant difference between grasslands and croplands was that the effect of PAR on maximum photosynthetic rate (A_m) was negative in the former but positive in the latter (Fig. 2c and d). In grasslands, the relatively low productivity and high water stress resulted in excess energy when the radiation increased, damaging the photosynthetic system and reducing A_m (Schulze et al., 2005). In contrast, high productivity and irrigation allowed the croplands to enhance A_m when the radiation increased. Another difference was that the indirect climatic effects on RE that occurred in grasslands was not found in croplands, which might be because the management (i.e. irrigation, fertilisation) in croplands hampered the influence of climate.

4.2. Regulation of multi-year water conditions

Our previous study suggested that the multi-year water conditions might be the most important regulator of the biotic responses to climatic variation (Shao et al., 2015). Current study also showed that the direct and indirect effects of climate on IAV in C fluxes were different among ecosystem groups with different aridity (Fig. 3). For example, WUE and apparent quantum yield (α , an indicator of light use efficiency) had significant effects on GPP in moist and dry, but not wet ecosystems. The discrepancy in importance of WUE among different ecosystem groups might be due to the different partitioning of evapotranspiration into evaporation and transpiration. The WUE in wet ecosystems was generally larger than that in other ecosystems due to the stronger stimulation of GPP compared to evapotranspiration when water availability increased (Xiao et al., 2013; Xue et al., 2015; Zhu et al., 2015). Higher WUE means that relatively more water is lost through transpiration, which is tightly coupled with photosynthesis. Therefore, the variation of WUE in wet ecosystems is smaller than that in other ecosystems, which ultimately results in the nonsignificant relationship between WUE and GPP. Likewise, light use efficiency also reaches its maximum when there is no water stress (Yuan et al., 2014), and this results in a non-significant relationship between light use efficiency and GPP. Interestingly, the water condition (i.e. VPD) was important to IAV in C fluxes, even for wet ecosystems (Fig. 3b). Due to flux partitioning and gap-filling, VPD can be related to C fluxes because it was influenced by air temperature and relative humidity. These are input variables of the artificial neural network model, and our results showed that it had little direct effect on C fluxes. This suggests that the effects of VPD on GPP in wet ecosystems were not a by-product of data processing, but a reflection of the underlying causal links. This might be due to a seasonal dry environment or an extreme drought year occurring in wet ecosystems, which can highlight the importance of water conditions (Tian et al., 2000; Lewis et al., 2011).

The moist ecosystems had more complicated indirect paths than wet and dry ecosystems, which might be because the former had a longer time lag (8-10 months) between the water balance and C fluxes compared with the latter ones (Vicente-Serrano et al., 2013). This lag effect might reflect the influence of autumn-winter precipitation stored in soil on the C fluxes of the following year. In order to investigate this issue, we added the effects of autumnwinter WBI on next year's physiological parameters and C fluxes. Although some significant correlations were found (Supplementary Table 3), none of these lag effects were selected in the final path models, suggesting that the lagged climatic effects were much weaker than the concurrent ones. Other studies found even longer lags (2-4 yr) between water conditions and C fluxes (Dunn et al., 2007; Ito, 2011). However, it is difficult to evaluate these longterm lags because of insufficient data.

4.3. Uncertainty and implications

Our study showed how climatic variations may directly and indirectly regulate the IAV in C fluxes in different biomes and ecosystems with different water conditions. However, there might be some uncertain patterns in the results. Overall, the explanatory ability of the path models was best for the IAV in GPP, followed by those in RE and NEE (Figs. 1-3). The difficulty of simulating IAV in RE and NEE might be due to the complex responses of both fluxes to climatic variations. For example, RE consists of autotrophic and heterotrophic respiration and each of these components includes multiple processes with different responses to environmental changes (Kuzyakov and Gavrichkova, 2010; Thornley, 2011; Zhou et al., 2014). Likewise, NEE is the difference between RE and GPP, whose responses to climate are quite different (Anderson-Teixeira et al., 2011; Shi et al., 2014).

Another source of uncertainty might be the autocorrelations between variables because they were not independently observed but were derived from a few observed variables (e.g. NEE and climatic variables). The correlation between GPP and RE was thought to be spurious because these were partitioned from NEE data (Vickers et al., 2009). However, recent studies suggested that this spurious correlation was relatively small (Lasslop et al., 2010; Baldocchi et al., 2015). The physiological parameters were also not independent because they were derived from the Michaelis-Menten and Q_{10} equations. Fortunately, these correlations were not strong enough to introduce large uncertainty into the results (Table 2). Strictly, the physiological parameters might be just byproducts of parameterisation process, which would lead to spurious relations in the path models. In this situation, there should be stronger relationships between climatic variables and C fluxes than those between climate and physiological parameters. However, we found that climatic variables were more highly correlated with physiological parameters rather than C fluxes, indicating that there was no obvious spurious correlation problem. Furthermore, it is conceptually reasonable to treat the physiological parameters as drivers rather than by-products of parameterisation because the physiological parameters determine the capacities of photosynthesis and respiration, which are also widely adopted in ecosystem models. In addition, although some relationships between climatic variables, physiological parameters and C fluxes could be expected (e.g. the relationship between PAR and photosynthetic parameters, and that between temperature and RE) due to the data processing procedure, they did not appear in all the ecosystem groups, indicating that the methodological uncertainty was not the main source of the correlations between variables.

Nevertheless, some implications and suggestions can be made for future studies. First, the weak correlations between climatic variables and physiological parameters may reflect the fact that NEE is not a biochemical process, so it is difficult to characterise its biotic drivers, or may indicate that the internal changes of ecosystems, and natural or human disturbance might be the primary drivers of the IAV in physiological parameters. These drivers may include the morphology, biochemistry and phenology of leaves, soil properties, nutrient availability, vegetation characteristics and human management (Buchmann and Schulze, 1999; Cook et al., 2008; de Beeck et al., 2010; Zhou et al., 2013; Madani et al., 2014; Duchemin et al., 2015; Gitelson and Gamon, 2015; Yan et al., 2015; Zhu et al., 2015). However, great caution should be applied when extrapolating findings from the seasonal scale up to the interannual scale, because the relationships between climate and physiological parameters might change across temporal scales. For example, Tan et al. (2015) found that VPD dominated WUE in a tropical rainforest at the instantaneous scale but became the second important factor at the seasonal scale and was not correlated to WUE at the interannual scale.

Second, the patterns of the direct and indirect climatic effects on the IAV in C fluxes differed among ecosystem types. However, identifying the different underlying mechanisms of C cycle dynamics and evaluating their importance to the IAV in C fluxes will require quantitative data describing the important features of the different ecosystems. For example, tree growth might be an important aspect for forest ecosystems and leaf phenology might regulate photosynthesis in deciduous broadleaf forests (Desai et al., 2005; Dragoni et al., 2011). Species composition may shift in the grasslands across several years (Kahmen et al., 2002), whereas human management regimens are critical to croplands (Lobell et al., 2006). In order to incorporate this information into C cycle models, longterm observations should be carried out and appropriate procedures that can better weigh the importance of these data should be developed in future studies.

5. Acknowledgements

This study was financially supported by the National Natural Science Foundation of China (Grant No. 31370489), the Program for Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning and the 'Thousand Young Talents' Program in China. The authors also thank Marc Aubinet, Dennis Baldocchi, Christian Bernhofer, Gil Bohrer, Paul Bolstad, Nina Buchmann, Alexander Cernusca, Reinhart Ceulemans, Kenneth Clark, Ankur Desai, Allen Goldstein, Andre Granier, David Hollinger, Gabriel Katul, Alexander Knohl, Werner Kutsch, Beverly Law, Anders Lindroth, Timothy Martin, Tilden Meyers, Eddy Moors, William Munger, Kimberly Novick, Ram Oren, Serge Rambal, Corinna Rebmann, Andrew Richardson, Maria-Jose Sanz, Russ Scott, Lise Soerensen, Riccardo Valentini and Timo Vesala for their generous permission to access the flux data.

References

- Aber, J. D., Reich, P. B. and Goulden, M. L. 1996. Extrapolating leaf CO₂ exchange to the canopy: a generalized model of forest photosynthesis compared with measurements by eddy correlation. *Oecologia*. **106**, 257–265.
- Ahlström, A., Raupach, M. R., Schurgers, G., Smith, B., Arneth, A. and co-authors. 2015. The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink. *Science*. 348, 895–899.
- Allard, V., Ourcival, J. M., Rambal, S., Joffre, R. and Rocheteau, A. 2008. Seasonal and annual variation of carbon exchange in an evergreen Mediterranean forest in southern France. *Global Change Biol.* 14, 714–725.
- Amthor, J. S. 2000. The McCree-de Wit-Penning de Vries-Thornley respiration paradigms: 30 years later. Ann. Bot. 86, 1–20.
- Anderson-Teixeira, K. J., Delong, J. P., Fox, A. M., Brese, D. A. and Litvak, M. E. 2011. Differential responses of production and respiration to temperature and moisture drive the carbon balance across a climatic gradient in New Mexico. *Global Change Biol.* 17, 410–424.
- Anthoni, P. M., Knohl, A., Rebmann, C., Freibauer, A., Mund, M. and co-authors. 2004. Forest and agricultural land-use-dependent CO₂ exchange in Thuringia, Germany. *Global Change Biol.* 10, 2005–2019.
- Aubinet, M., Chermanne, B., Vandenhaute, M., Longdoz, B., Yernaux, M. and co-authors. 2001. Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes. *Agric. Forest Meteorol.* 108, 293–315.
- Aurela, M. 2005. Carbon-Dioxide Exchange in Subarctic Ecosystems Measured by a Micrometeorological Technique. PhD Thesis. Finnish Meteorological Institute, Helsinki, Finland.
- Bahn, M., Rodeghiero, M., Anderson-Dunn, M., Dore, S., Gimeno, C. and co-authors. 2008. Soil respiration in European grasslands in relation to climate and assimilate supply. *Ecosystems.* 11, 1352–1367.
- Baldocchi, D., Sturtevant, C. and FLUXNET Contributors. 2015. Does day and night sampling reduce spurious correlation between canopy photosynthesis and ecosystem respiration? *Agric. Forest Meteorol.* 207, 117–126.
- Barr, A. G., Black, T. A., Hogg, E. H., Griffis, T. J., Morgenstern, K. and co-authors. 2007. Climatic controls on the carbon and water balances of a boreal aspen forest, 1994–2003. *Global Change Biol.* 13, 561–576.

- Beaujean, A. A. 2014. Latent Variable Modelling Using R: A Stepby-Step Guide. Routledge, New York.
- Bell, T. W., Menzer, O., Troyo-Diéquez, E. and Oechel, W. C. 2012. Carbon dioxide exchange over multiple temporal scales in an arid shrub ecosystem near La Paz, Baja California Sur, Mexico. *Global Change Biol.* 18, 2570–2582.
- Blanken, P. D., Williams, M. W., Burns, S. P., Monson, R. K., Knowles, J. and co-authors. 2009. A comparison of water and carbon dioxide exchange at a windy alpine tundra and subalpine forest site near Niwot Ridge, Colorado. *Biogeochemistry*. 95, 61–76.
- Bloom, A. J., Chapin, F. S. and Mooney, H. A. 1985. Resource limitation in plants: an economic analogy. *Ann. Rev. Ecol. Syst.* 16, 363–392.
- Buchmann, N. and Schulze, E.-D. 1999. Net CO₂ and H₂O fluxes of terrestrial ecosystem. *Global Biogeochem. Cycles.* 13, 751– 760.
- Canadell, J. G., Le Quéré, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T. and co-authors. 2007. Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks. *Proc. Natl. Acad. Sci. USA*. **104**, 18866–18870.
- Carrara, A., Kowalski, A. S., Neirynck, J., Janssens, I. A., Yuste, J. C. and co-authors. 2003. Net ecosystem CO₂ exchange of mixed forest in Belgium over 5 years. *Agric. Forest Meteorol.* 119, 209–227.
- Chen, Z., Yu, G., Zhu, X., Wang, Q., Niu, S. and co-authors. 2015. Covariation between gross primary production and ecosystem respiration across space and the underlying mechanisms: a global synthesis. *Agric. Forest Meteorol.* 203, 180–190.
- Cook, B. D., Bolstad, P. V., Martin, J. G., Heinsch, F. A., Davis, K. J. and co-authors. 2008. Using light-use and production efficiency models to predict photosynthesis and net carbon exchange during forest canopy disturbance. *Ecosystems*. 11, 26– 44.
- Cook, B. D., Davis, K. J., Wang, W., Desai, A., Berger, B. W. and co-authors. 2004. Carbon exchange and venting anomalies in an upland deciduous forest in northern Wisconsin, USA. *Agric. Forest Meteorol.* **126**, 271–295.
- Davidson, E. A., Richardson, A. D., Savage, K. E. and Hollinger, D. Y. 2006. A distinct seasonal pattern of the ratio of soil respiration to total ecosystem respiration in a spruce-dominated forest. *Global Change Biol.* **12**, 230–239.
- de Beeck, M. O., Gielen, B., Jonckheere, I., Samson, R., Janssens, I. A. and co-authors. 2010. Needle age-related and seasonal photosynthetic capacity variation is negligible for modelling yearly gas exchange of a sparse temperate Scots pine forest. *Biogeosciences*. 7, 199–215.
- DeLucia, E. H., Drake, J. E., Thomas, R. B. and Gonzalez-Meler, M. 2007. Forest carbon use efficiency: is respiration a constant fraction of gross primary production? *Global Change Biol.* 13, 1157–1167.
- Desai, A. R., Bolstad, P. V., Cook, B. D., Davis, K. J. and Carey, E. V. 2005. Comparing net ecosystem exchange of carbon dioxide between an old-growth and mature forest in the upper Midwest, USA. Agric. Forest Meteorol. 128, 33–55.

- Dolman, A. J., Moors, E. J. and Elbers, J. A. 2002. The carbon uptake of a mid latitude pine forest growing on sandy soil. *Agric. Forest Meteorol.* 111, 157–170.
- Dore, S., Kolb, T. E., Montes-Helu, M., Sullivan, B. W., Winslow, W. D. and co-authors. 2008. Long-term impact of a standreplacing fire on ecosystem CO₂ exchange of a *Ponderosa* pine forest. *Global Change Biol.* 14, 1801–1820.
- Dragoni, D., Schmid, H. P., Wayson, C. A., Potter, H., Grimmond, C. S. B. and co-authors. 2011. Evidence of increased net ecosystem productivity associated with a longer vegetated season in a deciduous forest in south-central Indiana, USA. *Global Change Biol.* 17, 886–897.
- Duchemin, B., Fieuzal, R., Rivera, M., Ezzahar, J., Jarlan, L. and co-authors. 2015. Impact of sowing date on yield and water use efficiency of wheat analyzed through spatial modelling and FORMOSAT-2 images. *Remote Sens.* 7, 5951–5979.
- Dunn, A., Barford, C. C., Wofsy, S. C., Goulden, M. L. and Daube, B. C. 2007. A long-term record of carbon exchange in a boreal black spruce forest: means, responses to interannual variability, and decadal trends. *Global Change Biol.* 13, 577–590.
- Eugster, W., Zeyer, K., Zeeman, M., Michna, P., Zingg, A. and co-authors. 2007. Methodical study of nitrous oxide eddy covariance measurements using quantum cascade laser spectrometery over a Swiss forest. *Biogeosciences.* 4, 927–939.
- Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M. and co-authors. 2001. Gap filling strategies for defensible annual sums of net ecosystem exchange. *Agric. Forest Meteorol.* **107**, 43–69.
- Fischer, M. L., Billesbach, D. P., Berry, J. A., Riley, W. J. and Torn, M. S. 2007. Spatiotemporal variations in growing season exchanges of CO₂, H₂O, and sensible heat in agricultural fields of the Southern Great Plains. *Earth Interact.* **11**, 1–21.
- Garbulsky, M. F., Peñuelas, J., Papale, D. and Filella, I. 2008. Remote estimation of carbon dioxide uptake by a Mediterranean forest. *Global Change Biol.* 14, 2860–2867.
- Gilmanov, T. G., Soussana, J. F., Aires, L., Allard, V., Ammann, C. and co-authors. 2007. Partitioning European grassland net ecosystem CO₂ exchange into gross primary productivity and ecosystem respiration using light response function analysis. *Agric. Ecosyst. Environ.* **121**, 93–120.
- Gilmanov, T. G., Tieszen, L. L., Wylie, B. K., Flanagan, L. B., Frank, A. B. and co-authors. 2005. Integration of CO₂ flux and remotely-sensed data for primary production and ecosystem respiration analyses in the Northern Great Plains: potential for quantitative spatial extrapolation. *Global Ecol. Biogeogr.* 14, 271–292.
- Gitelson, A. A. and Gamon, J. A. 2015. The need for a common basis for defining light-use efficiency: implications for productivity estimation. *Remote Sens. Environ.* **156**, 196–201.
- Givnish, T. J. 2002. Adaptive significance of evergreen vs. deciduous leaves: solving the triple paradox. *Silva Fenn.* **36**, 703–743.
- Goldstein, A. H., Hultman, N. E., Fracheboud, J. M., Bauer, M. R., Panek, J. A. and co-authors. 2000. Effects of climate variability on the carbon dioxide, water, and sensible heat fluxes above a *Ponderosa* pine plantation in the Sierra Nevada (CA). *Agric. Forest Meteorol.* 101, 113–129.

- Gordo, O. and Sanz, J. J. 2005. Phenology and climate change: a long-term study in a Mediterranean locality. *Global Change Ecol.* **146**, 484–495.
- Granier, A., Ceschia, E., Damesin, C., Dufrene, E., Epron, D. and co-authors. 2000. The carbon balance of a young beech forest. *Funct. Ecol.* 14, 312–325.
- Guan, D., Wu, J., Zhao, X., Han, S., Yu, G. and co-authors. 2006. CO₂ fluxes over an old, temperate mixed forest in northeastern China. *Agric. Forest Meteorol.* **137**, 138–149.
- Hardiman, B. S., Bohrer, G., Gough, C. M., Vogel, C. S. and Curtis, P. S. 2011. The role of canopy structural complexity in wood net primary production of a maturing northern deciduous forest. *Ecology*, **92**, 1818–1827.
- Högberg, P., Nordgren, A., Buchmann, N., Taylor, A. F. S., Ekblad, A. and co-authors. 2001. Large-scale forest girdling shows that current photosynthesis drives soil respiration. *Nature*. 411, 789–794.
- Högberg, P. and Read, D. J. 2006. Towards a more plant physiological perspective on soil ecology. *Trends Ecol. Evol.* 21, 548–554.
- Hollinger, D. Y., Ollinger, S. V., Richardson, A. D., Meyers, T. P., Dail, D. B. and co-authors. 2010. Albedo estimates for land surface models and support for a new paradigm based on foliage nitrogen concentration. *Global Change Biol.* 16, 696–710.
- Hollinger, S. E., Bernacchi, C. J. and Meyers, T. P. 2005. Carbon budget of mature no-till ecosystem in North Central Region of the United States. *Agric. Forest Meteorol.* **130**, 59–69.
- Humphreys, E. R. and Lafleur, P. M. 2011. Does earlier snowmelt lead to greater CO₂ sequestration in two low Arctic tundra ecosystems? *Geophys. Res. Lett.* 38, L09703. DOI: http://dx.doi. org/10.1029/2011GL047339
- IPCC. 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK, pp. 56–57.
- Ito, A. 2011. Decadal variability in the terrestrial carbon budget caused by the Pacific Decadal Oscillation and Atlantic Multidecadal Oscillation. J. Meteorol. Soc. Jpn. 89, 441–454.
- Jenkins, J. P., Richardson, A. D., Braswell, B. H., Ollinger, S. V., Hollinger, D. Y. and co-authors. 2007. Refining light-use efficiency calculations for a deciduous forest canopy using simultaneous tower-based carbon flux and radiometric measurements. *Agric. Forest Meteorol.* 143, 64–79.
- Johnson, Z. and Barber, R. T. 2003. The low-light reduction in the quantum yield of photosynthesis: potential errors and biases when calculating the maximum quantum yield. *Photosynth. Res.* 75, 85–95.
- Ju, W., Wang, S., Yu, G., Zhou, Y. and Wang, H. 2010. Modeling the impact of drought on canopy carbon and water fluxes for a subtropical evergreen coniferous plantation in southern China through parameter optimization using an ensemble Kalman filter. *Biogeosciences.* 7, 845–857.
- Kahmen, S., Poschlod, P. and Schreiber, K.-F. 2002. Conservation management of calcareous grasslands. Changes in plant species composition and response of functional traits during 25 years. *Biol. Conserv.* 104, 319–328.

- Kato, T., Tang, Y., Gu, S., Hirota, M., Du, M. and co-authors. 2006. Temperature and biomass influences on interannual changes in CO₂ exchange in an alpine meadow on the Qinghai–Tibetan Plateau. *Global Change Biol.* **12**, 1285–1298.
- Krishnan, P., Meyers, T. P., Scott, R. L., Kennedy, L. and Heuer, M. 2012. Energy exchange and evapotranspiration over two temperate semi-arid grasslands in North America. *Agric. Forest Meteorol.* 153, 31–44.
- Kutsch, W. L., Kolle, O., Rebmann, C., Knohl, A., Ziegler, W. and co-authors. 2008. Advection and resulting CO₂ exchange uncertainty in a tall forest in central Germany. *Ecol. Appl.* 18, 1391–1405.
- Kuzyakov, Y. and Gavrichkova, O. 2010. Time lag between photosynthesis and carbon dioxide efflux from soil: a review of mechanisms and controls. *Global Change Biol.* 16, 3386–3406.
- Lagergren, F., Eklundh, L., Grelle, A., Lundblad, M., Molder, M. and co-authors. 2005. Net primary production and light use efficiency in a mixed coniferous forest in Sweden. *Plant Cell Environ.* 28, 412–423.
- Lasslop, G., Reichstein, M., Papale, D., Richardson, A. D., Arneth, A. and co-authors. 2010. Separation of net ecosystem exchange into assimilation and respiration using a light response curve approach: critical issues and global evaluation. *Global Change Biol.* 16, 187–208.
- Law, B. E., Falge, E., Gu, L., Baldocchi, D. D., Bakwin, P. and co-authors. 2002. Environmental controls over carbon dioxide and water vapor exchange of terrestrial vegetation. *Agric. Forest Meteorol.* 113, 97–120.
- Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., Boden, T. A. and co-authors. 2014. Global carbon budget 2013. *Earth* Syst. Sci. Data. 6, 235–263.
- Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. F. and Nepstad, D. 2011. The 2010 Amazon drought. *Science*. 331, 554.
- Lobell, D. B., Bala, G. and Duffy, P. B. 2006. Biogeophysical impacts of cropland management changes on climate. *Geophys. Res. Lett.* 33, L06708. DOI: http://dx.doi.org/10.1029/ 2005GL025492
- Luo, Y., Keenan, T. F. and Smith, M. 2015. Predictability of the terrestrial carbon cycle. *Global Change Biol.* 21, 1737–1751.
- Ma, S., Baldocchi, D. D., Xu, L. and Hehn, T. 2007. Inter-annual variability in carbon dioxide exchange of an oak/grass Savanna and open grassland in California. *Agric. Forest Meteorol.* 147, 157–171.
- Madani, N., Kimball, J. S., Affleck, D. L. R., Kattge, J., Graham, J. and co-authors. 2014. Improving ecosystem productivity modeling through spatially explicit estimation of optimal light use efficiency: ecosystem optimal light use efficiency. J. Geophys. Res. 119, 1755–1769.
- Medlyn, B. E., Barton, C. V. M., Broadmeadow, M. S. J., Ceulemans, R., De Angelis, P. and co-authors. 2001. Stomatal conductance of forest species after long-term exposure to elevated CO₂ concentration: a synthesis. *New Phytol.* 149, 247–264.
- Migliavacca, M., Reichstein, M., Richardson, A. D., Colombo, R., Sutton, M. A. and co-authors. 2011. Semiempirical modeling of

abiotic and biotic factors controlling ecosystem respiration across eddy covariance sites. *Global Change Biol.* 17, 390-409.

- Milyukova, I. M., Kolle, O., Varlagin, A. V., Vygodskaya, N. N., Schulze, E.-D. and co-authors. 2002. Carbon balance of a southern taiga spruce stand in European Russia. *Tellus B.* **54**, 429–442.
- Moureaux, C., Debacq, A., Bodson, B., Heinesch, B. and Aubinet, M. 2006. Annual net ecosystem carbon exchange by a sugar beet crop. *Agric. Forest Meteorol.* 139, 25–39.
- Nagy, Z., Pintér, K., Czóbel, SZ., Balogh, J., Horváth, L. and co-authors. 2007. The carbon budget of semi-arid grassland in a wet and a dry year in Hungary. *Agric. Ecosyst. Environ.* **121**, 21– 29.
- Niu, S., Luo, Y., Fei, S., Yuan, W., Schimel, D. and co-authors. 2012. Thermal optimality of net ecosystem exchange of carbon dioxide and underlying mechanisms. *New Phytol.* 194, 775–783.
- Niu, S., Xing, X., Zhang, Z., Xia, J., Zhou, X. and co-authors. 2011. Water-use efficiency in response to climate change: from leaf to ecosystem in a temperate steppe. *Global Change Biol.* 17, 1073–1082.
- Noormets, A., Gavazzi, M. J., Mcnulty, S. G., Domec, J.-C., Sun, G. and co-authors. 2010. Response of carbon fluxes to drought in a coastal plain loblolly pine forest. *Global Change Biol.* **16**, 272–287.
- Papale, D., Reichstein, M. and Aubinet, M. 2006. Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation. *Biogeosciences.* 3, 571–583.
- Papale, D. and Valentini, R. 2003. A new assessment of European forests carbon exchanges by eddy fluxes and artificial neural network spatialization. *Global Change Biol.* 9, 525–535.
- Peichl, M., Sonnentag, O. and Nilsson, M. 2015. Bringing color into the picture: using digital repeat photography to investigate phenology controls of the carbon dioxide exchange in a boreal mire. *Ecosystems.* 18, 115–131.
- Pilegaard, K., Hummelshøj, P., Jensen, N. and Chen, Z. 2001. Two years of continuous CO₂ eddy-flux measurements over a Danish beech forest. *Agric. Forest Meteorol.* **107**, 29–41.
- Pintér, K., Barcza, Z., Balogh, J., Czóbel, S., Csintalan, Z. and coauthors. 2008. Interannual variability of grasslands' carbon balance depends on soil type. *Community Ecol.* 9, 43–48.
- Polgar, C. A. and Primack, R. B. 2011. Leaf-out phenology of temperate woody plants: from trees to ecosystems. *New Phytol.* 191, 926–941.
- Powell, T. L., Gholz, H. L., Clark, K. L., Starr, G., Cropper, W. P. and co-authors. 2008. Carbon exchange of a mature, naturally regenerated pine forest in north Florida. *Global Change Biol.* 14, 2523–2538.
- Prescher, A.-K., Grünwald, T. and Bernhofer, C. 2010. Land use regulates carbon budgets in eastern Germany: from NEE to NBP. Agric. Forest Meteorol. 150, 1016–1025.
- R Core Team. 2014. A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rebmann, C., Zeri, M., Lasslop, G., Mund, M., Kolle, O. and coauthors. 2010. Treatment and assessment of the CO₂-exchange

at a complex forest site in Thuringia, Germany. Agric. Forest Meteorol. 150, 684-691.

- Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M. and co-authors. 2005. On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm. *Global Change Biol.* **11**, 1424–1439.
- Rey, A., Pegoraro, E., Tedeschi, V., De Parri, I., Jarvis, G. and coauthors. 2002. Annual variation in soil respiration and its components in a coppice oak forest in Central Italy. *Global Change Biol.* 8, 851–866.
- Ricciuto, D. M., Butler, M. P., Davis, K. J., Cook, B. D., Bakwin, P. S. and co-authors. 2008. Causes of interannual variability in ecosystem–atmosphere CO₂ exchange in a northern Wisconsin forest using a Bayesian model calibration. *Agric. Forest Meteorol.* 148, 309–327.
- Richardson, A. D., Anderson, R. S., Arain, M. A., Barr, A. G., Bohrer, G. and co-authors. 2012. Terrestrial biosphere models need better representation of vegetation phenology: results from the North American Carbon Program Site Synthesis. *Global Change Biol.* 18, 566–584.
- Richardson, A. D. and Hollinger, D. Y. 2005. Statistical modeling of ecosystem respiration using eddy covariance data: maximum likelihood parameter estimation, and Monte Carlo simulation of model and parameter uncertainty, applied to three simple models. *Agric. Forest Meteorol.* 131, 191–208.
- Richardson, A. D., Hollinger, D. Y., Aber, J. D., Ollinger, S. V. and Braswell, B. H. 2007. Environmental variation is directly responsible for short- but not long-term variation in forest-atmosphere carbon exchange. *Global Change Biol.* 13, 788–803.
- Richardson, A. D., Hollinger, D. Y., Dail, D. B., Lee, J. T., Munger, J. W. and co-authors. 2009. Influence of spring phenology on seasonal and annual carbon balance in two contrasting New England forests. *Tree Physiol.* 29, 321–331.
- Rosseel, Y. 2012. Iavaan: an R package for structural equation modeling. J. Stat. Softw. 48, 1–36.
- Schimel, D. S., Braswell, B. H., Holland, E. A., McKeown, R., Ojima, D. S. and co-authors. 1994. Climatic, edaphic, and biotic controls over storage and turnover of carbon in soils. *Global Biogeochem. Cycles* 8, 279–293.
- Schulze, E.-D., Beck, E. and Müller-Hohenstein, K. 2005. Plant Ecology. Springer, Berlin, 23–44.
- Scott, R. L., Hamerlynck, E. P., Jenerette, G. D., Moran, M. S. and Barron-Gafford, G. A. 2010. Carbon dioxide exchange in a semidesert grassland through drought-induced vegetation change. J. Geophys. Res. 115, G03026. DOI: http://dx.doi.org/ 10.1029/2010JG001348
- Scott, R. L., Jenerette, G. D., Potts, D. L. and Huxman, T. E. 2009. Effects of seasonal drought on net carbon dioxide exchange from a woody-plant-encroached semiarid grassland. *J. Geophys. Res.* **114**, G04004. DOI: http://dx.doi.org/10.1029/ 2008JG000900
- Shao, J., Zhou, X., He, H., Yu, G., Wang, H. and co-authors. 2014. Partitioning climatic and biotic effects on interannual variability of ecosystem carbon exchange in three ecosystems. *Ecosystems*. 17, 1186–1201.

- Shao, J., Zhou, X., Luo, Y., Li, B., Aurela, M. and co-authors. 2015. Biotic and climatic controls on interannual variability in carbon fluxes across terrestrial ecosystems. *Agric. Forest Meteorol.* 205, 11–22.
- Shi, H., Li, L., Eamus, D., Cleverly, J., Huete, A. and co-authors. 2014. Intrinsic climate dependency of ecosystem light and wateruse-efficiencies across Australian biomes. *Environ. Res. Lett.* 9, 104002. DOI: http://dx.doi.org/10.1088/1748-9326/9/10/104002
- Shipley, B. 2004. A User's Guide to Path Analysis, Structural Equations and Causal Inference. Cambridge University Press, Cambridge, UK, pp. 100–135.
- Stoy, P. C., Katul, G. G., Siqueira, M. B. S., Juang, J.-Y., Novick, K. A. and co-authors. 2008. Role of vegetation in determining carbon sequestration along ecological succession in the southeastern United States. *Global Change Biol.* 14, 1409–1427.
- Sulman, B. N., Desai, A. R., Cook, B. D., Saliendra, N. and Mackay, D. S. 2009. Contrasting carbon dioxide fluxes between a drying shrub wetland in Northern Wisconsin, USA, and nearby forests. *Biogeosciences*. 6, 1115–1126.
- Suni, T., Berninger, F., Markkanen, T., Keronen, P., Rannik, Ü. and co-authors. 2003. Interannual variability and timing of growing-season CO₂ exchange in a boreal forest. J. Geophys. Res. 108, 4265. DOI: http://dx.doi.org/10.1029/2002JD002381
- Suyker, A. E., Verma, S. B., Burba, G. G., Arkebauer, T. J., Walters, D. T. and co-authors. 2004. Growing season carbon dioxide exchange in irrigated and rainfed maize. *Agric. Forest Meteorol.* 124, 1–13.
- Tan, Z.-H., Zhang, Y.-P., Deng, X.-B., Song, Q.-H., Liu, W.-J. and co-authors. 2015. Interannual and seasonal variability of water use efficiency in a tropical rainforest: results from a 9-year eddy flux time series. J. Geophys. Res. Atmos. 120, 464–479.
- Thomas, C. K., Law, B. E., Irvine, J., Martin, J. G., Pettijohn, J. C. and co-authors. 2009. Seasonal hydrology explains interannual and seasonal variation in carbon and water exchange in a semiarid mature *Ponderosa* pine forest in central Oregon. J. Geophys. Res. 114, G04006. DOI: http://dx.doi.org/ 10.1029/2009JG001010
- Thornley, J. H. M. 2011. Plant growth and respiration re-visited: maintenance respiration defined – it is an emergent property of, not a separate process within, the system – and why the respiration: photosynthesis ratio is conservative. *Ann. Bot.* **108**, 1365–1380.
- Tian, H., Melillo, J. M., Kicklighter, D. W., McGuire, A. D., Helfrich, J., III and co-authors. 2000. Climatic and biotic controls on annual carbon storage in Amazonian ecosystems. *Global Ecol. Biogeogr.* 9, 315–335.
- Tjoelker, M. G., Oleksyn, J. and Reich, P. B. 2001. Modelling respiration of vegetation: evidence for a general temperaturedependent Q₁₀. *Global Change Biol.* 7, 223–230.
- Urbanski, S., Barford, C., Wofsy, S., Kucharik, C., Pyle, E. and co-authors. 2007. Factors controlling CO₂ exchange on timescales from hourly to decadal at Harvard Forest. J. Geophys. Res. 112, G02020. DOI: http://dx.doi.org/10.1029/2006JG000293
- Vicente-Serrano, S. M., Gouveia, C., Camarero, J. J., Beguería, S., Trigo, R. and co-authors. 2013. Response of vegetation to drought time-scales across global land biomes. *Proc. Natl. Acad. Sci. USA*. 110, 52–57.

- Vickers, D., Thomas, C. K., Martin, J. G. and Law, B. 2009. Selfcorrelation between assimilation and respiration resulting from flux partitioning of eddy-covariance CO₂ fluxes. *Agric. Forest Meteorol.* 149, 1552–1555.
- Way, D. A. and Yamori, W. 2014. Thermal acclimation of photosynthesis: on the importance of adjusting our definitions and accounting for thermal acclimation of respiration. *Photosynth. Res.* **119**, 89–100.
- Welp, L. R., Randerson, J. T. and Liu, H. P. 2007. The sensitivity of carbon fluxes to spring warming and summer drought depends on plant functional type in boreal forest ecosystems. *Agric. Forest Meteorol.* 147, 172–185.
- Wohlfahrt, G., Hammerle, A., Haslwanter, A., Bahn, M., Tappeiner, U. and co-authors. 2008. Seasonal and inter-annual variability of the net ecosystem CO₂ exchange of a temperate mountain grassland: effects of weather and management. J. Geophys. Res. 113, D08110. DOI: http://dx.doi.org/10.1029/ 2007JD009286
- Wu, C., Chen, J. M., Black, T. A., Price, D. T., Kurz, W. A. and co-authors. 2013. Interannual variability of net ecosystem productivity in forests is explained by carbon flux phenology in autumn: autumn phenology and NEP. *Global Ecol. Biogeogr.* 22, 994–1006.
- Wythers, K. R., Reich, P. B. and Bradford, J. B. 2013. Incorporating temperature-sensitive Q_{10} and foliar respiration acclimation algorithms modifies modeled ecosystem responses to global change. J. Geophys. Res. **118**, 77–90.
- Xia, J., Niu, S., Ciais, P., Janssens, I. A., Chen, J. and co-authors. 2015. Joint control of terrestrial gross primary productivity by plant phenology and physiology. *Proc. Natl. Acad. Sci. USA*. 112, 2788–2793.
- Xiao, J., Sun, G., Chen, J., Chen, H., Chen, S. and co-authors. 2013. Carbon fluxes, evapotranspiration, and water use efficiency of terrestrial ecosystems in China. *Agric. Forest Meteorol.* 182–183, 76–90.
- Xiao, X., Zhang, Q., Hollinger, D., Aber, J. and Moore, B. 2005. Modeling gross primary production of an evergreen needleleaf forest using MODIS and climate data. *Ecol. Appl.* 15, 954–969.
- Xu, L. and Baldocchi, D. D. 2004. Seasonal variation in carbon dioxide exchange over a Mediterranean annual grassland in California. Agric. Forest Meteorol. 123, 79–96.
- Xu, L., Zhang, X., Shi, P., Li, W. and He, Y. 2007. Modeling the maximum apparent quantum use efficiency of alpine meadow ecosystem on Tibetan Plateau. *Ecol. Model.* 208, 129–134.
- Xue, B.-L., Guo, Q., Otto, A., Xiao, J., Tao, S. and co-authors. 2015. Global patterns, tends, and drivers of water use efficiency from 2000 to 2013. *Ecosphere*. 6, 174. DOI: http://dx.doi.org/10. 1890/ES14-00416.1

- Yamori, W., Hikosaka, K. and Way, D. A. 2014. Temperature response of photosynthesis in C3, C4, and CAM plants: temperature acclimation and temperature adaptation. *Photosynth. Res.* **119**, 101–117.
- Yan, W., Zhong, Y. and Shangguan, Z. 2015. The relationships and sensibility of wheat C:N:P stoichiometry and water use efficiency under nitrogen fertilization. *Plant Soil Environ*. 61, 201–207.
- Yang, B., Pallardy, S. G., Meyers, T. P., Gu, L.-H., Hanson, P. J. and co-authors. 2010. Environmental controls on water use efficiency during severe drought in an Ozark Forest in Missouri, USA. *Global Change Biol.* 16, 2252–2271.
- Yu, G.-R., Zhang, L.-M., Sun, X.-M., Fu, Y.-L., Wen, X.-F. and co-authors. 2008. Environmental controls over carbon exchange of three forest ecosystems in eastern China. *Global Change Biol.* 14, 2555–2571.
- Yuan, W., Cai, W., Xia, J., Chen, J., Liu, S. and co-authors. 2014. Global comparison of light use efficiency models for simulating terrestrial vegetation gross primary production based on the LaThuile database. *Agric. Forest Meteorol.* **192–193**, 108–120.
- Yuan, W., Luo, Y., Richardson, A. D., Oren, R., Luyssaert, S. and co-authors. 2009. Latitudinal patterns of magnitude and interannual variability in net ecosystem exchange regulated by biological and environmental variables. *Global Change Biol.* 15, 2905–2920.
- Zhang, L., Luo, Y., Yu, G. and Zhang, L. 2010. Estimated carbon residence times in three forest ecosystems of eastern China: applications of probabilistic inversion. J. Geophys. Res. 115, G01010. DOI: http://dx.doi.org/10.1029/2009JG001004
- Zhang, W., Wang, H., Yang, E., Yi, Y., Wen, X. and co-authors. 2011. Underestimated effects of low temperature during early growing season on carbon sequestration of a subtropical coniferous plantation. *Biogeosciences.* 8, 1667–1678.
- Zhou, L., Zhou, X., Zhang, B., Lu, M., Luo, Y. and co-authors. 2014. Different responses of soil respiration and its components to nitrogen addition among biomes: a meta-analysis. *Global Change Biol.* 20, 2332–2343.
- Zhou, Z., Guo, C. and Meng, H. 2013. Temperature sensitivity and basal rate of soil respiration and their determinants in temperate forests of North China. *PLoS One.* 8, e81793. DOI: http://dx.doi.org/10.1371/journal.pone.0081793
- Zhu, X.-J., Yu, G.-R., Wang, Q.-F., Hu, Z.-M., Zheng, H. and coauthors. 2015. Spatial variability of water use efficiency in China's terrestrial ecosystems. *Global Planet. Change*. **129**, 37– 44.