

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Biological Systems Engineering: Papers and Publications

Biological Systems Engineering

2019

Antibiotic resistance gene profile changes in cropland soil after manure application and rainfall

Morgan Meyers

USDA-ARS, morgan.meyers@usda.gov

Lisa Durso

United States Department of Agriculture, Agricultural Research Service, Lisa.Durso@ars.usda.gov

John E. Gilley

USDA-ARS, john.gilley@ars.usda.gov

Heidi Waldrip

USDA-ARS, heidi.waldrip@ars.usda.gov

Lana Castleberry

USDA-ARS, lana.castleberry@usda.gov

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/biosysengfacpub>



Part of the [Bioresource and Agricultural Engineering Commons](#), [Environmental Engineering Commons](#), and the [Other Civil and Environmental Engineering Commons](#)

Meyers, Morgan; Durso, Lisa; Gilley, John E.; Waldrip, Heidi; Castleberry, Lana; and Schmidt, Amy Millmier, "Antibiotic resistance gene profile changes in cropland soil after manure application and rainfall" (2019).

Biological Systems Engineering: Papers and Publications. 674.

<https://digitalcommons.unl.edu/biosysengfacpub/674>

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biological Systems Engineering: Papers and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Morgan Meyers, Lisa Durso, John E. Gilley, Heidi Waldrip, Lana Castleberry, and Amy Millmier Schmidt

Antibiotic resistance gene profile changes in cropland soil after manure application and rainfall

Morgan A. Meyers¹  | Lisa M. Durso¹  | John E. Gilley¹  | Heidi M. Waldrip²  | Lana Castleberry² | Amy Millmier-Schmidt³

¹USDA-ARS, Univ. of Nebraska-Lincoln East Campus, 251 Filley Hall, Lincoln, NE 68583, USA

²USDA-ARS Conservation and Production Research Lab., PO Drawer 10, Bushland, TX 79012, USA

³Dep. of Biological Systems Engineering, Univ. of Nebraska, 213 L. W. Chase Hall, PO Box 830726, Lincoln, NE 68583, USA

Correspondence

Lisa M. Durso, USDA-ARS, Univ. of Nebraska-Lincoln East Campus, 251 Filley Hall, Lincoln, NE 68583, USA.

Email: lisa.durso@ars.usda.gov

Assigned to Associate Editor Célia Manaia.

Abstract

Land application of manure introduces gastrointestinal microbes into the environment, including bacteria carrying antibiotic resistance genes (ARGs). Measuring soil ARGs is important for active stewardship efforts to minimize gene flow from agricultural production systems; however, the variety of sampling protocols and target genes makes it difficult to compare ARG results between studies. We used polymerase chain reaction (PCR) methods to characterize and/or quantify 27 ARG targets in soils from 20 replicate, long-term no-till plots, before and after swine manure application and simulated rainfall and runoff. All samples were negative for the 10 *b*-lactamase genes assayed. For tetracycline resistance, only source manure and post-application soil samples were positive. The mean number of macrolide, sulfonamide, and integrase genes increased in post-application soils when compared with source manure, but at plot level only, 1/20, 5/20, and 11/20 plots post-application showed an increase in *erm*(B), *sulI*, and *intI1*, respectively. Results confirmed the potential for temporary blooms of ARGs after manure application, likely linked to soil moisture levels. Results highlight uneven distribution of ARG targets, even within the same soil type and at the farm plot level. This heterogeneity presents a challenge for separating effects of manure application from background ARG noise under field conditions and needs to be considered when designing studies to evaluate the impact of best management practices to reduce ARG or for surveillance. We propose expressing normalized quantitative PCR (qPCR) ARG values as the number of ARG targets per 100,000 16S ribosomal RNA genes for ease of interpretation and to align with incidence rate data.

Abbreviations: ARG, antibiotic resistance gene; LOD, limit of detection; PCR, polymerase chain reaction; qPCR, quantitative polymerase chain reaction; rRNA, ribosomal RNA.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Published 2020. This article is a U.S. Government work and is in the public domain in the USA. *Journal of Environmental Quality* published by Wiley Periodicals, Inc. on behalf of American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America

1 | INTRODUCTION

The urgency and complexity of the global health crisis caused by antibiotic resistance is driving interest in understanding the prevalence, fate, and transport of agricultural and environmental antibiotic resistance. Soil is a natural

reservoir of antibiotic-resistant bacteria, and their genes can be found in soils across the globe (Cytryn, 2013; D'Costa, McGrann, Hughes, & Wright, 2006; Durso, Miller, & Wienhold, 2012). Human and animal feces are also a natural reservoir of antibiotic resistance genes (ARGs) (Stanton, Humphrey, & Stoffregen, 2011), and ARGs are enriched from individuals that receive antibiotics (Looft et al., 2012).

The application of animal manures to soil is an ancient practice (Bogaard, Heaton, Poulton, & Merbach, 2007), which continues today in both conventional and organic farming systems. Manure provides valuable nutrients for crops, increases soil organic matter and improves soil health (Doran & Zeiss, 2000; Edmeades, Thorrold, & Roberts, 2005; Garcia-Pausas, Rovira, Rabissi, & Romanyà, 2017). It also introduces gastrointestinal microbes into the environment, raising concerns related to infectious disease and antibiotic resistance (Marti et al., 2013; McKinney, Dungan, Moore, & Leytem, 2018).

The details of manure-borne ARG persistence in soils after land application remain unclear, with some studies reporting temporal declines reaching background levels over the course of a growing season (Chen et al., 2017; Durso, Miller, & Henry, 2018; Marti et al., 2013, 2014), whereas others report increases in specific ARG targets (Scott, Tien, Drury, Reynolds, & Topp, 2018). In one study, variable results were seen for *sulI* (sulfonamide), *erm(B)* (erythromycin), and *intI1* (Class 1 integron-integrase) genes immediately after land application of manures, where both exponential decay and increases were observed in the short-term during different years (Marti et al., 2014).

Deciphering the impact of manured soils on measures of antibiotic resistance in the environment is complicated by large areas to be tested, generally limited replication in field studies, and the known heterogeneity of soil systems (Kelley, 1922; Pepper & Brusseau, 2019). To explore the repeatability of field ARG measurements, we used a before-and-after framework common in land application studies to examine 20 replicate field plots receiving manure and simulated rainfall. We expected the ARG dynamics in all 20 plots to be similar, and pre-application soil ARG measurements to be lower than ARG measurements of recently manured soils.

2 | MATERIALS AND METHODS

2.1 | Sample collection

Soil samples (0–10 cm) were collected from plots at the University of Nebraska Rogers Memorial Farm (40°50'42" N, 96°28'19" W) (Gilley et al., 2017). Twenty no-till plots were established, perpendicular to the slope, on fields

Core Ideas

- There is uneven distribution of ARGs even in replicate plots of the same soil type and on the same farm.
- Temporary blooms of ARGs after manure application were likely linked to soil moisture levels.
- Heterogeneity of ARG distribution should be considered when planning environmental surveillance.

documented to have received no manure application since the farm was bequeathed to the university in 1947. The fields had previously been used for winter wheat (*Triticum aestivum* L.), soybean [*Glycine max* (L.) Merr.], corn (*Zea mays* L.), and grain sorghum [*Sorghum bicolor* (L.) Moench], with winter wheat harvested from the study site the previous year. The wheat residue was not chopped and provided 100% soil coverage. The Aksarben clay loam (fine, smectic, mesic Typic Arguidolls) was representative of soils found in southeastern Nebraska (Supplemental Table S1). Replicate plots were sampled at two time points, 1 wk apart (Supplemental Figure S1). Due to the labor-intensive nature of the field collection protocol involving simulated rainfall over the 20 plots (Gilley et al., 2017), two plots were processed each week during a 10-wk test period, with new manure applied weekly ($n = 10$ source manures).

Swine manure was collected weekly from a commercial deep pit swine operation in southeast Nebraska and transported in 20-L plastic buckets. An initial subsample of manure slurry was collected prior to the start of experiments and characterized for chemical and physical properties (Gilley et al., 2017). These values were used to calculate the application rate, with an estimated target of 151 kg N ha⁻¹ yr⁻¹, the rate needed to meet the annual N requirement for corn. Functionally, the addition of manure adds not only nutrients, but also antibiotic-resistant bacteria and ARGs to the soil. The slurry was surface applied by hand at a rate of 3.90×10^4 kg ha⁻¹, simulating a typical manure spreader. The manure was left on the soil surface without incorporation via tillage or other methods.

Twenty pre-application and 20 post-application samples were each collected from a 10-cm³ area that was 20 cm down-slope of the manure application zone, with samples from the two collection times collected directly adjacent to each other. Field collection utensils were wiped clean between uses and sprayed with ethanol. Soil and manure samples ($n = 10$ manure and 40 soil) were placed in coolers and transported immediately on ice to the laboratory for analyses.

2.2 | Microbiological analyses

Within 4 h of collection, the soil and manure samples were homogenized by hand mixing and stored in a -80°C freezer on arrival to the laboratory. The DNA was isolated from the 40 soil and 10 manure samples using Qiagen DNeasy Power Soil Kit (#12888-100) following manufacturer's instructions, except cell lysis was done with an OmniaBeadruptor12 (Omni International), at 2.40 m s^{-1} for 1 min. Standard polymerase chain reaction (PCR) was used to assay 14 tetracycline resistance genes, sulfonamide *sulI*, macrolide *erm(B)*, β -lactamase CTX-M-32, and integrase *intI1* (Supplemental Figure S1). All PCR and quantitative PCR (qPCR) details are presented in Supplemental Table S2. The PCR assay reaction mix consisted of $1 \times$ JumpStart REDTaq ReadyMix reaction mix (Sigma Chemical), $2.0\ \mu\text{M}$ primers, PCR-grade water, and DNA template. Thermocycling conditions are presented in Supplemental Table S3. Amplicons were visualized on a 2% agarose gel ($1 \times$ TAE) stained with SYBR Safe DNA gel stain (LifeTech) and imaged using the UVP GelDoc-It^{Ts3} Imager (Analytik Jena US). Pre-application soil results were confirmed with additional PCR testing, including controls for PCR inhibition. Detection of the 16S ribosomal RNA (rRNA) gene was used to confirm that PCR inhibitors had been removed during soil DNA extraction from all samples.

Individual qPCR assays were used to quantify sulfonamide, erythromycin, β -lactamase, integrase, and 16S rRNA genes. All samples were run in triplicate. Thermocycling conditions are listed in Supplemental Table S4. Quality conditions consisted of an efficiency between 90 and 110% and an R^2 value between .80 and 1.00. Melting curve analysis was performed for all assays to confirm authenticity of the PCR product. Limits of detection for the assays were determined by running, in triplicate, a series of dilutions using the gBlock (IDT) positive control, choosing the lowest dilution that amplified all samples with a standard deviation of Cq values (i.e., how many cycles were required to detect a measurable signal) that was <1.0 . The limits of detection (LODs) were four, three, four, and three copies per PCR reaction for *sulI*, *erm(B)*, CTX-M-32, and *intI1*, respectively. On a per-gram-soil basis, the LODs would be 1200–1600 copies g^{-1} (0.25 g soil extracted, 100 μl purified DNA extract, 1 μl amplified in each reaction). Samples were assayed for an additional nine clinically relevant β -lactamases using qPCR: CMY-2, CTX-M-15, CTX-M-14, OXA-48, IMP, VIM, DHA, KPC, and NDM, using the ARM-D Kit (Streck) for β -lactamase following manufacturer's instructions.

It is common to normalize ARG qPCR field measurements using the 16S rRNA gene, and these values are commonly displayed as the negative log of the normalized values. Here, we expressed these values as the number of ARG targets per




FIGURE 1 Percentage of 20 plots positive for individual tetracycline (TET) resistance genes. $N = 10$ manure. $N = 20$ post-application soils. Manure-borne antibiotic resistance genes (ARGs) account for many, but not all, of the ARGs enriched in soil after land application of swine slurry

100,000 16S rRNA genes, to align with incidence rate data commonly used to assess risk in public health settings.

3 | RESULTS

Antibiotic resistance gene detection varied, depending on the specific target measured, but all manure and soil samples were negative for the 10 β -lactamase genes surveyed and positive for 16S rRNA genes. Changes in soil antibiotic resistance gene profiles were observed for 17 out of 27 assayed genes after manure application and multiple rainfall events, with the remaining genes not detected in any samples.

3.1 | Qualitative or presence

Fourteen tetracycline resistance (TET^r) genes were assayed from the soil pre-application ($n = 20$), manure ($n = 10$), and soil post-application ($n = 20$) samples using standard PCR. None of the assayed TET^r genes were detected in pre-application soils, despite their presence in PCR positive controls. Out of the 14 TET^r genes assayed, 12 were detected in manure and post-application soils (Figure 1). All manure samples were positive for the same nine of the possible 14 TET^r genes. There was no change in the target ARG profile of the applied manure over the course of the experiment. The most frequently detected TET^r genes in the post-application soil were *tet(L)*, *tet(M)*, and *tet(O)*, which were found in 56% of post-application soil samples. Each of the 20 post-application soils had between zero and nine TET^r genes detected of the 14 that were assayed (Figure 1). Neither *tet(E)* nor *tetA(P)* were detected in any of the post-application soils. The mean




FIGURE 2 Carriage of *sull*, *erm(B)*, and *intI1* genes. Soil antibiotic resistance gene (ARG) profile changed after manure application and simulated rainfall

number of TET^r genes detected in manure and post-application soils were nine and six, respectively.

Standard PCR was also used to assay sulfonamide, macrolide, β -lactamase, and integrase genes (Figure 2). All samples were negative for β -lactamase (CTX-M-32) resistance genes. All manure samples were positive for sulfonamide resistance genes, macrolide resistance genes, and integrase. Of the post-application soil samples, 85% were positive for sulfonamide resistance genes, macrolide resistance genes, and integrase (17 of 20 post-application samples).

3.2 | qPCR (quantitative measures of antibiotic resistance)

The average number of copies of *sull*, *erm(B)*, CTX-M-32, *intI1*, and 16S rRNA in pre-application soil, manure, and post-application soil are presented in Table 1. Values were normalized based on the total number of 16S rRNA genes in the sample and are displayed in Figure 3. There were nine additional β -lactamase genes assayed using qPCR: CMY-2, CTX-M-15, CTX-M-14, OXA-48, IMP, VIM, DHA, KPC, and NDM. All soil and manure samples were negative for the nine additional β -lactamase genes.

4 | DISCUSSION

Methods that employ PCR are widely used in a research capacity for detecting antibiotic resistance in environmental samples (Marti et al., 2013; McKinney et al., 2018), offering the possibility to obtain information on target genes in hours as compared with culture-based methods that require days of laboratory work. However, PCR-based detection of ARGs from complex environmental samples also has limitations that make it difficult to compare results between studies or develop standardized environmental ARG monitoring efforts (Luby,




FIGURE 3 Quantification of antibiotic resistance gene (ARG) targets, normalized per 100,000 16S genes. Mean absolute values, although valuable for risk assessment purposes, obscure ecologically relevant information, such as variations in bacterial community, 16S copy number, and plot-based differences in ARG persistence. Lowest normalized copy number per 100,000 16S ribosomal RNA genes is zero

Ibekwe, Ziles, & Pruden, 2016). The first step to determine the potential for causal links between land application of animal manures and antibiotic resistance-derived adverse human or veterinary health outcomes (Williams-Nguyen et al., 2016) is to accurately measure antibiotic resistance in soil samples.

4.1 | Quantifying antibiotic resistance genes for individual plots versus mean

The qPCR method was used to quantify *sull*, *erm(B)*, CTX-M-32, *intI1*, and 16S rRNA genes (Gillings et al., 2015; Marti et al., 2014; Pruden, Pei, Sorteboom, & Carlson, 2006) from 20 replicate field plots, at two time points, and then normalized using 16S rRNA gene abundance. Looking at the normalized individual plot values contributing to the mean for these targets (Figure 3), it is evident that there was a large range of values across plots. For example, *sull* genes in post-application soil ranged from 0 to 12,400 copies. This highlights the heterogeneity associated with field measurements of ARGs. Mean values, although useful as a general summary tool, obscure the plot-level variability observed in this study. When collecting soils for field studies of manured soils, it is recommended to pool subsamples from multiple locations and homogenize thoroughly to minimize impact of soil heterogeneity on ARG conclusions.

The mean absolute values of the *sull*, *erm(B)*, and *intI1* ARGs per gram dry weight of soil in the post-application samples was higher than the combined mean values in the pre-application soil and applied manure (Table 1). This is likely due to the moist conditions after the rainfall events, in conjunction with an average Nebraskan July temperature of 32 °C—ideal conditions for manure- or soil-associated

TABLE 1 Quantitative polymerase chain reaction (qPCR) gene summary for sulfonamide *sulI*, erythromycin *erm(b)*, β -lactamase (*ctx-m-32*), and an integrase gene (*intI1*). An average moisture level of 90% was used to calculate the dry matter for manure (<https://extension2.missouri.edu/eq215>)

Gene	Pre-application soil	Manure	Post-application soil
	copies g ⁻¹ soil DM	copies g ⁻¹ manure DM	copies g ⁻¹ soil DM
<i>sulI</i>	2.12×10^3	1.91×10^6	4.37×10^7
<i>erm(B)</i>	ND	2.29×10^7	9.57×10^8
CTX-M-32	ND	ND	ND
<i>intI1</i>	3.17×10^3	3.29×10^5	4.12×10^7
16S ribosomal RNA	1.28×10^9	5.19×10^8	7.06×10^9

Note. DM, dry mass; ND, not detected.

microbes to proliferate and metabolize in the soil. Marti et al. (2014) hypothesized that the cooler and wetter spring weather was responsible for observed bacterial blooms, compared with warmer, drier conditions under which an immediate exponential decrease in bacteria was observed. Based on the current study and the results of Marti et al. (2014), it appears that temperature was less important than moisture in supporting proliferation of manure-borne bacteria after land application.

4.2 | Standard PCR screening for tetracycline resistance genes

Tetracycline resistance genes (TET^r) are frequently assayed in agricultural soils and are generally considered common (Cytryn, 2013; D'Costa et al., 2006; Durso et al., 2012). In the current study, none of the assayed TET^r genes were detected in pre-application soil samples. This result was unexpected, as previous studies have identified TET^r genes in nonmanured agricultural soils, organic farm soils, and prairie soils (Agga, Arthur, Durso, Harhay, & Schmidt, 2015; Cadena et al., 2018; Durso, Wedin, Gilley, Miller, & Marx, 2016). Unlike pre-application samples, multiple TET^r genes were detected in both manure and post-application soil samples (Figure 1); these data highlight the utility of TET^r genes as a marker for manure inputs at this site and provide evidence that applied manure is the primary source of the TET^r genes found in the post-application soils in this study.

Of note, however, is that three TET^r genes [*tet(A)*, *tet(D)*, and *tet(G)*] detected in post-application soils were absent in both pre-application soils and the source manure. We have two possible explanations:

1. Results could be due to soil heterogeneity. Post-application samples were collected directly adjacent to pre-application soils, but the destructive nature of field collection means the two samples were not identical.
2. It is possible that the genes were present in either the pre-application soils or source manure, but at a level below

our detection limit of three gene copies per PCR reaction (equivalent to 1.2×10^3 copies per gram of soil), and then the genes were enriched after the rainfall events.

Antibiotic resistance genes are widely considered to have originated in environmental bacteria (Perry, Waglechner, & Wright, 2016), and environmental bacteria remain an important reservoir for ARGs. (Gibson, Forsberg, & Dantas, 2015). Antibiotic resistance genes, including TET^r genes, are also common in feces, including feces of cattle, swine, humans, dogs, and fish (Brooks, Adeli, & McLaughlin, 2014; Durso et al., 2012). Tetracycline resistance genes evolved long before the discovery of antibiotics (Koike, Mackie, & Aminov, 2017; Rahman, Sakamoto, Kitamura, Nonaka, & Suzuki, 2015). However, the use of tetracycline, along with other selective pressures, has resulted in an increased proportion of bacteria harboring ARGs in clinical and environmental samples (Tan et al., 2018). This includes TET^r genes found in both organic and conventional farming systems (Brooks et al., 2014; Cadena et al., 2018; Marti et al., 2013, 2014). In this study, using plots that had never received any manure samples, the pre-application soils were negative for the TET^r genes assayed. In the current set of experiments, the swine manure harbored a distinct set of TET^r genes compared with the receiving agronomic soil.

For research and surveillance efforts, the specific ARG target measured has been shown to affect the conclusions of the study (Cadena et al., 2018; Durso et al., 2016; Liu, Jia, He, Zhang, & Ye, 2017; Walk et al., 2007). Even when associated with the same drug resistance category such as tetracycline, individual ARG targets, and sometimes even specific gene subtypes are not equivalent (Rahman et al., 2015). In the current study, we observed a smaller number of agronomic plots positive for all TET^r genes measured, compared with amounts originating from the source manure; however, the decrease varied considerably depending on the individual TET^r gene assayed (range 0–90%) (Figure 1). Contributing to this phenomenon is the fact that the types of genes detected in a sample depend, to a large extent, on the types of bacteria present (Durso et al., 2012; Durso, Miller, Schmidt, &

Callaway, 2017; Forsberg et al., 2014; Liu et al., 2017). This is particularly well documented for TET^r genes (Roberts & Schwarz, 2016; <https://faculty.washington.edu/marilynr/>).

The practical implications of individual TET^r genes being nonrandomly distributed in the environment is that target selection can influence the perceived efficacy of control measures, with different TET^r genes allowing for separate conclusions. For example, the short land application period for the current work resulted in manure-borne *tet(Q)* and *tet(S)* being nondetectable just days after manure application, but *tet(L)*, *tet(M)*, and *tet(O)* remained detectable in 90% of the plots. Thus, multiple TET^r gene targets need to be used to account for the differing persistence of individual ARGs. Furthermore, the background ARG profile of individual field sites must be considered (Durso et al., 2016).

Manure is an essential element of sustainable organic cropping systems, and responsible manure management is key to minimizing numerous adverse environmental and human health impacts. The addition of ARGs to soil, via manure application, results in an immediate increase in the number and types of soil ARG, due to a simple additive effect (Durso & Cook, 2014). There is also evidence that manure application enriches soil-borne bacteria, including indigenous bacteria that carry ARGs (Udikovic-Kolic, Wichmann, Broderick, & Handelsman, 2014), and that manure-borne bacteria can persist (Scott et al., 2018) and replicate in the environment (Walk, Alm, Calhoun, Mladonicky, & Whittam, 2007). Another important consideration in the study of ARG presence in manure-amended soils is high variability. Although the results from the current study demonstrated averaged potential increases in ARGs after land application of swine manure and rainfall, there was variability observed among the 20 replicate plots. These observations are supported by the work of Marti et al. (2014) and Udikovic-Kolic et al. (2014). However, it would be presumptuous to draw any broad conclusions on heterogeneity of ARG presence within plots based on these limited data. There were two data points that were significant outliers (Figure 3), potentially due to uneven distribution of the manure. Variability was also observed in the quantity of specific gene targets, supporting the idea that individual ARG targets are not equivalent for measuring general antibiotic resistance (Durso et al., 2016).

Data from this study highlight the challenge of differentiating the true effect of manure application from ARG background noise under field conditions when measuring frequency of ARG occurrence and abundance, even when there is a high degree of replication within the study. Our results support recommendations to collect background or baseline data, particularly when examining impacts of human or animal activities on ARG, so that results can be interpreted in light of the amounts and kinds of targets already present before the treatment or application began (Dungan,

McKinney, & Leytem, 2018; Durso & Cook, 2014; Rothrock et al., 2016).

ACKNOWLEDGMENTS

We wish to thank Katelyn Tucker and USDA-ARS technical staff for field and laboratory assistance, and Dr. Daniel N. Miller for helpful comments on the manuscript. This work was funded by the USDA-ARS National Program 212 Soil and Air. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture. USDA is an equal opportunity provider and employer.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

Morgan A. Meyers 

<https://orcid.org/0000-0003-0965-7466>

Lisa M. Durso  <https://orcid.org/0000-0002-9702-6463>

John E. Gilley  <https://orcid.org/0000-0002-8167-5362>

Heidi M. Waldrip  <https://orcid.org/0000-0001-7714-5417>

REFERENCES

- Agga, G. E., Arthur, T. M., Durso, L. M., Harhay, D. M., & Schmidt, J. W. (2015). Antimicrobial-resistant bacterial populations and antimicrobial resistance genes obtained from environments impacted by livestock and municipal waste. *PLOS One*, *10*, e0132586. <https://doi.org/10.1371/journal.pone.0132586>
- Bogaard, A., Heaton, T. H. E., Poulton, P., & Merbach, I. (2007). The impact of manuring on nitrogen isotope ratios in cereals: Archaeological implications for reconstruction of diet and crop management practices. *Journal of Archaeological Science*, *34*, 335–343. <https://doi.org/10.1016/j.jas.2006.04.009>
- Brooks, J. P., Adeli, A., & McLaughlin, M. R. (2014). Microbial ecology, bacterial pathogens, and antibiotic resistant genes in swine manure wastewater as influenced by three swine management systems. *Water Research*, *57*, 96–103. <https://doi.org/10.1016/j.watres.2014.03.017>
- Cadena, M., Durso, L. M., Miller, D. N., Waldrip, H. M., Castleberry, B. L., Drijber, R. A., & Wortmann, C. (2018). Tetracycline and sulfonamide antibiotic resistance genes in soils from Nebraska organic farming operations. *Frontiers in Microbiology*, *9*, 1283. <https://doi.org/10.3389/fmicb.2018.01283>
- Chen, Q. L., An, X. L., Li, H., Zhu, Y. G., Su, J. Q., & Cui, L. (2017). Do manure-borne or indigenous soil microorganisms influence the spread of antibiotic resistance genes in manured soil? *Soil Biology and Biochemistry*, *114*, 229–237. <https://doi.org/10.1016/j.soilbio.2017.07.022>
- Cytryn, E. (2013). The soil resistome: The anthropogenic, the native, and the unknown. *Soil Biology and Biochemistry*, *63*, 18–23. <https://doi.org/10.1016/j.soilbio.2013.03.017>
- D'Costa, V. M., McGrann, K. M., Hughes, D. W., & Wright, G. D. (2006). Sampling the antibiotic resistome. *Science*, *311*, 374–377. <https://doi.org/10.1126/science.1120800>

- Doran, J. W., & Zeiss, M. R. (2000). Soil health and sustainability: Managing the biotic component of soil quality. *Applied Soil Ecology*, *15*, 3–11. [https://doi.org/10.1016/S0929-1393\(00\)00067-6](https://doi.org/10.1016/S0929-1393(00)00067-6)
- Dungan, R. S., McKinney, C. W., & Leytem, A. B. (2018). Tracking antibiotic resistance genes in soil irrigated with dairy wastewater. *Science of the Total Environment*, *635*, 1477–1483. <https://doi.org/10.1016/j.scitotenv.2018.04.020>
- Durso, L. M., & Cook, K. L. (2014). Impacts of antibiotic use in agriculture: What are the benefits and risks? *Current Opinion in Microbiology*, *19*, 37–44. <https://doi.org/10.1016/j.mib.2014.05.019>
- Durso, L. M., Miller, D. N., & Henry, C. G. (2018). Impact of vegetative treatment systems on multiple measures of antibiotic resistance in agricultural wastewater. 2018. *International Journal of Environmental Research and Public Health*, *15*, 1295. <https://doi.org/10.3390/ijerph15071295>
- Durso, L. M., Miller, D. N., Schmidt, T. B., & Callaway, T. (2017). Tracking bacteria through the entire gastrointestinal tract of a beef steer. *Agricultural & Environmental Letters*, *2*, 170016. <https://doi.org/10.2134/ael2017.05.0016>
- Durso, L. M., Miller, D. N., & Wienhold, B. J. (2012). Distribution and quantification of antibiotic resistant genes and bacteria across agricultural and nonagricultural metagenomes. *PLOS One*, *7*, e48325. <https://doi.org/10.1371/journal.pone.0048325>
- Durso, L. M., Wedin, D., Gilley, J. E., Miller, D. N., & Marx, D. B. (2016). Assessment of selected antibiotic resistances in ungrazed native Nebraska prairie soils. *Journal of Environmental Quality*, *45*, 454–462. <https://doi.org/10.2134/jeq2015.06.0280>
- Edmeades, D. C., Thorrold, B. S., & Roberts, A. H. C. (2005). The diagnosis and correction of sulfur deficiency and the management of sulfur requirements in New Zealand pastures: A review. *Australian Journal of Experimental Agriculture*, *45*, 1205–1223. <https://doi.org/10.1071/EA01173>
- Forsberg, K. J., Patel, S., Gibson, M. K., Lauber, C. L., Knight, R., Fierer, N., & Dantas, G. (2014). Bacterial phylogeny structures soil resistomes across habitats. *Nature*, *509*, 612–616. <https://doi.org/10.1038/nature13377>
- Garcia-Pausas, J., Rovira, P., Rabissi, A., & Romanyà, J. (2017). Organic fertilisation increases C and N stocks and reduces soil organic matter stability in Mediterranean vegetable gardens. *Land Degradation & Development*, *28*, 691–698. <https://doi.org/10.1002/ldr.2569>
- Gibson, M. K., Forsberg, K. J., & Dantas, G. (2015). Improved annotation of antibiotic resistance determinants reveals microbial resistomes cluster by ecology. *ISME Journal*, *9*, 207–216. <https://doi.org/10.1038/ismej.2014.106>
- Gilley, J. E., Bartelt-Hunt, S. L., Eskridge, K., Li, X., Schmidt, A. M., & Snow, D. D. (2017). Setback distance requirements for removal of swine slurry constituents in runoff. *Transactions of the ASABE*, *60*, 1885–1894. <https://doi.org/10.13031/trans.12310>
- Gillings, M. R., Gaze, W. H., Pruden, A., Smalla, K., Tiedje, J. M., & Zhu, Y. G. (2015). Using the Class I integron-integrase gene as a proxy for anthropogenic pollution. *ISME Journal*, *9*, 1269–1279. <https://doi.org/10.1038/ismej.2014.226>
- Kelley, W. P. (1922). Variability of alkali soil. *Soil Science*, *14*, 177–190. <https://doi.org/10.1097/00010694-192209000-00002>
- Koike, S., Mackie, R., & Aminov, R. (2017). Agricultural use of antibiotics and antibiotic resistance. In S. Mirete & M. Lopez Perez (Eds.), *Antibiotic resistance genes in natural environments and long-term effects* (Chapter 8). Hauppauge, NY: Nova Science.
- Liu, P., Jia, S., He, X., Zhang, X., & Ye, L. (2017). Different impacts of manure and chemical fertilizers on bacterial community structure and antibiotic resistance genes in arable soils. *Chemosphere*, *188*, 455–464. <https://doi.org/10.1016/j.chemosphere.2017.08.162>
- Looft, T., Johnson, T. A., Allen, H. K., Bayles, D. O., Alt, D. P., Stedtfield, R. D., ... Stanton, T. B. (2012). In-feed antibiotic effects on the swine intestinal microbiome. *Proceedings of the National Academy of Sciences of the USA*, *109*, 1691–1696. <https://doi.org/10.1073/pnas.1120238109>
- Luby, E., Ibekwe, A. M., Ziles, J., & Pruden, A. (2016). Molecular methods for assessment of antibiotic resistance in agricultural ecosystems: Prospects and challenges. *Journal of Environmental Quality*, *45*, 441–453. <https://doi.org/10.2134/jeq2015.07.0367>
- Marti, R., Scott, A., Tien, Y. C., Murray, R., Sabourin, L., Zhang, Y., & Topp, E. (2013). Impact of manure fertilization on the abundance of antibiotic-resistant bacteria and frequency of detection of antibiotic resistance genes in soil and on vegetables at harvest. *Journal of Environmental Quality*, *79*, 5701–5709. <https://doi.org/10.1128/AEM.01682-13>
- Marti, R., Tien, Y. C., Murray, R., Scott, A., Sabourin, L., & Topp, E. (2014). Safely coupling livestock and crop production systems: How rapidly do antibiotic resistance genes dissipate in soil following a commercial application of swine or dairy manure? *Applied and Environmental Microbiology*, *80*, 3258–3265. <https://doi.org/10.1128/AEM.00231-14>
- McKinney, C. W., Dungan, R. S., Moore, A., & Leytem, A. B. (2018). Occurrence and abundance of antibiotic resistance genes in agricultural soil receiving dairy manure. *FEMS Microbiology Ecology*, *94*, fiy010. <https://doi.org/10.1093/femsec/fiy010>
- Pepper, I. L., and Brusseau, M. L. (2019). Physical-chemical characteristics of soils and the subsurface. In M. L. Brusseau, I. Pepper, & C. Gerba (Eds.), *Environmental and pollution science* (3rd ed., pp. 9–22). London: Academic Press. <https://doi.org/10.1016/B978-0-12-814719-1.00002-1>
- Perry, J., Waglechner, N., & Wright, G. (2016). The prehistory of antibiotic resistance. *Cold Spring Harbor Perspectives in Medicine*, *6*, a025197. <https://doi.org/10.1101/cshperspect.a025197>
- Pruden, A., Pei, R., Sorteboom, H., & Carlson, K. H. (2006). Antibiotic resistance genes as emerging contaminants: Studies in northern Colorado. *Environmental Science & Technology*, *40*, 7445–7450. <https://doi.org/10.1021/es0604131>
- Rahman, M. H., Sakamoto, K. Q., Kitamura, S. I., Nonaka, L., & Suzuki, S. (2015). Diversity of tetracycline-resistant bacteria and resistance gene *tet(M)* in fecal microbial community of Adélie penguin in Antarctica. *Polar Biology*, *38*, 1775–1781. <https://doi.org/10.1007/s00300-015-1732-x>
- Roberts, M. C., & Schwarz, S. (2016). Tetracycline and phenicol resistance genes and mechanisms: Importance for agriculture, the environment, and humans. *Journal of Environmental Quality*, *45*, 576–592. <https://doi.org/10.2134/jeq2015.04.0207>
- Rothrock, M. J., Keen, P. L., Cook, K. L., Durso, L. M., Franklin, A. M., & Dungan, R. S. (2016). How should we be determining background and baseline antibiotic resistance levels in agroecosystem research? *Journal of Environmental Quality*, *45*, 420–431. <https://doi.org/10.2134/jeq2015.06.0327>
- Scott, A., Tien, U. Y. C., Drury, C. F., Reynolds, W. D., & Topp, E. (2018). Enrichment of antibiotic resistance genes in soil receiving composts derived from swine manure, yard wastes, or food wastes, and evidence for multiyear persistence of swine

- Clostridium* spp. *Canadian Journal of Microbiology*, 64, 201–208. <https://doi.org/10.1139/cjm-2017-0642>
- Stanton, T. B., Humphrey, S. B., & Stoffregen, W. C. (2011). Chlortetracycline-resistant bacteria in organically raised and feral swine. *Applied and Environmental Microbiology*, 77, 7167–7170. <https://doi.org/10.1128/AEM.00688-11>
- Tan, L., Li, L., Ashbolt, N., Wang, X., Cui, Y., Zhu, X., ... Luo, Y. (2018). Arctic antibiotic resistance gene contamination, a result of anthropogenic activities and natural origin. *Science of the Total Environment*, 621, 1176–1184. <https://doi.org/10.1016/j.scitotenv.2017.10.110>
- Udikovic-Kolic, N., Wichmann, F., Broderick, N. A., & Handelsman, J. (2014). Bloom of resident antibiotic-resistant bacteria in soil following manure fertilization. *Proceedings of the National Academy of Sciences of the USA*, 111, 15202–15207. <https://doi.org/10.1073/pnas.1409836111>
- Walk, S. T., Alm, E. W., Calhoun, L. M., Mladonicky, J. M., & Whittam, T. S. (2007). Genetic diversity and population structure of *Escherichia coli* isolated from freshwater beaches. *Environmental Microbiology*, 9, 2274–2288. <https://doi.org/10.1111/j.1462-2920.2007.01341.x>
- Walk, S. T., Mladonicky, J. M., Middleton, J. A., Heidt, A. J., Cunningham, J. R., Bartlett, P., ... Whittam, T. S. (2007). Influence of antibiotic selection on genetic composition of *Escherichia coli* populations from conventional and organic dairy farms. *Applied and Environmental Microbiology*, 73, 5982–5989. <https://doi.org/10.1128/AEM.00709-07>
- Williams-Nguyen, J., Sallach, J. B., Bartelt Hunt, S., Boxall, A. B., Durso, L. M., McLain, J. E., ... Ziles, J. L. (2016). Antibiotics and antibiotic resistance in agroecosystems: State of the science. *Journal of Environmental Quality*, 45, 394–406. <https://doi.org/10.2134/jeq2015.07.0336>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Meyers MA, Durso LM, Gilley JE, Waldrip HM, Castleberry L, Millmier-Schmidt A. Antibiotic resistance gene profile changes in cropland soil after manure application and rainfall. *J. Environ. Qual.* 2020;49:754–761. <https://doi.org/10.1002/jeq2.20060>