

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Insecta Mundi

Center for Systematic Entomology, Gainesville,  
Florida

---

4-15-2011

## New collections and records for earwigflies and scorpionflies (Mecoptera: Meropeidae and Panorpididae) in Florida

Louis A. Somma

*Florida State Collection of Arthropods*, [somma@ufl.edu](mailto:somma@ufl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

---

Somma, Louis A., "New collections and records for earwigflies and scorpionflies (Mecoptera: Meropeidae and Panorpididae) in Florida" (2011). *Insecta Mundi*. 690.

<https://digitalcommons.unl.edu/insectamundi/690>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# INSECTA MUNDI

---

A Journal of World Insect Systematics

**0165**

New collections and records for earwigflies and scorpionflies  
(Mecoptera: Meropeidae and Panorpididae) in Florida

Louis A. Somma  
Florida State Collection of Arthropods  
Division of Plant Industry  
Florida Department of Agriculture & Consumer Services  
PO Box 147100  
Gainesville, FL 32614-7100

Date of Issue: April 15, 2011

Louis A. Somma

New collections and records for earwigflies and scorpionflies (Mecoptera: Meropidae and Panorpididae) in Florida  
*Insecta Mundi* 0165: 1-4

**Published in 2011 by**

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 U. S. A.

<http://www.centerforsystematicentomology.org/>

**Insecta Mundi** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

**Insecta Mundi** is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

**Managing editor:** Paul E. Skelley, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Production editor:** Michael C. Thomas & Ian Stocks, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Editorial board:** J. H. Frank, M. J. Paulsen

**Subject editors:** G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

**Printed copies deposited in libraries of:**

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, Great Britain

Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

**Electronic copies in PDF format:**

Printed CD mailed to all members at end of year.

Florida Center for Library Automation: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

**Author instructions** available on the *Insecta Mundi* page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

**Printed copies deposited in libraries** (ISSN 0749-6737)

**Electronic copies in PDF format** (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362)

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

## New collections and records for earwigflies and scorpionflies (Mecoptera: Meropeidae and Panorpidae) in Florida

Louis A. Somma

Florida State Collection of Arthropods

Division of Plant Industry

Florida Department of Agriculture & Consumer Services

PO Box 147100

Gainesville, FL 32614-7100

Louis.Somma@freshfromflorida.com

**Abstract.** I add new collection and phenological data on the North American earwigfly, *Merope tuber* Newman, and new county records for the red scorpionfly, *Panorpa rufa* Gray, and veined scorpionfly, *Panorpa venosa* Westwood, in Florida. Additionally, I report on a new Georgia county record for the extralimital species, *Panorpa ferruginea* Byers, the ferruginous scorpionfly, and speculate on its potential occurrence in Florida.

### Introduction

Mecoptera are a small assemblage of ancestral, largely understudied, and undoubtedly paraphyletic taxa of insects represented by fewer than 700 extant species arranged in as many as 33 genera and 9 families worldwide, and the subject of vigorous systematic debate (Penny 1975; Willmann 1987; [Novokshonov] 1997; Kluge 2004; Byers 2005; Grimaldi and Engel 2005; Beutel and Baum 2008; Beutel et al. 2008, 2009; Dunford and Somma 2008a, b; Mickoleit 2008; Machado et al. 2009; Ren et al. 2009; Bicha 2010; Friedrich and Beutel 2010; Minet et al. 2010). At least 12 species of mecopterans have been documented from Florida (Dunford et al. 2007; Somma and Dunford 2008, 2009a, b; Somma 2010), representing the families Meropeidae (one species), Bittacidae (4 species) and Panorpidae (7 species). Herein, I provide data on new collections and county records for Floridian Mecoptera. These new data are represented by the families Meropeidae, *Merope tuber* Newman 1838; and Panorpidae, *Panorpa rufa* Gray 1832 and *P. venosa* Westwood 1846. Additionally, I report on a southern Georgia county record for *Panorpa ferruginea* Byers 1993, near the Florida/Georgia border, and speculate on its possible occurrence in northern Florida. My ultimate goal is to stimulate greater interest in studying Mecoptera in Florida.

### Methods and Materials

I examined two *M. tuber* and identified an undetermined panorpid collected from the Apalachicola Bluffs and Ravines Preserve (ABRP), Liberty County, Florida, and deposited in the Bug Closet, University of Central Florida Collection of Arthropods (UCFC). All UCFC specimens have individual accession numbers. Moreover, Wesley J. Bicha examined a specimen of panorpid in the Florida State Collection of Arthropods (FSCA) listed as undetermined (Alachua County, Florida) by Somma and Dunford (2008), and another undetermined panorpid in FSCA collected from Georgia, and determined the species identifications for each.

### Results and Discussion

***Merope tuber* Newman**, the North American earwigfly, is one of two extant species of the family Meropeidae (Byers 1973, 2005; Abbott et al. 2007; Dunford et al. 2008a, b). The distribution of this seemingly secretive meropeid is in eastern North America, from southeastern Canada to northern Georgia and southern Alabama, west to eastern Kansas, central Iowa and Minnesota, with more recently discovered populations in Leon and Liberty counties in the Florida Panhandle (Byers 1973, 1993, 2002; Schiefer and Dunford 2005; Cheung et al. 2006; Dunford et al. 2006, 2007, 2008a, b; Somma and Dunford 2007; N. Penny pers. comm. 2010).

On 14 September 2008 a male *M. tuber* (UCFC 0402 713) was collected by Preserve staff with a Malaise trap set up on a seepage slope in a steephead ravine at ABRP, Liberty County, Florida (30.495183°N,

84.978461°W). On 28 December 2008 a female *M. tuber* (UCFC 0405 732) was collected from the same trap and locality by Preserve staff. These two specimens represent the first individuals collected from Florida in approximately 12 years (Dunford et al. 2007). Moreover, the female *M. tuber* represents the latest recorded season for which this species has ever been collected (Dunford et al. 2007).

***Panorpa rufa* Gray**, the red scorpionfly, is an elusive, xerophilous, autumnal-winter panorpid sparsely distributed throughout much of the southeastern United States, in sandy ecosystems east of the Mississippi River, and previously known from eight counties in northern Florida as far south as Alachua and Bay counties (Hagen 1861; Hine 1901a, b; Esben-Petersen 1921; Carpenter 1931; Byers 1954, 1993; Somma and Dunford 2008, 2009b).

A female *P. rufa* (UCFC 0414 950) was collected by Preserve staff with a Malaise trap on 16 November 2008 from ABRP, Liberty County, Florida (30.494725°N, 84.97955°W), in restored sandhills habitat dominated by longleaf pine, turkey oak and wiregrass. This specimen represents the first record for *P. rufa* in Liberty County; adding a ninth county from which this species has been collected in Florida.

***Panorpa venosa* Westwood**, the veined scorpionfly, is principally a coastal and eastern Georgian species that has recently been discovered in Baker and Leon counties, northern Florida (Byers 1993; Somma and Dunford 2008; W. Bicha pers. comm. 2009). This suggests an overall Coastal Plains distribution. Habitat for *P. venosa* collected by W. Bicha (pers. comm. 2010) in Baker County, Florida, was the upper leaf surfaces of a Japanese Honeysuckle thicket inside the edge of mixed hardwood forest on the outskirts of Glen St. Mary.

An adult female *P. venosa* in FSCA was collected with a Malaise trap from the J. & E. Turf Farm, north of LaCrosse, in the Santa Fe River floodplain east of State Route 121, Alachua County, Florida (29.8942°N, 82.4044°W), on 19-26 July 1996 by G. J. Steck and B. D. Sutton. This specimen was erroneously listed as collected in 1986 and described as an undetermined member of the *P. virginica* species group by Somma and Dunford (2008). The trap was periodically partially immersed in rising river water while in use (G. Steck pers. comm. 2010). This specimen is a county record, approximately 42.5 km southwest of the nearest recorded locality for specimens collected from Baker County in late November (Somma and Dunford 2008; W. Bicha pers. comm. 2010), and represents the southernmost record for *P. venosa* in continental North America.

***Panorpa ferruginea* Byers**, the ferruginous scorpionfly, is indigenous to west-central Georgia, extreme northeastern Mississippi, and much of eastern Alabama as far south as Wing, Covington County, approximately 3.2 km from the border of Okaloosa County in the Florida panhandle (Byers 1993). The preferred habitat of this panorpid is dense undergrowth in pine woods mixed with a scattering of hardwoods, and a ground cover of pine needles (Byers 1993).

A male *P. ferruginea* housed in FSCA was collected by E. I. Hazard on 12 November 1962 from an unspecified locality in Lowndes County, Georgia. This is a first county record for this species and approximately 243 km south-southeast from the nearest record in Georgia (Byers 1993). Lowndes County, Georgia, shares its southern border with Hamilton County in northern peninsular Florida. The records for *P. ferruginea* in southernmost Alabama and southernmost Georgia suggest that this species eventually could be found in northern Florida if searched for in suitable habitat.

Somma and Dunford (2008, 2009b) have previously emphasized the dearth of information on and collecting of mecopterans in Florida. Important new data on distribution and phenology of Mecoptera was derived from a mere five specimens reported herein, and clearly underscores the need for further intensive collecting. Undoubtedly *P. ferruginea* will be added to Florida's indigenous fauna in the future. The state of Florida could be the "undiscovered country" for North American Mecoptera.

## Acknowledgments

As usual I am greatly indebted to Wes Bicha (Oliver Springs, Tennessee) for carefully examining and identifying the FSCA panorpid in this study, reviewing the manuscript for this publication, and providing important insights based up his remarkable expertise. Stuart M. Fullerton and Shawn Kelly (Department of Biology, University of Central Florida) graciously allowed me to examine UCFC Bug Closet specimens collected by the diligent Preserve staff of ABRP, and provided additional data. Gary J. Steck (Curator of Diptera and Minor Orders, FSCA) and Bruce D. Sutton (Curator of Diptera, FSCA) unstintingly

provided invaluable data for the *P. venosa* specimen in this study. I am gratified that Norman D. Penny (California Academy of Sciences) took the time to critically review the manuscript for this publication. I thank Gerhard Mickoleit (Zoologische Schausammlung der Universität, Tübingen) and Beverly Pope (DPI Librarian) for providing important literature cited in this study. This is Entomology Contribution No. 1185, Bureau of Entomology, Nematology and Plant Pathology, DPI, Florida Department of Agriculture & Consumer Services, which also houses the *P. venosa* and *P. ferruginea* specimens examined, and provided technical and library support for this research.

### Literature Cited

- Abbott, I., T. Burbidge, and A. Wills. 2007.** *Austromerope poultoni* (Insecta, Mecoptera) in south-west Western Australia: Occurrence, modelled geographic distribution, and phenology. *Journal of the Royal Society of Western Australia* 90: 97-106.
- Beutel, R. G., and E. Baum. 2008.** A longstanding entomological problem finally solved? Head morphology of *Nannochorista* (Mecoptera, Insecta) and possible phylogenetic implications. *Journal of Zoological Systematics and Evolutionary Research* 46: 346-367.
- Beutel, R. G., F. Friedrich, and M. F. Whiting. 2008.** Head morphology of *Caurinus* (Mecoptera: Boreidae) and its phylogenetic implications. *Arthropod Structure & Development* 37: 418-433.
- Beutel, R. G., N. P. Kristensen, and H. Pohl. 2009.** Resolving insect phylogeny: The significance of cephalic structures of the Nannomecoptera in understanding endopterygote relationships. *Arthropod Structure & Development* 38: 427-460.
- Bicha, W. J. 2010.** A review of scorpionflies (Mecoptera) of Indochina with the description of a new species of *Neopanorpa* from northern Thailand. *Zootaxa* 2480: 61-67.
- Byers, G. W. 1954.** Notes on North American Mecoptera. *Annals of the Entomological Society of America* 47: 484-510.
- Byers, G. W. 1973.** Zoogeography of the Meropeidae (Mecoptera). *Journal of the Kansas Entomological Society* 46: 511-516.
- Byers, G. W. 1993.** Autumnal Mecoptera of southeastern United States. *University of Kansas Science Bulletin* 55: 57-96.
- Byers, G. W. 2002.** Scorpionflies, hangingflies, and other Mecoptera. *Kansas School Naturalist* 48: 3-15.
- Byers, G. W. 2005 [2004].** Order Mecoptera. Scorpionflies and hangingflies. p. 662-668. *In*: C. A. Triplehorn and N. F. Johnson (editors). *Borror and DeLong's introduction to the study of insects*. Seventh edition. Thomson Brooks/Cole; Belmont, CA. 864 p.
- Carpenter, F. M. 1931.** Revision of the Nearctic Mecoptera. *Bulletin of the Museum of Comparative Zoölogy at Harvard College* 72: 205-277, pl. 1-8.
- Cheung, D. K. B., S. A. Marshall, and D. W. Webb. 2006.** Mecoptera of Ontario. *Canadian Journal of Arthropod Identification* 1: 1-13.
- Dunford, J. C., P. W. Kovarik, L. A. Somma, and D. Serrano. 2007.** First state records for *Merope tuber* (Mecoptera: Meropeidae) in Florida and biogeographical implications. *Florida Entomologist* 90: 581-584.
- Dunford, J. C., D. Serrano, and L. A. Somma. 2006.** Earwigflies in the Great Smokies. *Southeastern Biology* 53: 27-29.
- Dunford, J. C., and L. A. Somma. 2008a.** Scorpionflies (Mecoptera). p. 3304-3310, pl. 97. *In*: J. L. Capinera (editor). *Encyclopedia of entomology*. Second ed. Vol. 4. S-Z. Springer; [Dordrecht].
- Dunford, J. C., and L. A. Somma. 2008b.** Scorpionflies (Mecoptera). Available on URL: <http://springerlink.com/content/134653515g813298/fulltext.html>. *In*: J. L. Capinera (editor). *Encyclopedia of entomology* (e-book). Springer Science + Business Media B. V.; [Dordrecht]. (Last accessed 28 October 2010.)
- Dunford, J. C., L. A. Somma, and D. Serrano. 2008a.** Earwigflies (Mecoptera: Meropeidae). p. 1261-1264. *In*: J. L. Capinera (editor). *Encyclopedia of entomology*. Second ed. Vol. 2. D-K. Springer; [Dordrecht].
- Dunford, J. C., L. A. Somma, and D. Serrano. 2008b.** Earwigflies (Mecoptera: Meropeidae). Available on URL: <http://springerlink.com/content/t26418954r355x4t/fulltext.html>. *In*: J. L. Capinera (editor).



- Encyclopedia of entomology (e-book). Springer Science + Business Media B. V.; [Dordrecht]. (Last accessed 28 October 2010.)
- Esben-Petersen, P. 1921.** Mecoptera. Monographic revision. Collections Zoologiques du Baron Edm. de Selys Longchamps; Catalogue Systématique et Descriptif, Publié par les Soins de ses Fils (Bruxelles) Fascicule 5(2): 1-172, pl. I-II.
- Friedrich, F., and R. G. Beutel. 2010.** The thoracic morphology of *Nannochorista* (Nannochoristidae) and its implications for the phylogeny of Mecoptera and Antliophora. *Journal of Zoological Systematics and Evolutionary Research* 48: 50-74.
- Grimaldi, D. [A.], and M. S. Engel. 2005.** Evolution of the insects. Cambridge University Press; Cambridge. 772 p.
- Hagen, H. 1861.** Synopsis of the Neuroptera of North America. With a list of South American species. *Smithsonian Miscellaneous Collections* 4(1): i-xx, 1-347.
- Hine, J. S. 1901a.** A review of the Panorpidae north of Mexico. *Bulletin of the Science Laboratories of Denison University* 11: 241-264, pl. LIX-LX.
- Hine, J. S. 1901b.** A review of the Panorpidae north of Mexico. *University Bulletin. Ohio State University, Series V* 7: 241-264, pl. LIX-LX.
- Kluge, N. J. 2004.** Larval leg structure of *Nannochorista* Tillyard, 1917 and characteristics of Mecoptera. *Russian Entomological Journal* 2003 12: 349-354.
- Machado, R. J. P., F. S. P. Godoi, and J. A. Rafael. 2009.** Neotropical Mecoptera (Insecta): New generic synonymies, new combinations, key to families and genera, and checklist of species. *Zootaxa* 2148: 27-38.
- Mickoleit, G. 2008.** Die Sperma-Auspreßvorrichtung der Nannochoristidae (Insecta: Mecoptera). *Entomologia Generalis* 31: 193-226.
- Minet, J., D.-Y. Huang, H. Wu, and A. Nel. 2010.** Early Mecopterida and the systematic position of the Microptysmatidae (Insecta: Endopterygota). *Annales de la Société Entomologique de France (N. S.)* 46: 262-270.
- [Novokshonov, V. G.] 1997.** [Early evolution of scorpionflies] (Insecta: Panorpidae). [Nauka, Moscow.] 140 p., plates I-IX. [in Russian]
- Penny, N. D. 1975.** Evolution of the extant Mecoptera. *Journal of the Kansas Entomological Society* 48: 331-350.
- Ren, D., C. C. Labandeira, J. A. Santiago-Blay, A. Rasnitsyn, CK Shih, A. Bashkuev, M. A. V. Logan, C. L. Hotton, and D. Dilcher. 2009.** A probable pollination mode before angiosperms: Eurasian, long-proboscid scorpionflies. *Science* 326: 840-847 + Supplement [online] available on URL: <http://www.sciencemag.org/cgi/content/full/326/5954/840/DC1>. (Last accessed 28 October 2010.)
- Schiefer, T. L., and J. C. Dunford. 2005.** New state record for *Merope tuber* Newman (Mecoptera: Meropeidae) in Alabama. *Journal of Entomological Science* 40: 471-473.
- Somma, L. A. 2010.** A state record for the Ocone scorpionfly, *Panorpa oconee* Byers (Mecoptera: Panorpidae), in Florida. *Insecta Mundi* 0135: 1-4.
- Somma, L. A., and J. C. Dunford. 2007.** Etymology of the Earwigfly, *Merope tuber* Newman (Mecoptera: Meropeidae): Simply dull or just inscrutable? *Insecta Mundi* 0013: 1-5.
- Somma, L. A., and J. C. Dunford. 2008.** Preliminary checklist of the Mecoptera of Florida: Earwigflies, hangingflies, and scorpionflies. *Insecta Mundi* 0042: 1-9.
- Somma, L. A., and J. C. Dunford. 2009a.** The Florida scorpionfly, *Panorpa floridana* Byers (Mecoptera: Panorpidae). Florida Department of Agriculture & Consumer Services, Division Plant Industry, Entomology Circular 420: 1-3. Also available on URL: <http://www.doacs.state.fl.us/pi/enpp/ento/entcirc/ent420.pdf>. (Last accessed 20 December 2010.)
- Somma, L. A., and J. C. Dunford. 2009b.** Records for *Bittacus* hangingflies and *Panorpa* scorpionflies (Mecoptera: Bittacidae and Panorpidae) in Florida. *Insecta Mundi* 0084: 1-5.
- Willmann, R. 1987.** The phylogenetic system of the Mecoptera. *Systematic Entomology* 12: 519-524.

Received November 17, 2010; Accepted February 1, 2011.