

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and
Plant Health Inspection Service

February 2007

BLACKBIRD USE OF WILDLIFE CONSERVATION SUNFLOWER PLOTS

George M. Linz

USDA/APHIS/WS National Wildlife Research Center, george_m_linz@yahoo.com

Jonathan M. Raetzman

North Dakota State University, Fargo

Heath M. Hagy

North Dakota State University, Fargo

H. Jeffrey Homan

USDA-APHIS-Wildlife Services

William J. Bleier

North Dakota State University, Fargo

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Environmental Sciences Commons](#)

Linz, George M.; Raetzman, Jonathan M.; Hagy, Heath M.; Homan, H. Jeffrey; and Bleier, William J., "BLACKBIRD USE OF WILDLIFE CONSERVATION SUNFLOWER PLOTS" (2007). *USDA National Wildlife Research Center - Staff Publications*. 707.

https://digitalcommons.unl.edu/icwdm_usdanwrc/707

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

BLACKBIRD USE OF WILDLIFE CONSERVATION SUNFLOWER PLOTS

GEORGE M. LINZ, USDA, Wildlife Services, National Wildlife Research Center, Bismarck, North Dakota, USA 58501

JONATHAN M. RAETZMAN, Department of Biological Sciences, North Dakota State University, Fargo, North Dakota, USA 58105

HEATH M. HAGY, Department of Biological Sciences, North Dakota State University, Fargo, North Dakota, USA 58105

H. JEFFREY HOMAN, USDA, Wildlife Services, National Wildlife Research Center, Bismarck, North Dakota, USA 58501

WILLIAM J. BLEIER, Department of Biological Sciences, North Dakota State University, Fargo, North Dakota, USA 58105

In North Dakota, annual blackbird damage to sunflower ranges from \$5-10 million. Blackbird damage to ripening sunflower has forced some growers to plant alternative crops. From 2004 to 2006, USDA-Wildlife Services cost-shared Wildlife Conservation Sunflower Plots (WCSP) with sunflower growers. The objective of the WCSP was to provide blackbirds an attractive nearby alternative food source to reduce damage to commercial fields. A secondary benefit of WCSP was to provide a safe-haven for other wildlife that frequently use shelterbelts and wetlands along the edges of sunflower fields. In 2004 (n=13), 2005 (n=21), and 2006 (n=25), sunflower damage in the WCSP's was 39%, 32%, and 60%, respectively. Damage in nearby commercial fields was 5% in 2004, 4% in 2005, and 18% in 2006. In 2006, dry conditions may have concentrated blackbirds into bigger roosts in the larger and deeper wetlands, contributing to higher levels of sunflower damage compared to 2004 and 2005. Lower levels of damage in 2004 and 2005 may have been caused by the greater availability of wetlands, which dispersed the flocks. We believe that avian use of WCSP was influenced by the nearness of shelterbelts, cattail-dominated wetlands, and contiguous blocks of commercial sunflower. We speculate that WCSP can reduce bird damage in nearby commercial fields. Based on these data, additional large-scale field tests appear to be warranted.