

2016

The Association of Exposure to Point-of-Sale Tobacco Marketing with Quit Attempt and Quit Success: Results from a Prospective Study of Smokers in the United States

Mohammad Siahpush

University of Nebraska Medical Center, msiahpush@unmc.edu

Raees A. Shaikh

University of Nebraska Medical Center, raees.shaikh@unmc.edu

Danielle Smith

Roswell Park Cancer Institute, Elm and Carlton Streets, Danielle.Smith@RoswellPark.org

Andrew Hyland

Roswell Park Cancer Institute, Elm and Carlton Streets, andrew.hyland@roswellpark.org

K. Michael Cummings

Medical University of South Carolina, cummingk@musc.edu

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/psychfacpub>

 Part of the [Psychology Commons](#)

Siahpush, Mohammad; Shaikh, Raees A.; Smith, Danielle; Hyland, Andrew; Cummings, K. Michael; Kessler, Asia Sikora; Dodd, Michael D.; Carlson, Les; Meza, Jane L.; and Wakefield, Melanie, "The Association of Exposure to Point-of-Sale Tobacco Marketing with Quit Attempt and Quit Success: Results from a Prospective Study of Smokers in the United States" (2016). *Faculty Publications, Department of Psychology*. 716.

<http://digitalcommons.unl.edu/psychfacpub/716>

This Article is brought to you for free and open access by the Psychology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Psychology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Mohammad Siahpush, Raees A. Shaikh, Danielle Smith, Andrew Hyland, K. Michael Cummings, Asia Sikora Kessler, Michael D. Dodd, Les Carlson, Jane L. Meza, and Melanie Wakefield



Article

The Association of Exposure to Point-of-Sale Tobacco Marketing with Quit Attempt and Quit Success: Results from a Prospective Study of Smokers in the United States

Mohammad Siahpush ^{1,*}, Raees A. Shaikh ¹, Danielle Smith ², Andrew Hyland ², K. Michael Cummings ³, Asia Sikora Kessler ¹, Michael D. Dodd ⁴, Les Carlson ⁵, Jane Meza ¹ and Melanie Wakefield ⁶

¹ College of Public Health, University of Nebraska Medical Center, 984365 Nebraska Medical Center, Omaha, NE 68198, USA; raees.shaikh@unmc.edu (R.A.S.); asia.sikora@unmc.edu (A.S.K.); jmeza@unmc.edu (J.M.)

² Roswell Park Cancer Institute, Elm and Carlton Streets | Buffalo, NY 14263, USA; Danielle.Smith@RoswellPark.org (D.S.); andrew.hyland@roswellpark.org (A.H.)

³ Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, 67 President Street, MSC 861, Charleston, SC 29425, USA; cummingk@musc.edu

⁴ Department of Psychology, University of Nebraska–Lincoln, 312 N 14th Street, Alexander West, Lincoln, NE 68588, USA; mdodd2@unl.edu

⁵ College of Business Administration, University of Nebraska–Lincoln, 312 N 14th Street, Alexander West, Lincoln, NE 68588, USA; lcarlson3@unl.edu

⁶ The Cancer Council Victoria, 615 St Kilda Road, Melbourne VIC 3094, Australia; melanie.wakefield@cancervic.org.au

* Correspondence: msiahpush@unmc.edu; Tel.: +1-402-559-3437; Fax: +1-402-559-3773

Academic Editors: Coral Gartner and Britta Wigginton

Received: 25 November 2015; Accepted: 2 February 2016; Published: 6 February 2016

Abstract: The aim was to assess the association of exposure to point-of-sale (POS) tobacco marketing with quit attempt and quit success in a prospective study of smokers in the United States. Data were collected via telephone-interview on exposure to POS tobacco marketing, sociodemographic and smoking-related variables from 999 smokers in Omaha, Nebraska, in the United States. Exposure to POS tobacco marketing was measured by asking respondents three questions about noticing pack displays, advertisements, and promotions in their respective neighborhoods stores. These three variables were combined into a scale of exposure to POS tobacco marketing. About 68% of the respondents participated in a six-month follow-up phone interview and provided data on quit attempts and smoking cessation. At the six-month follow-up, 39.9% of respondents reported to have made a quit attempt, and 21.8% of those who made a quit attempt succeeded in quitting. Exposure to POS marketing at baseline was not associated with the probability of having made a quit attempt as reported at the six-month follow-up ($p = 0.129$). However, higher exposure to POS marketing was associated with a lower probability of quit success among smokers who reported to have attempted to quit smoking at six-month follow-up ($p = 0.006$). Exposure to POS tobacco marketing is associated with lower chances of successfully quitting smoking. Policies that reduce the amount of exposure to POS marketing might result in higher smoking cessation rates.

Keywords: point-of-sale tobacco marketing; quit attempt; smoking cessation

1. Introduction

Tobacco is one of the most marketed products in the United States [1]. The 1998 Master Settlement Agreement (MSA), which imposed significant restrictions on tobacco marketing in most outdoor places, resulted in an increased focus of the tobacco industry on marketing activities at the point of sale (POS) [2–4]. In 2011, the tobacco industry spent \$7.5 billion on POS marketing [5] in three areas of (1) product displays; (2) advertisements; and (3) promotional and price incentives to consumers [3,6].

Despite the extensive amount of POS tobacco marketing that currently exists in the United States, very little research has addressed its impact on smoking cessation behavior. In a prospective study, Germain *et al.* examined the effect of “sensitivity” to POS tobacco displays on quitting behavior [7]. They conducted telephone surveys of 222 adult smokers in Australia at baseline and a follow-up 18 months later. Sensitivity to POS tobacco displays was measured with an index consisting of the following three variables: (1) the frequency of noticing tobacco displays; (2) unplanned purchasing behavior; and (3) deciding on brands based on POS displays. The results revealed that while sensitivity to POS marketing was not associated with making a quit attempt, lower sensitivity to POS marketing was associated with a higher probability of smoking cessation. In a qualitative study in New Zealand, Hoek *et al.* conducted semi-structured in-depth interviews with 20 participants who had attempted to quit smoking in the last six months [8]. At the time of the interview, 12 were still quit and eight had relapsed. The interviews suggested that tobacco product displays elicited emotional and physical reactions that undermined quit attempts. To our knowledge, there are no other studies that address how POS marketing influences smoking cessation. These studies only focused on a single aspect of POS tobacco marketing, namely cigarette pack displays. In the current research we employed a scale of exposure to POS tobacco marketing that included all three aspects of exposure to POS marketing (product display, advertisements, and promotional/price incentives) and examined its prospective association with quit attempt and quit success in a sample of adult smokers in a city in the United States.

2. Methods

2.1. Data and Design

This was a prospective study of a cohort of smokers in Omaha, Nebraska, who participated in a telephone interview at baseline and 6-month follow-up in the period between August 2013 and June 2014 [9]. At baseline we collected demographic, exposure to POS tobacco marketing, and smoking-related data on 999 participants using random digit dialing and placement of local advertisements in places such as the major daily newspaper and Craigslist to recruit volunteers. Our original plan was to use only random digit dialing, but this method proved to be more expensive than we had anticipated and, as a result, we sought other methods of recruitment. Those included in the study spoke English, were 18 years of age or older, had smoked more than 100 cigarettes in their life, and smoked five or more cigarettes a day at the time of the recruitment. Those who responded “never” to the following question were excluded from the study: “How often do you visit the stores in the neighborhood where you live? By stores, we mean such places as convenience stores, gas stations, grocery stores, supermarkets, drug stores, liquor stores, and tobacco stores. (never/sometimes/frequently/always)”. About 68% of respondents completed the 6-month follow-up survey and provided information on quitting behavior such as quit attempt and successful smoking cessation. The follow-up survey included questions on exposure to POS marketing as well, but we did not use these in the current research. The University of Nebraska Medical Center Institutional Review Board provided ethics approval for the study. The Institutional IRB approval number is 490-12-EP.

2.2. Measurement of Outcomes

Both engaging in a quit attempt and quit success were assessed at 6-month follow-up. Participants were regarded as having made a quit attempt if they responded affirmatively to the

question, “Have you made any attempts to quit smoking since we last spoke with you in (month of last interview)?” [10–12]. Those who had made a quit attempt were asked “Are you back smoking or are you still stopped?” [13–16]. If they indicated they were still stopped, we regarded them as having successfully quit smoking. About 98% of those who reported to be still stopped had quit one week or longer prior to the interview.

2.3. Measurement of Independent Variables

To measure the main independent variable POS cigarette marketing, the baseline survey asked each respondent the following questions [7,9,17,18]: “When you are in a store in your neighborhood, how often do you notice tobacco ads?”; “When you are in a store in your neighborhood, how often do you notice tobacco promotions such as special prices, multi-pack discounts, or free gift with purchase of cigarettes?”; and “When you are in a store in your neighborhood, how often do you notice cigarette pack displays?”. Possible responses to each question were: 1 = never, 2 = rarely, 3 = sometimes, 4 = often, 5 = always. The responses to the three questions were summed to create a scale of reported exposure to POS tobacco marketing with scores ranging from 3 (low marketing) to 15 (high marketing) and a Cronbach’s alpha of 0.6. This level of alpha has been considered satisfactory, especially in the early stages of research [19]. Research on the effect of cigarette marketing on smoking cessation is in its infancy. We have used this scale in previous research and have shown that it is associated with having urges to buy and impulse purchases of cigarettes [18].

Other independent variables, all measured at baseline, included urge to buy cigarettes, quit attempt history, motivation to quit, intention to quit, self-efficacy to quit, Heaviness of Smoking Index (HSI), sex, age, race/ethnicity, education, method of recruitment, and frequency of visits to stores in one’s neighborhood. Urge to buy cigarettes was measured with the question “When you are in a store in your neighborhood, how often do you get an urge to buy cigarettes? (1 = never; 2 = rarely; 3 = sometimes; 4 = often; 5 = always)” [17,18]. Quit attempt was measured with the question “Have you ever tried to quit smoking? (yes/no)” [20,21]. Motivation to quit was measured with the question “On a scale of one to ten, with “0” being “not at all” and “10” being “extremely”, how motivated you to give are up smoking right now?” Intention to quit was measured with the question “Are you planning to quit smoking: within the next month . . . within the next 6 months . . . sometime in the future, beyond 6 months . . . or not planning to quit?” [10,22,23]. Those who were planning to quit within the next six months were distinguished from those who were not. We measured self-efficacy by asking all respondents “If you decided to give up smoking completely in the next 6 months, how sure are you that you would succeed?” [22,23]. Respondents who said “somewhat sure”, “very sure” or “extremely sure” were distinguished from those who said “not at all sure” as having self-efficacy to quit. We dichotomized intention to quit and self-efficacy in order to be consistent with previous similar research using data from the United States, Canada, United Kingdom, and Australia and because the distribution of these variables was highly skewed [22,23]. HSI scores range from 0 to 6 and were calculated by summing the points for time to first cigarette after waking and number of cigarettes smoked per day. Time to first cigarette is scored as follows: <5 min = 3 points; 6–30 min = 2 points; 31–60 min = 1 point; and >60 min = 0 point. Number of cigarettes smoked per day is scored as follows: 1–10 = 0 point; 11–20 = 1 point; 21–30 = 2 points; and >31 = 3 points. Higher HSI scores indicate higher nicotine dependence [24,25]. Age was measured using the question “What is your age?”. To assess race/ethnicity, we asked the following two questions: “Are you Hispanic or Latino?” and “Which one or more of the following would you say is your race? White, Black or African American, Asian, Native Hawaiian or Other Pacific Islander, American Indian or Alaska Native, and other”. Education was measured using the question “What is the highest grade or year of school you completed?”.

2.4. Statistical Analysis

The quit attempt analysis included respondents who completed both baseline and follow-up interviews. The quit success analysis included only those who at follow-up indicated that they had made a quit attempt since the baseline interview. In both quit attempt and quit success analyses, we

omitted observations that had a missing value for any of the analysis variables. This constituted 4.2% of the sample in the quit attempt analysis (final sample $n = 649$) and 4.7% of the sample in the quit success analysis (final sample $n = 257$). We used logistic regression models to examine the association of POS tobacco marketing with quit attempt and quit success. Independent variables whose p -values were greater than 0.05 in the bivariate models were not included in the multivariate models.

In the multivariate analyses we assessed the severity of multicollinearity by examining the variance inflation factor (VIF) associated with each independent variable and found no evidence of an estimation problem arising from multicollinearity. Additionally, using Akaike Information Criterion (AIC) and Bayesian Information Criterion (BIC) for goodness-of-fit, we compared models with POS marketing as a continuous variable and models with POS marketing included as a categorical variable and found that the former provided a better fit to the data, supporting the assumption of a linear relationship between POS marketing and log odds of the outcomes.

3. Results

Table 1 shows the characteristics of the sample. Table 2 shows the unadjusted and adjusted effects of exposure to POS marketing and other independent variables on making a quit attempt. The unadjusted results indicated that exposure to POS marketing, urge to buy cigarettes, baseline quit attempt, motivation to quit, intention to quit, self-efficacy to quit, race/ethnicity, and method of recruitment had an association with quit attempt. In the adjusted model, exposure to POS marketing was not associated with quit attempt ($p = 0.185$). In this model, having made a quit attempt reported at baseline ($p = 0.003$), having a higher level of motivation to quit smoking ($p < 0.001$), and having an intention to quit at baseline ($p = 0.001$) were associated with a higher probability of making a quit attempt. In Supplementary Materials analyses (Table S1), we defined quit attempt as making an attempt that resulted in not smoking for longer than 24 hours. This definition did not change the conclusions of the study and there was no evidence of an association between POS marketing and quit attempt ($p = 0.209$).

Table 1. Sample characteristics for the quit attempt ($n = 649$) and quit success ($n = 257$) analysis.

Variables ^a	Quit Attempt	Quit Success
	% (n) or Mean (Range)	% (n) or Mean (Range)
Quit attempt (measured at 6-month follow-up)		
Attempted	39.91 (259)	–
Did not attempt	60.09 (390)	–
Quit success (measured at 6-month follow-up)		
Succeeded	–	21.79 (56)
Did not succeed	–	78.21 (201)
Exposure to Point-of-sale marketing (POS)	8.83 (3–15)	9.37 (3–15)
Urge to buy cigarettes	2.95 (1–5)	3.14 (1–5)
Baseline quit attempt		
Attempted	83.67 (543)	91.44 (235)
Did not attempt	16.33 (106)	8.56 (22)
Motivation	4.73 (1–10)	5.74 (1–10)
Intention to quit		
Yes	31.74 (206)	46.3 (119)
No	68.26 (443)	53.7 (138)
Self-efficacy to quit		
Yes	60.55 (393)	65.76 (169)
No	39.45 (256)	34.24 (88)
HSI	3.24 (1–6)	3.22 (1–6)
Sex		
Male	40.83 (265)	39.3 (101)
Female	59.17 (384)	60.7 (156)

Table 1. Cont.

Variables ^a	Quit Attempt	Quit Success
	% (n) or Mean (Range)	% (n) or Mean (Range)
Age		
18–39	21.73 (141)	24.12 (62)
40–54	37.75 (245)	40.08 (103)
55+	40.52 (263)	35.8 (92)
Race/ethnicity		
Non-Hispanic White	67.95 (441)	59.92 (154)
Other	32.05 (208)	40.08 (103)
Education		
High school graduate or below	49.15 (319)	47.47 (122)
At least some college	50.85 (330)	52.53 (135)
Method of recruitment		
Random digit dialing	50.4 (327)	43.97 (144)
Other	49.6 (322)	56.03 (113)
Frequency of visits to stores		
Sometimes	12.02 (78)	10.51 (27)
Frequently	38.06 (247)	36.58 (94)
Always	49.92 (324)	52.92 (136)

^a All variables were measured at baseline unless otherwise stated in the table.

Table 2. Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette marketing and other independent variables on the odds of making a quit attempt ($n = 649$).

Independent Variables ^a	Unadjusted Odds Ratio (95%CI)	<i>p</i> -Value	Adjusted ^b Odds Ratio (95%)	<i>p</i> -Value
POS cigarette marketing	1.09 (1.03–1.15)	<0.001	1.04 (0.98–1.11)	0.18
Urge to buy cigarettes	1.19 (1.06–1.34)	0.004	1.12 (0.97–1.29)	0.111
Baseline quit attempt		<0.001		0.003
Attempted	2.77 (1.7–4.54)		2.17 (1.29–3.65)	
Did not attempt	1.00		1.00	
Motivation	1.25 (1.17–1.32)	<0.001	1.14 (1.06–1.23)	<0.001
Intention to quit				0.001
Yes	3.05 (2.17–4.30)	<0.001	2.02 (1.36–3.01)	
No	1.00		1.00	
Self-efficacy to quit		<0.001		0.691
Yes	1.43 (1.03–1.98)		1.08 (0.75–1.55)	
No	1.00		1.00	
HSI	0.97 (0.81–1.15)	0.691	–	–
Sex		0.54		–
Male	0.9 (0.66–1.25)		–	
Female	1.00		–	
Age		0.187		–
18–39	1.00		–	
40–54	0.92 (0.61–1.4)		–	
55+	0.71 (0.67–1.08)		–	
Race/ethnicity		<0.001		0.033
Non-Hispanic White	1.00		1.00	
Other	1.84 (1.32–2.58)		1.5 (1.03–2.19)	
Education		0.395		–
High school graduate or below	1.00		–	
At least some college	1.15 (0.85–1.57)		–	
Method of recruitment		0.02		0.497
Random digit dialing	0.69 (0.5–0.94)		0.88 (0.61–1.27)	
Other	1.00		1.00	
Frequency of visits to stores		0.44		–
Sometimes	1.00		–	
Frequently	1.1		–	
Always	1.31		–	

^a All independent variables were measured at baseline; ^b Adjusted for the effect of variables with $p < 0.5$ in the unadjusted models.

Table 3 shows the unadjusted and adjusted effects of exposure to POS marketing and other independent variables on quit success. The unadjusted results indicated that exposure to POS marketing and self-efficacy to quit had an association with quit success. In the adjusted model, only exposure to POS marketing was associated with quit success ($p = 0.006$). A one point, increase in the scale of exposure to POS marketing was associated with a 12% decrease in the odds of quit success.

In additional Supplementary Materials analyses (Tables S2–S7), we examined the association of each of the three components of the POS marketing scale with quit attempt and quit success. Quit attempt was not associated with exposure to displays ($p = 0.275$), advertisements ($p = 0.583$), and promotions ($p = 0.202$). Quit success was associated with exposure to displays ($p = 0.013$) and advertisements ($p = 0.01$), but not with exposure to promotions ($p = 0.249$).

Table 3. Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette marketing and other independent variables on the odds of quit success ($n = 257$).

Independent Variables ^a	Unadjusted Odds Ratio (95%CI)	<i>p</i> -Value	Adjusted ^b Odds Ratio (95%)	<i>p</i> -Value
POS cigarette marketing	0.88 (0.8–0.96)	0.006	0.88 (0.8–0.96)	0.006
Urge to buy cigarettes	0.95 (0.76–1.19)	0.657	–	–
Baseline quit attempt		0.911		–
Attempted	0.94 (0.33–2.68)		–	
Did not attempt	1.00		–	
Motivation	1.07 (0.95–1.19)	0.251	–	–
Intention to quit		0.353		–
Yes	1.32 (0.73–2.4)		–	
No	1.00		–	
Self-efficacy to quit		0.044		0.052
Yes	1.97 (0.66–3.90)		1.99 (0.99–3.97)	
No	1.00		1.00	
HSI	0.93 (0.67–1.3)	0.686	–	–
Sex		0.538		–
Male	1.21 (0.66–2.2)		–	
Female	1.00		–	
Age		0.164		–
18–39	1.00		–	
40–54	1.6 (0.68–3.74)		–	
55+	2.2 (0.94–5.1)		–	
Race/ethnicity		0.089		–
Non-Hispanic White	1.00		–	
Other	0.58 (0.31–1.1)		–	
Education		0.9		–
High school graduate or below	1.00		–	
At least some college	0.96 (0.53–1.74)		–	
Method of recruitment		0.103		–
Random digit dialing	1.64 (0.9–2.98)		–	
Other	1.00		–	
Frequency of visits to stores		0.2		–
Sometimes	1.00		–	
Frequently	2.2		–	
Always	1.36		–	

^a All independent variables were measured at baseline. ^b Adjusted for the effect of variables with $p < 0.5$ in the unadjusted models.

4. Discussion

In this prospective study, we examined the association of exposure to POS cigarette marketing reported at baseline with quit attempt and quit success reported at a six-month follow up. We developed an exposure to POS marketing scale which included product displays, advertisements, and promotions. We found that while exposure to POS marketing was not associated with quit attempt, higher levels of exposure to POS marketing were associated with a lower probability of successfully quitting.

The results were consistent with Germain *et al.*'s study that showed sensitivity to POS cigarette display is not associated with quit attempts but is predictive of smoking cessation [7]. The results were also consistent with Hoek *et al.* qualitative study that suggested POS cigarette displays undermine smoking cessation [8]. Of note is our finding that exposure to POS marketing was the only variable among a set of smoking-related variables including HSI and sociodemographic factors that was associated with smoking cessation. Other studies have shown higher HSI to be associated with a lower probability of smoking cessation [7,10,11]. The effect of HSI in our analysis was in the expected direction but perhaps because of our small sample size in the quit success analysis, there was insufficient power to detect an effect. In relation to the effect of other independent variables on smoking cessation, our findings were in accordance with several other studies which reported no association between smoking cessation and sex, age, socioeconomic status, ethnicity, intention to quit, self-efficacy to quit, and quit attempt [7,10,11]. However, our results were not consistent with other studies that showed a higher probability of smoking cessation to be associated with older age [26], higher levels of education, [26–29] and non-Hispanic white race/ethnicity [26–29].

A strength of the study was its prospective design where the measurement of exposure took place prior to the assessment of the outcome; exposure to POS marketing was measured at baseline and quit attempt and quit success were measured at follow up. Another strength of the study was that it developed a scale of exposure to POS marketing which included its three aspects (displays, advertisements, and promotions), instead of merely focusing on exposure to POS cigarette displays as other previous studies of the relationship between marketing and smoking cessation have done [7,8]. A weakness of the study was that the sample was not a probability sample and it may not be completely representative of the smoking population. However, we note that the sociodemographic distribution of the sample was similar to the subsample of smokers in the center city of Nebraska Metropolitan Statistical Area in the Behavioral Risk Factor Surveillance System (BRFSS) [30]. For example, the mean age in our sample was 50.3 years in our sample and 53 years in BRFSS, and the percentage of respondents with a high school diploma or a lower level of education was 49.9 in our sample and 46.3 in BRFSS. Another weakness of the study is that we did not include independent variables such as poverty or financial stress that might influence both noticing cigarette promotions such as discounts and successful smoking cessation.[11,31] Finally, because we did not ask respondents from which stores in their neighborhoods they normally purchased cigarettes, we were not able to investigate whether the effect of marketing on smoking cessation varied by the type of store.

5. Conclusions

There is a large evidence base about the effectiveness of population-level tobacco control policies in increasing smoking cessation rates. Policies such as increasing the price of tobacco products [32], restricting smoking in public places [33–35], and anti-smoking mass media campaigns [36] have been shown by numerous studies to promote smoking cessation. The current study adds to this body of knowledge by suggesting that policies that reduce the amount of exposure to POS cigarette marketing might result in higher smoking cessation rates.

Supplementary Materials: The following are available online at www.mdpi.com/www.mdpi.com/1660-4601/13/2/203/s1. **Table S1.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette marketing and other independent variables on the odds of making a quit attempt (defined as making an attempt that resulted not smoking for longer than 24 h) ($n = 649$). **Table S2.** Logistic regression results for the effect of exposure

to point-of-sale (POS) cigarette pack displays and other independent variables on the odds of making a quit attempt ($n = 649$). **Table S3.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette advertisements and other independent variables on the odds of making a quit attempt ($n = 649$). **Table S4.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette promotions and other independent variables on the odds of making a quit attempt ($n = 649$). **Table S5.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette pack displays and other independent variables on the odds of quit success ($n = 257$). **Table S6.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette advertisements and other independent variables on the odds of quit success ($n = 257$). **Table S7.** Logistic regression results for the effect of exposure to point-of-sale (POS) cigarette promotions and other independent variables on the odds of quit success ($n = 257$).

Acknowledgments: Research reported in this publication was supported by National Institutes of Health (NIH) grant R01CA166156 and Food and Drug Administration Center for Tobacco Products (CTP). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the Food and Drug Administration.

Author Contributions: We assure the reader that all authors included on this paper fulfill the criteria of authorship. All have contributed to the conception and design, analysis and interpretation of data, drafting of the article, and revising it critically for important intellectual content. All authors approve the final version of the manuscript. In addition, we also assure the reader that there is no one else who fulfills the criteria but has not been included as an author. Mohammad Siahpush was instrumental in conceptualization of the research study, data analysis, and writing of the initial draft of the manuscript. Danielle Smith, Andrew Hyland, K. Michael Cummings, Michael D. Dodd, Les Carlson, Jane Meza and Melanie Wakefield contributed to the development of study, data analysis, and helped with writing the manuscript and editing it for final submission. Raees A. Shaikh helped with literature review, provided inputs for the methods section, and contributed to writing and editing the manuscript. Asia Sikora Kessler was involved with literature review and data analysis and provided input to the entire manuscript.

Conflicts of Interest: K. Michael Cummings has received grant funding from the Pfizer, Inc., to study the impact of a hospital based tobacco cessation intervention. K. Michael Cummings also receives funding as an expert witness in litigation filed against the tobacco industry. None of the other authors have conflicts to report.

References

1. American Lung Association. *Tobacco Industry Marketing*; Publisher: Chicago, IL, USA, 2014; Volume 2015.
2. Wakefield, M.A.; Terry-McElrath, Y.M.; Chaloupka, F.J.; Barker, D.C.; Slater, S.J.; Clark, P.I.; Giovino, G.A. Tobacco industry marketing at point of purchase after the 1998 MSA billboard advertising ban. *Am. J. Public Health* **2002**, *92*, 937–940. [[CrossRef](#)] [[PubMed](#)]
3. Lavack, A.M.; Toth, G. Tobacco point-of-purchase promotion: Examining tobacco industry documents. *Tob. Control* **2006**, *15*, 377–384. [[CrossRef](#)] [[PubMed](#)]
4. Loomis, B.R.; Farrelly, M.C.; Mann, N.H. The association of retail promotions for cigarettes with the master settlement agreement, tobacco control programmes and cigarette excise taxes. *Tob. Control* **2006**, *15*, 458–463. [[CrossRef](#)] [[PubMed](#)]
5. Federal Trade Commission. *Federal Trade Commission Cigarette Report for 2011*; Federal Trade Commission: Washington, DC, USA, 2013.
6. Bloom, P.N. Role of slotting fees and trade promotions in shaping how tobacco is marketed in retail stores. *Tobacco Control* **2001**, *10*, 340–344. [[CrossRef](#)] [[PubMed](#)]
7. Germain, D.; McCarthy, M.; Wakefield, M. Smoker sensitivity to retail tobacco displays and quitting: A cohort study. *Addiction* **2009**, *105*, 159–163. [[CrossRef](#)] [[PubMed](#)]
8. Hoek, J.; Gifford, H.; Pirikahu, G.; Thomson, G.; Edwards, R. How do tobacco retail displays affect cessation attempts? Findings from a qualitative study. *Tob. Control* **2010**, *19*, 334–337. [[CrossRef](#)] [[PubMed](#)]
9. Siahpush, M.; Shaikh, R.A.; Cummings, K.M.; Hyland, A.; Dodd, M.; Carlson, L.; Kessler, A.S.; Meza, J.; Wan, N.; Wakefield, M. The association of point-of-sale cigarette marketing with cravings to smoke: Results from a cross-sectional population-based study. *Tob. Control* **2015**. [[CrossRef](#)] [[PubMed](#)]
10. Hyland, A.; Borland, R.; Li, Q.; Yong, H.-H.; McNeill, A.; Fong, G.T.; O'Connor, R.J.; Cummings, K.M. Individual-level predictors of cessation behaviors among participants in the international tobacco control (ITC) policy evaluation study. *Tob. Control* **2006**, *15*. [[CrossRef](#)] [[PubMed](#)]
11. Siahpush, M.; Yong, H.H.; Borland, R.; Reid, J.L.; Hammond, D. Smokers with financial stress are more likely to want to quit but less likely to try or succeed: Findings from the international tobacco control (ITC) four country survey. *Addiction* **2009**, *104*, 1382–1390. [[CrossRef](#)] [[PubMed](#)]

12. Siahpush, M.; Borland, R.; Yong, H.H. Sociodemographic and psychosocial correlates of smoking-induced deprivation and its effect on quitting: Findings from the international tobacco control policy evaluation survey. *Tob. Control* **2007**, *16*. [[CrossRef](#)] [[PubMed](#)]
13. Nagelhout, G.E.; de Vries, H.; Fong, G.T.; Candel, M.J.; Thrasher, J.F.; van den Putte, B.; Thompson, M.E.; Cummings, K.M.; Willemsen, M.C. Pathways of change explaining the effect of smoke-free legislation on smoking cessation in the netherlands. An application of the international tobacco control conceptual model. *Nicot. Tob. Res.* **2012**, *14*, 1474–1482. [[CrossRef](#)] [[PubMed](#)]
14. Rennen, E.; Nagelhout, G.E.; van den Putte, B.; Janssen, E.; Mons, U.; Guignard, R.; Beck, F.; de Vries, H.; Thrasher, J.F.; Willemsen, M.C. Associations between tobacco control policy awareness, social acceptability of smoking and smoking cessation. Findings from the international tobacco control (ITC) Europe surveys. *Health Educ. Res.* **2014**, *29*, 72–82. [[CrossRef](#)] [[PubMed](#)]
15. IARC. *Iarc handbooks of Cancer Prevention, Tobacco Control, Volume 12: Methods for Evaluating Tobacco Control Policies*; International Agency for Research on Cancer: Lyon, France, 2008.
16. Cheong, Y.; Yong, H.-H.; Borland, R. Does how you quit affect success? A comparison between abrupt and gradual methods using data from the international tobacco control policy evaluation study. *Nicot. Tob. Res.* **2007**, *9*. [[CrossRef](#)] [[PubMed](#)]
17. Wakefield, M.; Germain, D.; Henriksen, L. The effect of retail cigarette pack displays on impulse purchase. *Addiction* **2008**, *103*, 322–328. [[CrossRef](#)] [[PubMed](#)]
18. Siahpush, M.; Shaikh, R.A.; Hyland, A.; Smith, D.; Kessler, A.S.; Meza, J.; Wan, N.; Wakefield, M. Point-of-sale cigarette marketing, urge to buy cigarettes, and impulse purchases of cigarettes: Results from a population-based survey. *Nicot. Tob. Res.* **2015**, *2015*. [[CrossRef](#)] [[PubMed](#)]
19. Nunnally, J.C.; Bernstein, I.H.; Berge, J.M.T. *Psychometric Theory*; McGraw-Hill: New York, NY, USA, 1967; Volume 226.
20. Sharma, A.; Szatkowski, L. Characteristics of smokers who have never tried to quit: Evidence from the british opinions and lifestyle survey. *BMC Public Health* **2014**, *14*. [[CrossRef](#)] [[PubMed](#)]
21. Hosking, W.; Borland, R.; Yong, H.-H.; Fong, G.; Zanna, M.; Laux, F.; Thrasher, J.; Lee, W.B.; Sirirassamee, B.; Omar, M. The effects of smoking norms and attitudes on quitting intentions in malaysia, thailand and four western nations: A cross-cultural comparison. *Psychol. Health* **2009**, *24*, 95–107. [[CrossRef](#)] [[PubMed](#)]
22. Siahpush, M.; Borland, R.; Yong, H.H.; Kin, F.; Sirirassamee, B. Socio-economic variations in tobacco consumption, intention to quit and self-efficacy to quit among male smokers in thailand and malaysia: Results from the nternational tobacco Control-South-East Asia (ITC-SEA) survey. *Addiction* **2008**, *103*, 502–508. [[CrossRef](#)] [[PubMed](#)]
23. Siahpush, M.; McNeill, A.; Borland, R.; Fong, G.T. Socioeconomic variations in nicotine dependence, self-efficacy and intention to quit across four countries: Findings from the international tobacco control policy evaluation survey. *Tob. Control* **2006**, *15*. [[CrossRef](#)] [[PubMed](#)]
24. Kozlowski, L.; Porter, C.Q.; Orleans, C.T.; Pope, M.A.; Heatherton, T.F. Predicting smoking cessation with self-reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug Alcohol Depend.* **1994**, *34*, 211–216. [[CrossRef](#)]
25. Heatherton, T.F.; Kozlowski, L.; Frecker, R.C.; Rickert, W.; Robinson, J. Measuring the heaviness of smoking: Using self-reported time to the first cigarette of the day and number of cigarettes smoked per day. *Br. J. Add.* **1989**, *84*, 791–799. [[CrossRef](#)]
26. Lee, C.W.; Kahende, J. Factors associated with successful smoking cessation in the United States, 2000. *Am. J. Public Health* **2007**, *97*, 1503–1509. [[CrossRef](#)] [[PubMed](#)]
27. Biener, L.; Hamilton, W.L.; Siegel, M.; Sullivan, E.M. Individual, social-normative, and policy predictors of smoking cessation: A multilevel longitudinal analysis. *Am. J. Public Health* **2010**, *100*, 547–554. [[CrossRef](#)] [[PubMed](#)]
28. Reid, J.L.; Hammond, D.; Boudreau, C.; Fong, G.T.; Siahpush, M.; Collaboration, I.T.C. Socioeconomic disparities in quit intentions, quit attempts, and smoking abstinence among smokers in four western countries: Findings from the international tobacco control four country survey. *Nicot. Tob. Res. Off. J. Soc. Res. Nicot. Tob.* **2010**, *12*, S20–S33. [[CrossRef](#)] [[PubMed](#)]
29. Centers for Disease Control and Prevention. Quitting smoking among adults—United States, 2001–2010. *MMWR. Morbid. Mortal. Wkly. Rep.* **2011**, *60*, 1513–1519.

30. Centers for Disease Control and Prevention. *Smart: Bfss City and County Data*; Centers for Disease Control and Prevention: Atlanta, GA, USA, 2013; Volume 2015.
31. Siahpush, M.; Carlin, J.B. Financial stress, smoking cessation and relapse: Results from a prospective study of an Australian national sample. *Addiction* **2006**, *110*, 121–127. [[CrossRef](#)] [[PubMed](#)]
32. Chaloupka, F.J.; Straif, K.; Leon, M.E. Effectiveness of tax and price policies in tobacco control. *Tob. Control* **2011**, *20*, 235–238. [[CrossRef](#)] [[PubMed](#)]
33. Farkas, A.J.; Gilpin, E.A.; Distefan, J.M.; Pierce, J.P. The effects of household and workplace smoking restrictions on quitting behaviours. *Tob. Control* **1999**, *8*, 261–265. [[CrossRef](#)] [[PubMed](#)]
34. Borland, R.; Owen, N.; Hocking, B. Changes in smoking behavior after a total workplace smoking ban. *Austr. J. Public Health* **1991**, *15*, 130–134. [[CrossRef](#)]
35. Fowkes, F.J.; Stewart, M.C.; Fowkes, F.G.; Amos, A.; Price, J.F. Scottish smoke-free legislation and trends in smoking cessation. *Addiction* **2008**, *103*, 1888–1895. [[CrossRef](#)] [[PubMed](#)]
36. National Cancer Institute. *The Role of the Media in Promoting and Reducing Tobacco Use. Tobacco Control Monograph No. 19*; Davis, R.M., Gilpin, E.A., Loken, B., Viswanath, K., Wakefield, M.A., Eds.; U.S. Department of Health and Human Services, National Institute of Health, National Cancer Institute: Bethesda, MD, USA, 2008.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).