

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

3-23-2012

First record of *Megischus brunneus* Cresson, 1865 (Hymenoptera: Stephanidae) from Hispaniola, the Antilles

Julio A. Genaro

Florida State Collection of Arthropods, polimita@hotmail.com

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

Genaro, Julio A., "First record of *Megischus brunneus* Cresson, 1865 (Hymenoptera: Stephanidae) from Hispaniola, the Antilles" (2012). *Insecta Mundi*. 729.

<https://digitalcommons.unl.edu/insectamundi/729>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0216

First record of *Megischus brunneus* Cresson, 1865 (Hymenoptera:
Stephanidae) from Hispaniola, the Antilles

Julio A. Genaro
Florida State Collection of Arthropods
1911 SW 34th Street
Gainesville, FL 32608-1268, USA

Date of Issue: March 23, 2012

Julio A. Genaro

First record of *Megischus brunneus* Cresson, 1865 (Hymenoptera: Stephanidae) from Hispaniola, the Antilles
Insecta Mundi 0216: 1-3

Published in 2012 by

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 U. S. A.

<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Production editor: Michael C. Thomas, Brian Armitage, and Ian Stocks

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

Spanish editors: Julieta Brambila, Angélico Asenjo

Printed copies (ISSN 0749-6737) deposited in libraries of:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and AgriFood Canada, Ottawa, ON, Canada

The Natural History Museum, London, Great Britain

Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD mailed to all members at end of year.

Florida Center for Library Automation: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

Author instructions available on the Insecta Mundi page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

First record of *Megischus brunneus* Cresson, 1865 (Hymenoptera: Stephanidae) from Hispaniola, the Antilles

Julio A. Genaro

Florida State Collection of Arthropods
1911 SW 34th Street
Gainesville, FL 32608-1268, USA
polimita@hotmail.com

Abstract. The occurrence of *Megischus brunneus* Cresson (Hymenoptera: Stephanidae) is recorded for the first time from Hispaniola, the Antilles. The species was previously known from southern Florida and Cuba. This finding further demonstrates the similarities between the Cuban and Hispaniolan biota.

Key words. Hymenoptera, Stephanidae, *Megischus brunneus*, Hispaniola, first record.

Resumen. Se registra por primera vez la presencia de *Megischus brunneus* Cresson (Hymenoptera: Stephanidae) para la Hispaniola. Esta especie se conocía anteriormente del sur de la Florida y Cuba. Este hallazgo constituye otro ejemplo de las similitudes de la biota cubana e hispaniolana.

Palabras claves. Hymenoptera, Stephanidae, *Megischus brunneus*, Hispaniola, primer registro.

Introduction

The only moderately diverse family Stephanidae (Hymenoptera) contains species of solitary idiobiont ectoparasitoids of wood boring insect larvae (332 species worldwide; see Aguiar and Johnson 2003; Aguiar 2006a, b, c). Stephanid wasps are remarkable insects, with a slender and richly sculptured body, highly modified hind legs, and a somewhat spherical head - set out on a long neck - which bears a crown of tubercles around the median ocellum (Aguiar 2006a, b).

Megischus Brullé is a cosmopolitan genus (Aguiar 2004; Hong et al. 2011). Aguiar and Johnson (2003) provided data on geographical distribution, biology and a key to the North American species of *Megischus*, including the Antilles. Aguiar (2006c) subsequently also revised the Mexican species. Aguiar (2006c) revised the Mexican species, describing new ones and offering a key.

Megischus brunneus Cresson can be recognize by having a pronotum with four sharp transverse carinae, and mesopleuron and prosternum with dense foveolae, each separated by less than their diameter (Aguiar and Johnson 2003). The species was previously known from the United States (Southern Florida) and Cuba (Townes 1949; Alayo 1972, 1973; Aguiar and Johnson 2003; Aguiar 2004). This study provides the first record of the occurrence of *M. brunneus* for Hispaniola, the Antilles.

Codens for collections mentioned here are as follows: Department of Biology, University of Puerto Rico at Mayagüez (UPRM) (located in Mayagüez); personal collection of the author (JAGA).

Results

In November of 2003 evidence was gathered on the occurrence of this species in the Dominican Republic. This was possible because of a direct sighting and capture of a single female in the town of Oviedo, Pedernales, in the backyard of the Jaragua Group facilities. Unfortunately the material was subsequently stolen. Nevertheless, further collecting in Jaragua yielded another female and another female was found in the UPRM collection, which validates this new record.

The specimen collected at Oviedo lake was resting on a mangrove's trunk (Fig. 1). The habitat (Fig. 2) is characterized by the following mangrove species: *Rhizophora mangle* (L.) Rhizophoraceae, *Avicennia germinans* (L.) L. Avicenniaceae, *Conocarpus erectus* L. Combretaceae and *Laguncularia racemosa* (L.) Gaertn. f. Combretaceae. Similar habitats have been reported elsewhere for this species (Aguiar and



Figures 1-2. *Megischus brunneus*. 1) Female resting on a mangrove trunk of *Laguncularia racemosa* at Oviedo Lake, Pedernales. 2) Habitat at Oviedo Lake constituted by salt marsh vegetation and mangroves

Johnson 2003). The distributional records (Fig. 3) show that *M. brunneus* is present in dry areas of the southwestern Dominican Republic.

Material examined. HISPANIOLA, Dominican Republic, Oviedo lake, Pedernales, Jaragua National Park, 21.viii.2010, coll. J. A. Genaro (female, JAGA) (body length, excluding ovipositor 29.2 mm; hind wing length 14.8 mm); Dominican Republic, San José de Ocoa, ca 3km N along road towards El Canal, empty riverbed, night coll. (Hg and UV lights), 18°33'53"N 070°30'57"W, 500 m asl, 6.vi.2008, Franz lab-DR 08 (female, UPRM) (body length, excluding ovipositor 28 mm; hind wing length 13.5 mm).

Megischus brunneus likely originated on the continent and dispersed to Cuba, perhaps via the Bahamas, as Cuba and the Florida have never been connected by emerged lands. Portions of Cuba and Hispaniola were connected during Latest Eocene-Early Oligocene (35-33 Ma), a time of general uplift and a maximum area of subaerial land (Iturralde-Vinent 2006). This offered favorable conditions for sharing biota. After the Late Oligocene (29-27 Ma) isolation of land areas resulting in the subdivision and separation of block-terraces previously acting as continuous landmasses (Iturralde-Vinent 2006). Perhaps vicariance has isolated populations of *M. brunneus* creating the present-day distribution patterns.

Acknowledgments

I am grateful to Victor González who encouraged the study of insects and generously provided logistic support for field trips during several years. I thank the personnel of Jaragua group for logistical support of expeditions and especially Ivonne Arias and Héctor Andujar also for facilitation of collecting permits and use of Jaragua National Park infrastructure. I thank to my colleagues G. Alayón, E. Gutiérrez and A. Pérez-Asso for collaboration during several years of expeditions. Thanks to Nico Franz for facilitating the access to the UPRM insect collection to study the specimen. Cooperation of Secretaría de Estado de Medio Ambiente y Recursos Naturales for obtaining field work permits is greatly appreciated. Fabiola Areces (University of Puerto Rico at Río Piedras) provided identifications and information about vegetation. Carlos Suriel and Celeste Mir (Museo Nacional de Historia Natural de Santo Domingo) always supported with pleasure our research in the island and permitted the use of museum installations. I extend special thanks to James Carpenter, Nico Franz and José L. Fernández for reviewing the manuscript and offering valuable suggestions.




Figure 3. New locality records for *Megischus brunneus* from Hispaniola.

Literature Cited

- Alayo, P. 1972.** Introducción al estudio de los himenópteros de Cuba. I. Superfamilias Tenthredinoidea, Siricoidea, Ichneumonoidea. Escuela de Agronomía, Universidad Central de las Villas; Cuba. 71 p.
- Alayo, P. 1973.** Catálogo de los himenópteros de Cuba. Ed. Pueblo y Educación; La Habana. 218 p.
- Aguiar, A. P. 2004.** World catalog of the Stephanidae (Hymenoptera: Stephanoidea). *Zootaxa* 753: 1-120.
- Aguiar, A. P. 2006a.** Superfamilia Stephanoidea y familia Stephanidae. p. 273- 278. *In*: M. Sharkey and F. Fernández (Eds.). Introducción a los Hymenoptera de la Región Neotropical. Sociedad Colombiana de Entomología y Universidad Nacional de Colombia; Bogotá. 894 p.
- Aguiar, A. P. 2006b.** Stephanidae. p. 205-209. *In*: P. E. Hanson and I. D. Gauld (Eds.). Los himenópteros del Neotrópico. *Memoirs of the American Entomological Institute* 77: 1-994.
- Aguiar, A. P. 2006c.** The Stephanidae (Hymenoptera) of Mexico, with description new species and key to western *Foenatopus* Smith. *Zootaxa* 1186: 1-56.
- Aguiar, A. P., and N. F. Johnson. 2003.** Stephanidae (Hymenoptera) of America North of Mexico. *Proceedings of the Entomological Society of Washington* 105(2): 467-483.
- Hong, C, C. van Achterberg, and Z. Xu. 2011.** A revision of the Chinese Stephanidae (Hymenoptera, Stephanoidea). *ZooKeys* 110: 1-108.
- Iturralde-Vinent, M. A. 2006.** Meso-Cenozoic Caribbean Paleogeography: Implications for the Historical Biogeography of the Region. *International Geology Review* 48: 91-827.
- Townes, H. 1949.** The Nearctic species of the family Stephanidae (Hymenoptera). *Proceedings of the United States Natural Museum* 99 (3243): 361-370.

Received December 6, 2011; Accepted February 28, 2012.

