

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Nebraska Beef Cattle Reports

Animal Science Department

---

2013

## Statistics Used in the Nebraska Beef Report and Their Purpose

Follow this and additional works at: <https://digitalcommons.unl.edu/animalscinbcr>

---

"Statistics Used in the Nebraska Beef Report and Their Purpose" (2013). *Nebraska Beef Cattle Reports*. 752.

<https://digitalcommons.unl.edu/animalscinbcr/752>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Beef Cattle Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Statistics Used in the Nebraska Beef Report and Their Purpose

The purpose of beef cattle and beef product research at University of Nebraska–Lincoln is to provide reference information that represents the various populations (cows, calves, heifers, feeders, carcasses, retail products, etc.) of beef production. Obviously, the researcher cannot apply treatments to every member of a population; therefore, he/she must sample the population. The use of statistics allows the researcher and readers of the *Nebraska Beef Report* the opportunity to evaluate separation of random (chance) occurrences and real biological effects of a treatment. Following is a brief description of the major statistics used in the beef report. For a more detailed description of the expectations of authors and parameters used in animal science see *Journal of Animal Science Style and Form* at: <http://jas.fass.org/misc/ifora.shtml>.

- **Mean** — Data for individual experimental units (cows, steers, steaks) exposed to the same treatment are generally averaged and reported in the text, tables and figures. The statistical term representing the average of a group of data points is mean.
- **Variability** — The inconsistency among the individual experimental units used to calculate a mean for the item measured is the variance. For example, if the ADG for *all* the steers used to calculate the mean for a treatment is 3.5 lb then the variance is zero. But, this situation never happens! However, if ADG for individual steers used to calculate the mean for a treatment range from 1.0 lb to 5.0 lb, then the variance is large. The variance may be reported as standard deviation (square root of the variance) or as standard error of the mean. The standard error is the standard deviation of the mean as if we had done repeated samplings of data to calculate multiple means for a given treatment. In most cases treatment means and their measure of variability will be expressed as follows:  $3.5 \pm 0.15$ . This would be a mean of 3.5 followed by the standard error of the mean of 0.15. A helpful step combining both the mean and the variability from an experiment to conclude whether the treatment results in a real biological effect is to calculate a 95% confidence interval. This interval would be twice the standard error added to and subtracted from the mean. In the example above, this interval is 3.2-3.8 lb. If in an experiment, these intervals calculated for treatments of interest overlap, the experiment does not provide satisfactory evidence to conclude that treatments effects are different.
- **P Value** — Probability (*P Value*) refers to the likelihood the observed differences among treatment means are due to chance. For example, if the author reports  $P \leq 0.05$  as the significance level for a test of the differences between treatments as they affect ADG, the reader may conclude there is less than a 5% chance the differences observed between the means are a random occurrence and the treatments do not affect ADG. Hence we conclude that, because this probability of chance occurrence is small, there must be difference between the treatments in their effect on ADG. It is generally accepted among researchers when *P* values are less than or equal to 0.05, observed differences are deemed due to important treatment effects. Authors occasionally conclude that an effect is significant, hence real, if *P* values are between 0.05 and 0.10. Further, some authors may include a statement indicating there was a “tendency” or “trend” in the data. Authors often use these statements when *P* values are between 0.10 and 0.15, because they are not confident the differences among treatment means are real treatment effects. With *P* values of 0.10 and 0.15 the chance random sampling caused the observed differences is 1 in 10 and 1 in 6.7, respectively.
- **Linear & Quadratic Contrasts** — Some articles contain linear (L) and quadratic (Q) responses to treatments. These parameters are used when the research involves increasing amounts of a factor as treatments. Examples are increasing amounts of a ration ingredient (corn, by-product, or feed additive) or increasing amounts of a nutrient (protein, calcium, or vitamin E). The L and Q contrasts provide information regarding the shape of the response. Linear indicates a straight line response and quadratic indicates a curved response. *P*-values for these contrasts have the same interpretation as described above.
- **Correlation (r)** — Correlation indicates amount of linear relationship of two measurements. The correlation coefficient can range from -1 to 1. Values near zero indicate a weak relationship, values near 1 indicate a strong positive relationship, and a value of -1 indicates a strong negative relationship.



# Animal Science

<http://animalscience.unl.edu>

**Curriculum:** The curriculum of the Animal Science Department at the University of Nebraska–Lincoln is designed so that each student can select from a variety of options oriented to specific career goals in professions ranging from animal production to veterinary medicine. With unique opportunities to double major in **Grazing Livestock Systems** (<http://gls.unl.edu>) or complete the **Feedlot Management Internship Program** (<http://feedlot.unl.edu/intern>)

## Careers:

Animal Health  
Banking and Finance  
Animal Management  
Consultant

Education  
Marketing  
Technical Service  
Meat Processing

Meat Safety  
Quality Assurance  
Research and Development  
Veterinary Medicine

**Scholarships:** The Animal Science Department also offers scholarships to incoming freshmen and upperclassmen. The department awards over \$30,000 each year to Animal Science students.

ABS Global Scholarship  
Baltzell-Agri-Products, Inc. Scholarship  
Maurice E. Boeckenhauer Memorial  
Scholarship  
Mike Cull Judging and Activities Scholarship  
Don Geweke Memorial Award  
Parr Young Senior Merit Award  
Nebraska Pork Producers Association  
Scholarship  
Waldo Family Farms Scholarship  
Frank and Mary Bruning Scholarship  
Art and Ruth Raun Scholarship  
Animal Science Department Freshman  
Scholarship  
Feedlot Management Scholarship  
Robert Boeckenhauer Memorial Scholarship  
Burnell Scholarship Fund  
Doane Scholarship  
Lincoln Coca-Cola Bottling Company  
Scholarship.

William J. and Hazel J. Loeffel Scholarship  
Nutrition Service Associates Scholarship  
Parr Family Student Support Fund  
Chris and Sarah Raun Memorial Scholarship  
Walter A. and Alice V. Rockwell Scholarship  
Standard Manufacturing Co. Scholarship  
Max and Ora Mae Stark Scholarship  
D.V. and Ernestine Stephens Memorial  
Scholarship  
Dwight F. Stephens Scholarship  
Arthur W. and Viola Thompson Scholarship  
Thomas H. Wake, III Scholarship  
Frank E. Card Scholarship  
Derrick Family Scholarship  
G. H. Francke Livestock Judging Scholarship  
Eric Peterson Memorial Award  
Winkler Memorial Livestock Judging  
Scholarship