

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Papers and Publications in Animal
Science

Animal Science Department

2013

REVIEW: Life-cycle, total-industry genetic improvement of feed efficiency in beef cattle: Blueprint for the Beef Improvement Federation

Merlyn K. Nielsen

University of Nebraska-Lincoln, mnielsen1@unl.edu

M. D. MacNeil

Delta G, Miles City

J. C. M. Dekkers

Iowa State University

D. H. Crews Jr.

Colorado State University

T. A. Rathje

Danbred North America

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/animalscifacpub>



Part of the [Genetics and Genomics Commons](#), and the [Meat Science Commons](#)

Nielsen, Merlyn K.; MacNeil, M. D.; Dekkers, J. C. M.; Crews, D. H. Jr.; Rathje, T. A.; Enns, R. Mark; and Weaber, R. L., "REVIEW: Life-cycle, total-industry genetic improvement of feed efficiency in beef cattle: Blueprint for the Beef Improvement Federation" (2013). *Faculty Papers and Publications in Animal Science*. 804.

<https://digitalcommons.unl.edu/animalscifacpub/804>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Merlyn K. Nielsen, M. D. MacNeil, J. C. M. Dekkers, D. H. Crews Jr., T. A. Rathje, R. Mark Enns, and R. L. Weaber

REVIEW

REVIEW: Life-cycle, total-industry genetic improvement of feed efficiency in beef cattle: Blueprint for the Beef Improvement Federation¹

M. K. Nielsen,* M. D. MacNeil,† J. C. M. Dekkers,‡ D. H. Crews Jr.,§ T. A. Rathje,#
R. M. Enns,§² and R. L. Weaber||

*Department of Animal Science, University of Nebraska, Lincoln 68583; †Delta G, Miles City, MT 59301; ‡Department of Animal Science, Iowa State University, Ames 50011; §Department of Animal Sciences, Colorado State University, Fort Collins 80523; #Danbred North America, Columbus, NE 68601; and ||Department of Animal Sciences, Kansas State University, Manhattan 66506

ABSTRACT

On a life-cycle basis, beef animals are able to consume large amounts of low-cost, low-quality forages relative to higher-cost concentrates compared with pigs and chickens. However, of the 3, beef is still more expensive to produce on a cost-per-edible pound basis. Accordingly, there is need for genetic programs and management changes that will improve efficiency, sustainability, and profitability of beef production. Options include improving reproductive rate, reducing feed used for maintenance, or both, while not reducing output. A goal for improving efficiency of feed utilization is to reduce the amount or proportion of feed

used for maintenance. Such reduction is a target for genetic improvement, but such a goal does not include defining a single measure of efficiency. A single efficiency measure would likely lead to single-trait selection and not account for any potentially antagonistic effects on other production characteristics. Because we are not able to explain all variation in individual-animal intake from only knowledge of BW maintained and level of production, measuring feed intake is necessary. Therefore, our recommendation is that national cattle evaluation systems analyze feed intake as an economically relevant trait with incorporation of appropriate indicator traits for an EPD for feed intake requirements that could then be used in a multiple-trait setting such as in a selection index. With improvements in technology for measurement of feed intake, individual measures of feed intake should continually be collected to facilitate development of genetic

predictors that enhance accuracy of prediction of progeny differences in national cattle evaluations.

Key words: beef cattle, feed utilization, intake

INTRODUCTION

Beef, as a protein source for humans, has 2 major positive characteristics relative to pork and chicken: 1) consumers, on average, place greater preference on beef in its eating characteristics and 2) beef animals, on an industry-wide life-cycle basis, consume large amounts of lower-cost forages as compared with higher-cost concentrates. Although these positive characteristics exist, beef production still needs to improve cost per unit of product because it has greater cost per edible pound than does chicken and pork. If one compares edible

¹ The development of this commentary was supported by the Beef Improvement Federation.

² Corresponding author: Mark.Enns@Colostate.edu

product per unit of feed energy input, beef production is about one-third as efficient as pork production and about one-fifth to one-sixth as efficient as broiler production (adapted from Dickerson, 1978). Greatly lower reproduction per breeding female in cattle is a major contributor to the inefficiency, and adding the consumer-desired intramuscular fat in beef contributes to slaughter beef animals having greater total-carcass waste fat compared with slaughter pigs and broilers.

Implementing genetic programs and management changes that can improve efficiency of beef production requires answers to several questions. Some of these questions follow, and our goal in this paper is to provide answers to these questions, based on current knowledge. From an industry-wide perspective, what are the opportunities for improving efficiency of feed utilization? What can we learn from the pork and broiler industries in how they have approached genetic improvement of efficiency of feed utilization? Are there potential antagonisms between feed utilization or efficiency measurements and other economically relevant traits in beef cattle? What phenotypic and genomic data collections are warranted, and how will these be incorporated into National Cattle Evaluation programs? Where are the holes in our knowledge base, and what are the needs for future research to generate answers?

REVIEW AND DISCUSSION

Do We Need to Measure Feed?

Efficiency has been conventionally expressed as the ratio of output per unit of input. However, expressing efficiency in a linear form as output minus input has better statistical properties and comes closer to economic measures such as net return (value of output minus cost of input). If we express feed efficiency of the beef life cycle on an average dam basis and in linear form, we have the following (adapted from Dickerson, 1970):

$$\begin{aligned} & (\text{Dam BW} \times \text{Lean Value of Dam} \\ & + \text{No. Progeny} \times \text{Progeny BW} \\ & \quad \times \text{Lean Value of Progeny}) \\ & - (\text{Dam Feed} \times \text{Value of Feed for} \\ & \text{Dam} + \text{No. Progeny} \times \text{Progeny Feed} \\ & \quad \times \text{Value of Feed for Progeny}). \end{aligned}$$

Note, there is no requirement that the value terms be expressed in monetary units. They could equally well be expressed in biological units (e.g., kcal) to reflect biological efficiency.

In the positive income component we have the output from harvesting the dam (or fraction of the dam accounting for death loss) and from harvesting progeny (again, accounting for death loss); these are multiplied by different per unit prices to obtain the total value output. The negative feed cost component accounts for the input of feed energy, where we can account for different feedstuffs in the calculation of energy. The number of progeny per dam is in both components, and thus, increasing number of progeny will increase efficiency. By simply increasing number of progeny per dam through either selection, heterosis from crossing, or better management, we will increase efficiency of production. We do not need to measure feed intake to get this improvement in feed efficiency.

If we look at feed efficiency of a single animal, we also find that there are possible improvements in efficiency that can be achieved again without measurement of feed intake. To visualize this, first imagine that we can separate feed intake, at least conceptually, into 1) feed required to meet maintenance requirements (**M**, basal metabolism, tissue repair, thermal regulation, locomotor activity, and so on) or the energy required for keeping BW constant; 2) feed required to create new product (**P**, e.g., growth, milk, new offspring); and 3) feed that goes unused (**U**, waste products). For a growing calf, efficiency can be shown simply as

$$\begin{aligned} & \text{Calf BW Gain} \times \text{Calf BW Value} \\ & - (\text{Feed}_M + \text{Feed}_P + \text{Feed}_U) \\ & \quad \times \text{Feed Value.} \end{aligned}$$

For a pair of calves with the *same* starting and ending BW but with one animal gaining BW more quickly, thus requiring fewer days and less *maintenance* to reach market BW, the faster-growing calf would be more efficient. This can occur with *no* difference in efficiency of feed use for either maintenance or creation of new product; it is “all mathematical.” Similarly, with an improvement in reproduction, there is no need to measure feed intake to capitalize on methods to improve efficiency. The same would be true for an individual cow; if there is more output per day and no difference in cow size and in partial costs for maintenance and for production, then the cow with a greater rate of output will be the more efficient.

For a reproducing cow herd, we can express efficiency based on the BW of calf and cull cow as the summed outputs, and total feed intake for the 2 production components as the feed costs. This gets a bit more complicated compared with the growing calf example above. But, we can express this as

$$\begin{aligned} & [\text{Calf BW} \times \text{Calf BW Value} \\ & + (\text{Culling Rate} \times \text{Cull Cow BW} \\ & \times \text{Cow BW Value})] - [\text{Feed}_M(\text{cow}) \\ & + \text{Feed}_P(\text{cow}) + \text{Feed}_U(\text{cow})] \\ & \times \text{Cow Feed Value} - [\text{Feed}_M(\text{calf}) \\ & + \text{Feed}_P(\text{calf}) + \text{Feed}_U(\text{calf})] \\ & \times \text{Calf Feed Value} - [\text{Feed}_M(\text{heifer}) \\ & + \text{Feed}_P(\text{heifer}) + \text{Feed}_U(\text{heifer})] \\ & \quad \times \text{Heifer Feed Value.} \end{aligned}$$

So again, there is 1) feed for maintenance, 2) feed for production, and 3) feed that is wasted. So, one goal for improving efficiency of feed utilization, whether with a growing calf in a feedlot or with a reproducing cow and calf in a cow herd, must be to reduce

the feed being used for maintenance, while not reducing output. This also means that we could lose efficiency if we reduce rate of output, and hence reduce feed above maintenance required to produce output. Thus, instead of focusing on single traits, yearling bull buying decisions must consider the multiple-trait associations of feed intake and the implications of making selection decisions in the multiple-trait sense based mainly on data collected from growing bulls, especially with regard to the replacement daughters of selected bulls.

Thus, if the size of animals, rates of product formation (growth, milk), and reproduction are known, then why is it necessary to measure feed intake? Feed measurement is costly because of the need for special facilities and equipment and because of labor. To complicate matters, this costly measurement is amenable for high-energy diets but not for low-energy, high-roughage diets, in particular, pasture-based systems. The reproducing cow herd, including calves to weaning, consumes the greater fraction of annual feed energy required for beef production, as compared with calves grown from weaning to slaughter. The reproducing cow herd consumes mostly roughages, thus any measurement of feed intake of a cow herd is not feasible at this time. We can, however, use research results to formulate appropriate multiple-trait index tools that account for synergisms and antagonisms that may exist among feed intake and other economically relevant traits.

The reason why we might consider measuring feed intake is because we cannot explain all the variation in individual-animal feed intake from simply knowledge of BW maintained and level of production. Animals differ in their ability to digest feedstuffs and their ability to transform feed energy to meet these needs. Within these, the main deficiency in being able to explain feed intake is in predicting the cost of meeting maintenance requirements, adjusted for body size. As noted above, maintenance includes all energy costs to hold BW

and body energy content constant in the particular production situation. During extremely cold ambient conditions, maintenance for a given animal will be greater than for more moderate ambient conditions. Maintenance in an extensive, open grazing situation will be greater than in a confined feeding situation. From a total life-cycle perspective, energy costs for maintenance are estimated to be about 70% of the total energy intake in the beef production system.

Thus, reducing energy for maintenance, while accounting for possible negative effects on other performance characteristics, becomes a clear target for genetic improvement programs. The focus of these programs should not be to define a single measurement of efficiency, which may lead to inappropriate use of single-trait selection, but to define optimal measures that conform to the marketing practices (e.g., profitability) in the industry.

We can write an expression for feed intake that is unique for each animal in a defined production scenario: Feed Intake = Feed for Maintenance + Feed Above Maintenance for Production. This expression can be expanded to

$$\text{Feed Intake} = b_M \times (\text{BW})^{0.75} + b_P \\ \times (\text{Amount of Production}) + e.$$

Both b_M and b_P are partial efficiencies, as they represent the amount of feed required per unit of metabolic body size (b_M) or per unit of production (b_P). The error term (e) denotes the feed not used for either maintenance or producing products. If desired, production can be further subdivided into fat and lean gain in the case of a growing calf, or into lactation and fetal growth in the case of a lactating or pregnant cow, respectively.

Differences between animals in b_M , including those due to genetics or breeding value that can be changed via selection, have been demonstrated following selection in mice (Nielsen et al., 1997). And, the magnitude of these differences is relatively large enough to lead us to believe that re-

ductions of 15 to 20% in maintenance costs would be achievable through long-term selection. Less clear is whether there are differences between animals in b_P , and if so, what amount might be due to underlying genetic causes. Work by Eggert and Nielsen (2006), using selected lines of mice, did not reveal genetic differences in b_P .

Selection Practices in Swine and Broiler Production

The general characteristics of broiler and swine selection programs are fairly similar and quite different from beef cattle. In part because of intensive production, lower value of individual animals, and perhaps most importantly, much shorter generation intervals, broiler and swine selection programs are controlled and directed by a small number of companies. In broiler breeding, only a handful of multinational companies control the genetic improvement programs. Both broiler and swine breeding programs are centered around a limited number of nucleus flocks and herds, respectively, and multiple sets of contemporary groups occur each year to make for more cost-effective year-round utilization of data collection technology. Costs of implementation in breeding companies are balanced against predicted genetic change and its value recovery to make decisions on implementation of selection programs, including methods, traits, population sizes, and phenotypic and genomic data collected.

In beef cattle, the breeding pyramid structure is less fully defined, and thus, evaluation of costs of implementation and value recovery are not easy. The long generation interval sets a long horizon for recovery of costs in selection programs. With many breeders trying to contribute at a nucleus level, inefficiencies become evident, as compared with the industrial organization that exists in swine and broiler breeding companies.

Broiler breeding programs emphasize feed conversion as the most important trait for improvement. An

example of the importance placed on measurement and subsequent selection for decreasing feed needs is that by Aviagen (D. Emmerson, 2012, Aviagen Group, Huntsville, AL, personal communication), an international leader in broiler breeding. This company annually collects feed-intake data on more than 150,000 birds in individual pens for selection decisions, and in addition, collects another 50,000 feed records, measured on full-sib families, to assess response under commercial conditions. Adjusted feed conversion has been the trait of focus for selection, although feed-efficiency selection in egg-laying populations has used residual feed intake (**RFI**, feed intake adjusted for BW maintained and level of output) as the trait of focus. Because the commercial goal is to reduce feed required to grow birds to a constant, defined market weight, emphasis has long been placed on selecting faster-growing birds. Yet for a feed measurement program, a fixed age schedule is employed, and thus, adjustments for BW are required when using feed conversion as the primary selection trait (Emmerson, 1997).

Data collection of feed intake is for 7- to 14-d periods, starting in the last phase of production (~35 d of age) before market BW is attained. As noted above, in a broiler system, hatches throughout the year keep new birds in queue for use of feed measurement equipment. Aviagen also uses their own proprietary technology for individual bird measurement of feeding behavior (number of visits, length of visits, size of meals, and so on; Howie et al., 2011). Estimated genetic correlations between feeding behavioral traits and feed intake have not been large, thus behavioral indicators of feed intake have not been uncovered.

The swine industry has increasingly adopted a structure similar to the broiler industry. The largest 25 producers in the United States manage ~3 million sows, which represent just less than 50% of the sow inventory. The vast majority of genetics provided to this market originate from

only 3 to 4 suppliers. As in poultry, feed conversion is the dominant trait in the selection objective for terminal lines, and increasingly for maternal lines.

Swine breeding companies leverage unique selection objectives within specialized populations to create commercial sows (maternal lines) and market pigs (terminal lines). Historically, litter size has been the dominant trait in the selection objective of maternal lines. However, as litter size has increased, the incremental value of each additional pig decreases relative to the other traits in the selection objective. Recent changes in the cost of feed, and the performance level achieved for litter size, have increased the relative emphasis on feed conversion in maternal lines. Today, feed efficiency would account for slightly more than 50% of the selection objective in a terminal line and between 30 and 40% of the objective in a maternal line.

In general, feed conversion is measured only in the growing pig, specifically during the finishing period. The Danbred breeding program is a typical example that would be representative of most companies. In this program, feed intake is recorded during the period between 11 and 22 wk of age. Pigs are maintained in pens of 12 to 15 animals that are equipped with a feeding station designed to record feed intake. Each pig is equipped with an electronic ear tag that identifies the pig as it enters the feeding station. The number of visits to the feeder, the amount of time spent at the feeder, and the amount of feed consumed is recorded for each individual pig. These data are recorded on a sample of males that originate from the highest indexing litters resulting in the accuracy for the adjusted feed conversion (**FCR**) breeding value being the highest for those animals most likely to be selected to produce the next generation.

Adjusted feed conversion (adjusted to a fixed end BW) is used as the trait in the selection index. Feed conversion data from the boars are combined with information on growth

rate and body composition (percent lean) from the boars and all remaining males and females that undergo the same performance test less feed intake information. A breeding value for FCR is predicted on all animals in the population and weighted by its economic value in the overall index.

The use of feed intake measures in swine has, as in broilers, been used to improve the efficiency of the market pig, which represents the largest cost of producing pork. There is increasing interest in modifying these programs, particularly in maternal lines, to more directly (or perhaps reliably) affect the maintenance energy requirements of sows while maintaining progress in feed utilization by market pigs. It is particularly important that sows are able to maintain high levels of feed intake during the lactation period to maintain body condition, support high levels of milk production, and prepare for the next reproductive cycle. At the same time, the ideal sow has relatively low maintenance energy requirements during the gestation period. Selection for RFI is one method to address these needs in a more direct manner compared with the current emphasis on FCR during the finishing phase.

Improving Efficiency of Beef Production

As noted earlier in this paper, simply reducing feed intake is not the sole goal for a selection program. Care must be taken to not reduce production or output while we attempt to reduce feed intake for maintenance. Thus, a selection program with ranking criteria for selection and culling decisions that maximizes efficiency will include multiple traits that include both output(s) and input(s). A selection index that considers both cow or calf performance and post-weaning growth and carcass merit characteristics in the definition of net merit breeding value will be optimum.

The efficiency with which an animal utilizes feed can be expressed in different ways. For growing animals, traditional ratio measures are feed

efficiency (feed efficiency = G:F) and feed conversion ratio (FCR = F:G). Although feed efficiency and FCR are often used in production settings, these traits are problematic because they are the ratio of 2 traits (Gunsett, 1984). Koch et al. (1963) introduced the concept of using residuals for expression of efficiency. Residual feed intake (RFI) is defined in a beef growing animal as the amount of feed that the animal consumed adjusted for expected consumption based on requirements for maintenance and growth. Residual gain (RG) is defined as the amount of gain adjusted for feed intake. Thus, animals with negative RFI consume less than expected and are deemed more efficient, and animals with positive RG grow more rapidly than is expected and are thus deemed more efficient. Further elaboration on RG has been given by MacNeil et al. (2011) and Crowley et al. (2010). Arthur et al. (1996, 2001) and Crews (2005) further elaborate on RFI. For the purpose of improving production efficiency, it is recommended that RFI, RG, or both be computed using genetic regression coefficients based on estimates of genetic (co)variances (Kennedy et al., 1993), preferably from a single multiple-trait mixed model analysis. Both RFI and RG have been found to be moderately heritable.

From a genetic improvement perspective, it is important to recognize that selection for feed efficiency does not require an explicit measure of feed efficiency to be computed. Instead, selection for feed efficiency can be accomplished by selection on a linear index of traits that measure components of output (e.g., BW gain) and input (i.e., feed intake), with output traits receiving positive weights and input traits negative weights (MacNeil et al., 2013). Furthermore, Kennedy et al. (1993) showed the equivalence of selection indexes that incorporated intake or RFI, when the economic weights were calculated correctly.

One of the challenges, and a lingering question and need for further research, is to assess and understand the genetic correlation between feed

energy requirement for maintenance per unit size (the b_M coefficient described above) in a growing calf in a feedlot, which consumes a high proportion of grain or grain coproducts from ethanol production, versus in a reproducing cow, which consumes mostly forages in a range or pasture environment. Basarab et al. (2007) was unable to detect any significant antagonisms among feed intake and reproductive merit or lifetime productivity of dams that were the mothers of calves with different efficiency. Future results from a project nearing completion at the US Meat Animal Research Center with collection of cow feed intake, combined with growing calf data (Rolfe et al., 2011), will provide further insights. In addition, data from the USDA-supported National Program for Genetic Improvement of Feed Efficiency project (<http://www.beefefficiency.org>) will add clarification to this question.

Past work using breed differences as a method to infer possible genetic correlations has pointed to strong, positive genetic relationships between growing calf and reproducing cow energy requirements, per unit body size, for maintenance (b_M ; Montaña-Bermudez et al., 1990). In addition, Archer et al. (2002) found a strong and positive genetic correlation between RFI of growing calves in a feedlot and RFI of cows in a feedlot. The data on feed intake and utilization on heifers, before making replacement selection decisions, is lacking. Therefore, more study is needed to appropriately account for the associations of feed intake and other economically relevant traits in the female.

All of the measures of efficiency discussed above are favorably related to life-cycle production efficiency. Feed conversion ratio is used in many broiler and swine breeding company programs as the measure of feed efficiency. Measurement is done over a fairly well-defined BW and age interval within each company. Because new hatches or farrowings occur frequently, with new birds or animals ready for feed measurement at the starting BW and age, feed conversion

is measured in a narrowly defined window, and new batches enter and use the feed measurement equipment year-round. Statistical adjustments for BW range are still made to fine-tune the measurement, but the level of adjustment is relatively small. With the annual calving interval of beef cattle, many young cattle are ready for feed measurement at the same time, which results in expensive equipment then sitting idle for much of the year.

Especially for broiler production, but also for swine production, the magnitude of total feed consumption that is used by the reproducing female flock or herd is much smaller than what we observe with cattle. Low reproductive rate (<1 calf/cow per yr) and long time between successive reproductive cycles (1 yr) result in almost 65% of the feed energy in total-system or life-cycle being used by the reproducing cow herd, as opposed to the growing calf to slaughter. Although we cannot easily measure feed intake for grazing cows, improving the efficiency of feed utilization of cows is still of paramount importance. Thus, either protocols for measuring intake at grazing, or indicator traits indicative of intake at grazing, need to be identified to maximize improvement in production system efficiency.

Data Recording

Given the desirability of recording feed intake to enhance improvement in efficiency relative to that attainable from output traits alone, protocols for data recording take on increased importance. Archer et al. (1997) have demonstrated that a measurement period of 70 d will provide adequate precision in measurement of relevant performance traits in growing calves. The main limitation and need for a minimum of 70 d is precision in measurement of BW gain, rather than feed intake. At this point, protocols for data collection from reproducing females are less well established. However, historical studies of factors affecting life-cycle efficiency considered individually fed cows and calves for an entire production cycle (e.g.,

Davis et al., 1983; Kirkpatrick et al., 1985).

To make effective use of expensive feed measurement equipment, at least a couple of groups from the same calving season will have data collected in the system. These calves are likely to be more variable in age and BW than what happens in swine and broiler breeding programs. Opportunities to use central testing facilities, where animals from multiple herds can have feed intake data collected, can also reduce cost of collection of individual feed intake. However, variation in age and BW at the time of data collection will likely remain.

A National Cattle Evaluation Program

National cattle evaluation (NCE) seeks to provide producers with information regarding the genetic basis for economically relevant traits. To the extent that feed costs money, feed intake is an economically relevant trait. Alternative measures, derived from feed intake and performance, provide no additional information beyond that contained in the traits used in their calculation (Kennedy et al., 1993). Therefore, it is recommended that NCE analyze feed intake.

Conditions vary between different contemporary groups (BIF, 2010), potentially affecting not only the mean but also the variance of observations. Across testing facilities, different equipment may be employed, with ramifications for the observed variance in feed intake. Even in the same herd, the variance of feed intake may be altered because of environmental conditions and perhaps diets fed, the latter perhaps varying in both composition and form that may affect animal behavior. Finally, differences in sophistication in operating the test may result in feed intake being reported in different units (e.g., as fed, DM, or ME), which can be assumed to differ only by unobserved multiplicative constants. Furthermore, contemporary groups in which feed intake is recorded tend to be fairly large. Thus, standardization or normalization of

data, as proposed by MacNeil et al. (2011), has the desired result that measurement of feed intake within a contemporary group will have mean zero and standard deviation one.

Feed intake is known to be genetically and phenotypically correlated with other phenotypes that are more easily obtained, for instance post-weaning growth. Including these indicator trait phenotypes in NCE for feed intake has potential for increasing accuracy in the evaluation of feed intake and extending the evaluation of feed intake to many animals beyond those for which feed intake was observed.

Because of reliance on relatively expensive testing facilities, with limited capacity for collection of data on feed intake, only a selected sample of animals may be evaluated. Thus, to overcome selection bias, an NCE for feed intake should also contain correlated trait(s) recorded for all animals in the contemporary groups from which the evaluated animals were selected.

MacNeil et al. (2011) provide concepts and then an example for an NCE program in which measures of feed intake plus more easily measured indicator traits are incorporated to predict EPD for feed intake requirements. The genetic relationship matrix, in addition, greatly aids in prediction of breeding value for animals that have no measurement of feed intake on them.

Because of the cost of measurement, feed intake (hence, efficiency of feed utilization) will benefit from further development of genomic predictors to enhance prediction of breeding value in an NCE program. In a multibreed population of steers at the US Meat Animal Research Center, the best 96 SNP drawn from the BovineSNP50 Chip explained approximately three-fourths of the breeding value variance in genetic RFI (Snelling et al., 2011). Similarly, Rolf et al. (2012) found 55 to 65 SNP explained approximately 55% of the additive genetic variance of feed intake in growing Angus steers. A continuing requirement, for the use of genetic markers to enhance the evaluation of feed intake, is an

ongoing commitment to collection of phenotypic data for training marker prediction panels. Ideally, these data are collected on animals from different breeds and crossbreds to yield robust predictions across many genetic types.

Several opportunities exist to derive genetic predictors of merit and efficiency following the NCE program analysis of feed intake. These include EPD for RG, RFI, and residual intake and BW gain (Berry and Crowley, 2012), as well as selection indexes. If the feed intake data were standardized before the NCE program, then incorporation of the predicted genetic values into indexes or decision support systems may require back adjustment to a given diet formulation and environment, where the mean and standard deviation have estimated or assumed values.

The choice of which measure of feed intake or efficiency should be derived from the NCE program of feed intake and provided to breeders in terms of an EPD should be driven primarily by the goal to provide an EPD that promotes the proper use of the information provided by breeders in a multiple-trait setting. Thus, assuming that not all breeders use selection indexes and that many breeders are concerned about the effect of reducing feed intake capacity, it may be desirable to provide EPD for RG or RFI, rather than EPD for feed intake. Although there are compelling reasons for phenotypic measures of efficiency in other contexts, selection decisions in genetic improvement programs should be based on genetic predictions from the multiple-trait genetic evaluation of feed intake (MacNeil et al., 2013). The measure should also ensure that it addresses efficiencies both during the growing period and cow-calf phase of production.

IMPLICATIONS

Improvement of production-system efficiency is important to the profitability and sustainability of beef production. Substantial improvement results solely from increasing rate of production, by reducing *per diem*

costs associated with maintenance, as well as increasing reproduction by minimizing losses from feeding non-productive females. However, because there is variation between animals in utilization of feed energy, especially for maintenance, further improvement is possible through appropriate consideration of feed intake measurement in selection decisions. This consideration should be facilitated by an NCE program. Difficulty in measuring feed intake of grazing animals, especially reproducing females, may limit improvement of life-cycle efficiency. For the near term, measurement of feed intake will be centered in growing animals. Choosing which measure of feed intake or efficiency should be derived from the NCE program of feed intake and provided to breeders as EPD should be driven primarily by the goal to provide an EPD that promotes proper use of that information in a multiple-trait setting.

LITERATURE CITED

- Archer, J. A., P. F. Arthur, R. M. Herd, P. F. Parnell, and W. S. Pitchford. 1997. Optimum postweaning test for measurement of growth rate, feed intake, and feed efficiency in British breed cattle. *J. Anim. Sci.* 75:2024–2032.
- Archer, J. A., A. Reverter, R. M. Herd, D. J. Johnston, and P. F. Arther. 2002. Genetic variation in feed intake and efficiency of mature beef cows and relationships with postweaning measurements. 7th World Cong. Genet. Appl. Livest. Prod. CD-ROM Comm. 10–07.
- Arthur, P. F., J. A. Archer, D. J. Johnston, R. M. Herd, E. C. Richardson, and P. F. Parnell. 2001. Genetic and phenotypic variance and covariance components for feed intake, feed efficiency, and other postweaning traits in Angus cattle. *J. Anim. Sci.* 79:2805–2811.
- Arthur, P. F., R. M. Herd, J. Wright, G. Xu, K. Dibley, and E. C. Richardson. 1996. Net feed conversion efficiency and its relationship with other traits in beef cattle. *Proc. Aust. Soc. Anim. Prod.* 21:107–110.
- Basarab, J. A., D. McCartney, E. K. Okine, and V. S. Baron. 2007. Relationships between progeny residual feed intake and dam lifetime production efficiency traits. *Can. J. Anim. Sci.* 87:489–502.
- Berry, D. P., and J. J. Crowley. 2012. Residual intake and body weight gain: A new measure of efficiency in growing cattle. *J. Anim. Sci.* 90:109–115.
- BIF. 2010. Guidelines for Uniform Beef Improvement Programs. 9th ed. Beef Improvement Federation. www.beefimprovement.org.
- Crews, D. H., Jr. 2005. Genetics of efficient feed utilization and national cattle evaluation: A review. *Genet. Mol. Res.* 4:152–165.
- Crowley, J. J., M. McGee, D. A. Kenny, D. H. Crews Jr., R. D. Evans, and D. P. Berry. 2010. Phenotypic and genetic parameters for different measures of feed efficiency in different breeds of Irish performance-tested beef bulls. *J. Anim. Sci.* 88:885–894.
- Davis, M. E., J. J. Rutledge, L. V. Cundiff, and E. R. Hauser. 1983. Life-cycle efficiency of beef production. 1. Cow efficiency ratios for progeny weaned. *J. Anim. Sci.* 57:832–851.
- Dickerson, G. 1970. Efficiency of animal production—Molding the biological components. *J. Anim. Sci.* 30:849–859.
- Dickerson, G. E. 1978. Animal size and efficiency: Basic concepts. *Anim. Prod.* 27:367–379.
- Eggert, D. L., and M. K. Nielsen. 2006. Comparison of feed energy costs of maintenance, lean deposition, and fat deposition in three lines of mice selected for heat loss. *J. Anim. Sci.* 84:276–282.
- Emmerson, D. 1997. Commercial approaches to genetic selection for growth and feed conversion in domestic poultry. *Poult. Sci.* 76:1121–1125.
- Gunsett, F. C. 1984. Linear index selection to improve traits defined as ratios. *J. Anim. Sci.* 58:1185–1193.
- Howie, J. A., S. Avendano, B. J. Tolkamp, and I. Kyriazakis. 2011. Genetic parameters of feeding behavior traits and their relationship with live performance traits in modern broiler lines. *Poult. Sci.* 90:1197–1205.
- Kennedy, B. W., J. H. J. van der Werf, and T. H. E. Meuwissen. 1993. Genetic and statistical properties of residual feed intake. *J. Anim. Sci.* 71:3239–3250.
- Kirkpatrick, B. W., C. A. Dinkel, J. J. Rutledge, and E. R. Hauser. 1985. Prediction equations of beef cow efficiency. *J. Anim. Sci.* 60:964–969.
- Koch, R. M., L. A. Swiger, D. Chambers, and K. E. Gregory. 1963. Efficiency of feed use in beef cattle. *J. Anim. Sci.* 22:486–494.
- MacNeil, M. D., N. Lopez-Villalobos, and S. L. Northcutt. 2011. A prototype national cattle evaluation for feed intake and efficiency of Angus cattle. *J. Anim. Sci.* 89:3917–3923.
- MacNeil, M. D., M. M. Scholtz, and A. Maiwashe. 2013. Estimates of variance components for postweaning feed intake and growth in Bonsmara bulls and evaluation of alternative measures of feed efficiency. *S. Afr. J. Anim. Sci.* 43:18–24.
- Montaño-Bermudez, M., M. K. Nielsen, and G. H. Deutscher. 1990. Energy requirements for maintenance of crossbred beef cattle with different genetic potential for milk. *J. Anim. Sci.* 68:2279–2288.
- Nielsen, M. K., B. A. Freking, L. D. Jones, S. M. Nelson, T. L. Vorderstrasse, and B. A. Hussey. 1997. Divergent selection for heat loss in mice. II. Correlated responses in feed intake, body mass, body composition, and number born through fifteen generations. *J. Anim. Sci.* 75:1469–1476.
- Rolf, M. M., J. F. Taylor, R. D. Schnabel, S. D. McKay, M. C. McClure, S. L. Northcutt, M. S. Kerley, and R. L. Weaver. 2012. Genome-wide association analysis for feed efficiency in Angus cattle. *Anim. Genet.* 43:367–374.
- Rolfe, K. M., W. M. Snelling, M. K. Nielsen, H. C. Freetly, C. L. Ferrell, and T. G. Jenkins. 2011. Genetic and phenotypic parameter estimates for feed intake and other traits in growing beef cattle and opportunities for selection. *J. Anim. Sci.* 89:3452–3459.
- Snelling, W. M., M. A. Allan, J. W. Keele, L. A. Kuehn, R. M. Thallman, G. L. Bennett, C. L. Ferrell, T. G. Jenkins, H. C. Freetly, M. K. Nielsen, and K. M. Rolfe. 2011. Partial-genome evaluation of postweaning feed intake and efficiency of crossbred beef cattle. *J. Anim. Sci.* 89:1731–1741.