

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Historical Materials from University of  
Nebraska-Lincoln Extension

Extension

---

2001

## NF01-467 Turf Disease Fact Sheet No. 7: Management Program for Anthracnose

John E. Watkins

University of Nebraska--Lincoln, [jwatkins1@unl.edu](mailto:jwatkins1@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>



Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

---

Watkins, John E., "NF01-467 Turf Disease Fact Sheet No. 7: Management Program for Anthracnose" (2001). *Historical Materials from University of Nebraska-Lincoln Extension*. 906.  
<https://digitalcommons.unl.edu/extensionhist/906>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



# NebFact



Published by Cooperative Extension, Institute of Agriculture and Natural Resources,  
University of Nebraska-Lincoln

## Turf Disease Fact Sheet No. 7 Management Program for Anthracnose

by John E. Watkins, Extension Plant Pathologist

### Cause, Hosts and Occurrence

Anthracnose:            Cause: *Colletotrichum graminicola*  
                              Primary Hosts: Annual bluegrass, creeping bentgrass  
                              Occurrence: July - August

### Key Symptoms and Signs

- Irregularly shaped patches of yellow-bronze turf.
- Elongated reddish brown spots on the leaves.
- Black, spiny fungal structures on infected leaves, visible with 10X hand lens.

### Cultural/Maintenance Practices

- Provide sufficient nitrogen to maintain moderate growth rate through the summer.
- Irrigate to maintain plant vigor and avoid drought stress.

### Fungicide Program

- Preventive applications of reduced-rate tank mixes of a DMI fungicide and chlorothalonil at two-week intervals.

Products reported to provide fair to excellent control of anthracnose include:

#### Commercial Products

Active ingredient(s)	Product name(s)
----------------------	-----------------

azoxystrobin	Heritage
chlorothalonil	Daconil Ultrex, Thalonil, Manicure Ultrex
fenarimol	Rubigan
myclobutanil (DMI)	Eagle
propiconazole (DMI)	Banner MAXX
thiophanate-methyl	Cleary's 3336, Fungo, Cavalier
triadimefon (DMI)	Bayleton, Lesco Granular Turf Fungicide, Accost
trifloxystrobin (DMI)	Compass
chlorothalonil + thiophanate-methyl	Spectro 90

### **Home Lawn Products**

<i>Active ingredient(s)</i>	<i>Product name(s)</i>
propiconazole	ferti-lome Liquid Systemic Fungicide
thiophanate-methyl	Green Light Fung-Away II Systemic Fungicide, ferti-lome Halt Systemic, Dragon Systemic Fungicide 3336WP
triadimefon	Green Light Systemic Lawn Fungicide, Bonide Lawn Fungicide with Bayleton

Fungicides listed represent the best information available. No criticism is intended of products not listed, nor is endorsement by the University of Nebraska given to those listed. Read and follow all product label directions for mixing and application.

---

***File NF01-467 under PLANT DISEASES***

***F-8, Turf***

***Issued April 2001***

---

*Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Dean and Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.*

*University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.*