University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Harold W. Manter
Laboratory of Parasitology

Parasitology, Harold W. Manter Laboratory of

12-1972

Digenetic Trematodes of the Chesapeake Bay

Robin M. Overstreet

Gulf Coast Research Laboratory, robin.overstreet@usm.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/parasitologyfacpubs>

Part of the [Parasitology Commons](#)

Overstreet, Robin M., "Digenetic Trematodes of the Chesapeake Bay" (1972). *Faculty Publications from the Harold W. Manter Laboratory of Parasitology*. 903.

<http://digitalcommons.unl.edu/parasitologyfacpubs/903>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Harold W. Manter Laboratory of Parasitology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in *Chesapeake Science* 13 (December 1972), pp. S106–S107; doi: 10.2307/1350661
Copyright © 1972 Coastal and Estuarine Research Federation; published by Springer-Verlag. Used by permission.

Digenetic Trematodes of the Chesapeake Bay

Robin M. Overstreet

Gulf Coast Research Laboratory, Ocean Springs, Mississippi, USA

Present Status of Taxonomical Knowledge

It is safe to say that the taxonomy of Digenea from fishes and invertebrates from Chesapeake Bay is in need of considerable attention. First, the knowledge of what flukes are present is lacking, with the exception of a few scattered papers dealing with a limited number of species. A few such papers about adult Digenea from fishes are by Hopkins (1) and Anderson (2) and about larval forms from second intermediate hosts are by Stunkard and Uzman (3), Dillon (4), and Perkins (5). A list of the Digenea from Chesapeake Bay compiled by D. E. Zwerner and A. R. Lawler is most likely incomplete from the standpoint of actual species present. Second, there are probably many Digenea present that are identical to those found in fishes from North Carolina and Massachusetts. Much of the knowledge about Digenea from those areas is based on numerous reports by Edwin Linton dating from the late 1800s to 1940 (see Linton, (6), and (7) in particular). Many of the descriptions of the species are insufficient by modern standards. Some of the species are not valid, whereas others should be split into several species. Many of the problems created in the early literature by several authors have yet to be solved.

Present Status of Knowledge about the Distribution and Abundance of the Group

The viscera from several preserved fishes from Chesapeake Bay were given to me by Adrian R. Lawler to examine for parasites for a list of parasites from Chesapeake Bay. Most of the fish were infected with Digenea and most were new records for the parasite to be found in the particular host and in the Bay area. These findings suggest that Digenea are abundant and their distribution poorly known.

Present Status of Knowledge Concerning the Biology of the Group

The biology of several trematodes from near Woods Hole and from other areas along the Atlantic coast have been studied by R. M. Cable, H. W. Stunkard, W. E. Martin, and others. Some of the studied worms probably occur in the Bay. In general, however, the life histories and other aspects of the biology of marine and estuarine trematodes are poorly understood.

Present Status of Knowledge Concerning the Role of the Group in the Bay Ecosystem

Without a knowledge of what species are present, little can be said of the status of Digenea in the ecosystem. Digenea are seldom severely pathogenic to a definitive host, but they may have a profound effect on the first or second intermediate hosts, especially during specific seasons or in partially or entirely enclosed areas.

Present Status of Knowledge Concerning the Sensitivity of the Group to Man-Induced Environmental Changes

Digenea are especially valuable as indicators of environmental conditions. Molluscs are the first intermediate host for all known Digenea but two. In order for a trematode larva to infect a mollusc and asexually produce larvae capable of infecting a second intermediate host, the trematode must depend on both a receptive host and satisfactory environmental conditions. Proper conditions are also necessary for development in the second or additional intermediate hosts and in the definitive host. A breakdown in the availability of susceptible hosts or any of the favorable environmental conditions accompanying the various developmental and infective stages could eliminate the parasite. It should be pointed out that the presence of Digenea in an area is not usually detrimental to the involved hosts or the ecosystem.

References

1. Hopkins, S. H. 1954. The American species of trematode confused with *Bucephalus* (*Bucephalopsis*) *haimeanus*. *Parasitology* 44: 353–370.
2. Anderson, H. G. 1970. Annotated list of parasites of the bluefish *Pomatomus saltatrix*. US Bur. Sport Fish. Wild. Tech. Pap. 54, 15 p.
3. Stunkard, H. W., and J. R. Uzzmann. 1955. The killifish, *Fundulus heteroclitus*, second intermediate host of the trematode *Ascocotyle* (*Phagicola*) *diminuta*. *Biol. Bull.* 109(3): 475–483.
4. Dillon, W. A. 1966. Provisional list of parasites occurring on *Fundulus* spp. *Va. J. Sci.* 17(1):21–31.
5. Perkins, F. O. 1971. Sporulation in the trematode hyperparasite *Urosporidium crescens* De Turk, 1940 (Haplosporida: Haplosporidiidae)—an electron microscope study. *J. Parasit.* 57(1): 9–23.
6. Linton, E. 1905. Parasites of fishes of Beaufort, North Carolina. *US Fish Comm. Bull.* (1904) 24: 321–428.
7. Linton, E. 1940. Trematodes from fishes mainly from the Woods Hole region, Massachusetts. *U.S. Nat. Mus., Proc.* 88:1–172.