

9-16-2016

Sybra alternans (Wiedemann) (Lamiinae: Apomecynini): an Asian cerambycid established on Easter Island, Chile

José Mondaca

Servicio Agrícola y Ganadero (SAG), jose.mondaca@sag.gob.cl

Francisco Ramírez

Sociedad Chilena de Entomología, Santiago, ramcua@hotmail.com

Sergio Rothmann

Servicio Agrícola y Ganadero (SAG), sergio.rothmann@sag.gob.cl

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

Mondaca, José; Ramírez, Francisco; and Rothmann, Sergio, "*Sybra alternans* (Wiedemann) (Lamiinae: Apomecynini): an Asian cerambycid established on Easter Island, Chile" (2016). *Insecta Mundi*. 1009.

<http://digitalcommons.unl.edu/insectamundi/1009>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0503

Sybra alternans (Wiedemann) (Lamiinae: Apomecynini):
an Asian cerambycid established on Easter Island, Chile

José Mondaca
Servicio Agrícola y Ganadero (SAG)
Avenida Portales N° 3.396
Santiago, Chile

Francisco Ramírez
Sociedad Chilena de Entomología
Santiago, Chile

Sergio Rothmann
Unidad de Entomología
Departamento Laboratorios y Estaciones Cuarentenarias
Servicio Agrícola y Ganadero (SAG)
Santiago, Chile

Date of Issue: September 16, 2016

José Mondaca, Francisco Ramírez, and Sergio Rothmann
Sybra alternans (Wiedemann) (Lamiinae: Apomecynini): an Asian cerambycid
established on Easter Island, Chile
Insecta Mundi 0503: 1–6

ZooBank Registered: urn:lsid:zoobank.org:pub:13A5A379-D164-46A9-AD41-A9EF933BC7B6

Published in 2016 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com
Assistant Editor: David Plotkin, e-mail: insectamundi@gmail.com
Head Layout Editor: Eugenio H. Nearn
Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas
Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and Submission Requirements available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and AgriFood Canada, Ottawa, ON, Canada
The Natural History Museum, London, UK
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

Sybra alternans (Wiedemann) (Lamiinae: Apomecynini): an Asian cerambycid established on Easter Island, Chile

José Mondaca

Servicio Agrícola y Ganadero (SAG)
Avenida Portales N° 3.396, Santiago, Chile
jose.mondaca@sag.gob.cl

Francisco Ramírez

Sociedad Chilena de Entomología
Santiago, Chile
ramcua@hotmail.com

Sergio Rothmann

Unidad de Entomología, Departamento Laboratorios y Estaciones Cuarentenarias
Servicio Agrícola y Ganadero (SAG)
Santiago, Chile
sergio.rothmann@sag.gob.cl

Abstract. Seven specimens of *Sybra alternans* (Wiedemann) (Cerambycidae: Lamiinae: Apomecynini) were captured with a UV light trap and by beating branches of trees in the urban area of Easter Island (Chile) during 2011–2016, representing the first record of the species in this Chilean insular territory, but not continental Chile. *Sybra alternans* is native to Southeast Asia and has been introduced accidentally to Hawaii and Florida in the United States of America. Data are presented from the literature on this species' distribution, host plants, and biological information.

Key words. Alien species, longhorned beetles, new introductions.

Resumen. Siete ejemplares de *Sybra alternans* (Wiedemann) (Cerambycidae: Lamiinae: Apomecynini) fueron capturados con una trampa de luz UV y mediante el sacudido de ramas de árboles en el área urbana de la Isla de Pascua (Chile) durante los años 2011-2016, representando el primer registro de la especie en este territorio insular chileno, no estando presente hasta ahora en el continente. A pesar de que *S. alternans* es una especie nativa del sudeste asiático, se ha introducido accidentalmente en Hawaii y Florida en los Estados Unidos. De acuerdo a lo señalado en la literatura, se entrega información sobre su distribución, plantas hospedantes y aspectos biológicos de la especie.

Palabras clave. Escarabajo de cuernos largos, especies foráneas, nuevas introducciones.

Introduction

Of the 180 species of longhorned beetles (Coleoptera: Cerambycidae) known to occur in Chile (Elgueta 2000), eight are recognized as alien species (Mondaca and Zavala 2016). Four of these species have been detected or identified for the first time in the past decade (Barriga and Cepeda 2007; Mondaca 2008; SAG 2008; Mondaca and Zavala 2016), representing new introductions far from their native geographic ranges. The eight exotic cerambycids already reported as established in Chile were summarized by Mondaca and Zavala (2016) and include six species introduced and established in the mainland of Chile: *Nathrius brevipennis* (Mulsant) [native to Europe], *Phoracantha semipunctata* Fabricius and *P. recurva* Newman [both native to Australia], *Ambeodontus tristis* (Fabricius) [native to New Zealand and Tasmania], *Xylotrechus rusticus* (Linnaeus) [native to Europe], and recently *Psapharochrus jaspideus* (Germar) [native to South America but not including Chile] (González 1989; Sandoval 2002; Barriga and Cepeda 2007; SAG 2008; Mondaca and Zavala 2016); and two species restricted to Easter Island: *Ceresium unicolor* (Linnaeus) [native to Asia] and *Lagocheirus obsoletus* (Thomson) [native to Central America] (Cerdeña 1991; Mondaca 2008). All of these species were likely introduced to Chile as result of

international trade or tourism, and with the exception of the two *Phoracantha* species, the remaining six species do not represent major threats to forestry or natural ecosystems in Chile.

Collections made with a static UV light trap installed by staff of the local office of the Servicio Agrícola y Ganadero of Chile (SAG) in the residential area of Hanga Roa village, Easter Island, during 2011–2015, plus a collection made in 2016 by one of the authors (FR) while shaking dry branches of *Erythrina* sp. tree (Fabaceae, “ceibo”), provided seven adult cerambycid specimens that were later identified as *Sybra alternans* (Wiedemann, 1823) (Lamiinae: Apomecynini). To our knowledge, this is the first report of this species from insular Chile. The objective of this paper is to report details on the first collections of *S. alternans* on Easter Island, which makes it the third cerambycid to be introduced and established on this remote South Pacific island.

Sybra Pascoe is the most speciose genus within the tribe Apomecynini and it is widespread throughout the eastern Palearctic, Oriental, and Australian Regions (Slipinski and Escalona 2013). The genus comprises more than 420 species, and only one of them, *S. alternans*, is known to be found outside its natural distribution area, being successfully established in Hawaii and Florida in the United States of America (Gressitt 1956; Thomas 2000; Chen et al. 2001; Haack 2006).

Materials

The generic and specific identification was completed using the key of Rondon and Breuning (1970). Photographs of the adult beetle were taken using a digital camera Nikon D300s of 12 megapixels. The specimens collected in this study are deposited in the collection of the Unidad de Entomología, Subdepartamento de Laboratorios y Estaciones Cuarentenarias, Servicio Agrícola y Ganadero (CSAG) (Santiago, Chile), and the private collection of Francisco Ramírez (Santiago, Chile).

Sybra alternans (Wiedemann, 1823)

(Fig. 1)

Material Examined. Chile, Valparaíso Region: Easter Island [Rapa Nui], UV light trap, VI-2011, III-2014, 22-XII-2014, col. C. Valdés (3 females). Valparaíso Region, Rapa Nui, UV light trap, 1-X-2014, col. S. Ríos (1 female). Valparaíso Region, Rapa Nui, UV light trap, 23-III-2015, col. S. Ríos (2 females). Chile, Valparaíso Region, Easter Island, sector Hanga Kio’e, beating *Erythrina* sp., 20-IV-2016, leg. F. Ramírez (1 female).

Host Plants. *Sybra alternans* is a polyphagous species that infests numerous host plants. It has been detected attacking some fruit trees and crops such as *Ficus carica* L. (Moraceae, “fig tree”), *Ananas comosus* (L.) Merr. (Bromeliaceae, “pineapple”), *Musa paradisiaca* L. (Musaceae, “banana”), *Phaseolus vulgaris* L. (Fabaceae, “bean”), *Ocimum basilicum* L. (Lamiaceae, “basil”), *Gossypium hirsutum* L. (Malvaceae, “cotton”), *Saccharum officinarum* L. (Poaceae, “sugar cane”), and some ornamental plants typical of the South Pacific island. For a full list of host and respective citations, see Chen et al. (2001). In Florida, it was collected from dead limbs of *Ficus aurea* Nutt. (Moraceae, “golden fig”) (Thomas 2000), while on Easter Island an adult was captured by shaking the dry branches of *Erythrina* sp. trees (Fabaceae, “ceibo”).

Distribution. *Sybra alternans* is an Asian species distributed in Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Taiwan, Indonesia, Mariana Islands, Caroline Islands, Marshall Islands, and the Philippines (Rondon and Breuning 1970; Roguet 2016). It was introduced in the early twentieth century to Hawaii (see details in Chen et al. 2001) and more recently to Florida in the United States of America (Thomas 2000). It may have arrived in Florida via military flights from the Hawaiian Oahu military bases to those at Homestead, Florida (Samuelson and Howarth 2013). In Chile, *S. alternans* is known to occur only on to Easter Island, Valparaíso Region (**new record**) (Fig. 2).

Biology. The life cycle of *S. alternans* has been described briefly by Swezey (1950), who mentions that the life cycle can be completed in about four months. Adults are nocturnally active and seem to oviposit preferentially on host plants that are thoroughly dried. This cerambycid feeds on the inner fermenting and decaying bark and the outer sapwood. The larvae bore into the wood to pupate for their final transformation to adults.

Remarks. Repeated collections of *S. alternans* during six years in two different parts of Easter Island (Chile) provide strong evidence that a population of this cerambycid has successfully established on the island. Because the collection sites of *S. alternans* are located near the airport of Mataverí (see Fig. 2), it is likely that this species reached Easter Island via international commercial flights. Samuelson and Howarth (2013) proposed a similar hypothesis (but through military flights) to try to explain the introduction of this species from Hawaii to Florida.

Acknowledgments

We thank Marcelo Guerrero for the photograph illustrating the adult beetle, M. J. Paulsen (University of Nebraska State Museum) for improving the English, and Mario Elgueta (Museo Nacional de Historia Natural de Santiago, Chile) and Robert A. Haack (USDA Forest Service) for critical review of the paper.

Literature Cited

- Barriga, J. E., and D. E. Cepeda. 2007.** Nuevas sinonimias en Cerambycidae de Chile (Coleoptera). *Revista Chilena de Entomología* 33: 5–13.
- Cerda, M. 1991.** Presencia de *Ceresium unicolor unicolor* (Fabr.) en Isla de Pascua (Coleoptera: Cerambycidae). *Acta Entomológica Chilena* 16: 271–272.
- Chen, H., A. Ota, and G. E. Fonash. 2001.** Infestation of *Sybra alternans* (Cerambycidae: Coleoptera) in a Hawaii banana plantation. *Proceedings of the Hawaiian Entomological Society* 35: 119–122.
- Elgueta, M. 2000.** Coleoptera de Chile. p. 145–154. *In:* F. Martín-Piera, J. J. Morrone, and A. Melic (eds.). *Hacia un proyecto CYTED para el inventario y estimación de la diversidad entomológica Iberoamericana*, Monografías Tercer Milenio, Vol. 1. Sociedad Entomológica Aragonesa, Zaragoza, España.
- González, R. H. 1989.** Insectos y ácaros de importancia agrícola y cuarentenaria en Chile. Editorial Ograma. Santiago, Chile. 310 p.
- Gressitt, J. L. 1956.** Insects of Micronesia Coleoptera: Cerambycidae. B. P. Bishop Museum, *Insects of Micronesia* 17(2): 61–183.
- Haack, R. A. 2006.** Exotic bark- and wood-boring Coleoptera in the United States: recent establishments and interceptions. *Canadian Journal of Forest Research* 36: 269–288.
- Mondaca, J. 2008.** Primer registro de *Lagocheirus undatus* (Voet) (Coleoptera: Cerambycidae: Acanthocinini) en Isla de Pascua, Chile. *Revista Chilena de Entomología* 34: 63–68.
- Mondaca, J., and J. Zavala. 2016.** *Psapharochrus jaspideus* (Germar) (Coleoptera: Cerambycidae: Lamiinae: Acanthoderini), a new exotic longhorn beetle in Chile. *The Coleopterists Bulletin* 70(2): 391–394.
- Roguet, J.-P. 2016.** Lamiinae du Monde (Coleoptera Cerambycidae Lamiinae). (Available at ~ www.lamiinae.org . Last accessed 14 July 2016).
- Rondon, J. A., and S. Breuning. 1970.** Lamiinae du Laos. p. 315–571. *In:* J. L. Gressitt, J. A. Rondon, and S. Breuning (eds.). *Cerambycid-beetles of Laos (Longicornes du Laos)*. Pacific Insects Monograph 24. Bishop Museum, Hawaii. 651 p.
- SAG [Servicio Agrícola y Ganadero]. 2008.** *Rusticoclytus rusticus* (Linnaeus) (Coleoptera, Cerambycidae) en Chile. Ficha técnica N°5.
- Samuelson, G. A., and F. G. Howarth. 2013.** Notes on a recently established darkling beetle (Coleoptera: Tenebrionidae) in Hawaii. *Bishop Museum Occasional Papers* 114: 67–68.

- Sandoval, A. 2002.** Control biológico en Chile. p. 102–103. *In*: J. Macías, M. Arguedas, and L. Hilje (eds.). Plagas Forestales Neotropicales. Manejo Integrado de Plagas y Agroecología, 64. Costa Rica.
- Slipinski, A., and H. E. Escalona. 2013.** Australian Longhorn Beetles (Coleoptera: Cerambycidae). Vol. 1: Introduction and Subfamily Lamiinae. CSIRO Publishing, Collingwood, Australia. 484 p.
- Swezey, O. H. 1950.** Notes on the life cycle of certain introduced Cerambycid beetles. Proceedings of the Hawaiian Entomological Society 14(1): 187–188.
- Thomas, M. C. 2000.** *Ficus aurea* (Florida strangler fig, golden fig) – *Sybra alternans* (Wiedemann), a longhorned beetle. Tri-ology Technical Report 39(3). Florida Division of Plant Industry, Gainesville, FL. (Available at ~ <http://www.freshfromflorida.com/Divisions-Offices/Plant-Industry/Plant-Industry-Publications/Tri-ology-FDACS-DPI/Triology-Vol.-39-No.-3>. Last accessed July 2016.)

Received August 18, 2016; Accepted August 25, 2016.

Review editor Michael L. Ferro.



Figure 1. Female specimen of *Sybra alternans* (Wiedemann) collected on Easter Island (Chile), dorsal habitus. Scale bar = 2 mm.




Figure 2. Collection sites for *Sybra alternans* on Easter Island, Chile (black circles). Rapa Nui National Park is indicated in light grey.