

2017

Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of forages for Arctic caribou: effects of location, phenology and simulated digestion

Lindsay L. Vansomeren
University of Alaska, Fairbanks

Perry S. Barboza
Texas A&M University, psbarboza@tamu.edu

David D. Gustine
US Geological Survey

M. Syndonia Bret-Harte
University of Alaska, Fairbanks

Follow this and additional works at: <http://digitalcommons.unl.edu/usgsstaffpub>

 Part of the [Geology Commons](#), [Oceanography and Atmospheric Sciences and Meteorology Commons](#), [Other Earth Sciences Commons](#), and the [Other Environmental Sciences Commons](#)

Vansomeren, Lindsay L.; Barboza, Perry S.; Gustine, David D.; and Bret-Harte, M. Syndonia, "Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of forages for Arctic caribou: effects of location, phenology and simulated digestion" (2017). *USGS Staff -- Published Research*. 1002.
<http://digitalcommons.unl.edu/usgsstaffpub/1002>

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Rapid Commun. Mass Spectrom. 2017, 31, 813–820
(wileyonlinelibrary.com) DOI: 10.1002/rcm.7849

Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of forages for Arctic caribou: effects of location, phenology and simulated digestion

Lindsay L. Vansomeren¹, Perry S. Barboza^{2*},† , David D. Gustine^{3‡} and M. Syndonia Bret-Harte²

¹Department of Biology and Wildlife, University of Alaska Fairbanks, Fairbanks, AK 99775-7000, USA

²Institute of Arctic Biology, Department of Biology and Wildlife, University of Alaska Fairbanks, Fairbanks, AK 99775-7000, USA

³U.S. Geological Survey, Alaska Science Center, 4210 University Drive, Anchorage, AK 99508, USA

RATIONALE: The use of stable isotopes for dietary estimates of wildlife assumes that there are consistent differences in isotopic ratios among diet items, and that the differences in these ratios between the diet item and the animal tissues (i.e., fractionation) are predictable. However, variation in isotopic ratios and fractionation of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values among locations, seasons, and forages are poorly described for arctic herbivores especially migratory species such as caribou (*Rangifer tarandus*).

METHODS: We measured the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of seven species of forage growing along a 200-km transect through the range of the Central Arctic caribou herd on the North Slope of Alaska over 2 years. We compared forages available at the beginning (May; $n = 175$) and the end ($n = 157$) of the growing season (September). Purified enzymes were used to measure N digestibility and to assess isotopic fractionation in response to nutrient digestibility during simulated digestion.

RESULTS: Values for $\delta^{13}\text{C}$ declined by 1.38 ‰ with increasing latitude across the transect, and increased by 0.44 ‰ from the beginning to the end of the season. The range of values for $\delta^{15}\text{N}$ was greater than that for $\delta^{13}\text{C}$ (13.29 vs 5.60 ‰). Differences in values for $\delta^{13}\text{C}$ between graminoids (*Eriophorum* and *Carex* spp.) and shrubs (*Betula* and *Salix* spp.) were small but $\delta^{15}\text{N}$ values distinguished graminoids (1.87 ± 1.02 ‰) from shrubs (-2.87 ± 2.93 ‰) consistently across season and latitude. However, undigested residues of forages were enriched in ^{15}N when the digestibility of N was less than 0.67.

CONCLUSIONS: Although $\delta^{15}\text{N}$ values can distinguish plant groups in the diet of arctic herbivores, variation in the digestibility of dietary items may need to be considered in applying fractionation values for ^{15}N to caribou and other herbivores that select highly digestible items (e.g. forbs) as well as heavily defended plants (e.g. woody browse). Published in 2017. This article is a U.S. Government work and is in the public domain in the USA.

The stable isotopes of C and N have been used to measure the diet of wild herbivores,^[1] but their use for fine-scale dietary analysis in the Arctic has been problematic. The proportions of diet composed of each forage species are calculated using mixing models, and these estimates are more precise and accurate when forage species have large and distinct differences in stable isotope ratios. In fact, most dietary analyses for herbivores rely on assigning their diets to C3 woody browse and C4 grass categories because of the large differences in $\delta^{13}\text{C}$ values between these groups. In northern ecosystems, however, C4 plants are absent,^[2] leaving only small differences in $\delta^{13}\text{C}$ values among the C3 plant species to resolve the contributions of different plant groups to the

diet. Nevertheless, both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values have been used successfully for broad-scale dietary analysis in northern herbivores,^[3,4] although less frequently than in temperate and tropical systems.^[1,5,6]

Mixing models also require adjusting diet components for fractionation (i.e. diet discrimination factors^[7]), which is the difference in isotopic ratio between the dietary item and the tissue of the animal due to digestion and metabolism.^[8] Most studies use constant values for fractionation (e.g., +3‰ for $\delta^{15}\text{N}$ values) between forage plants and herbivores, but recent work has suggested that diet-tissue fractionation values may be influenced by the quality of forage, because differing fractions of ^{13}C and ^{15}N are presented to the herbivore according to the digestibility^[9] and/or protein quality^[10] of food items. Accurate diet-feces fractionation values of ^{15}N are difficult to obtain due to contamination of ^{15}N -enriched microbial and endogenous compounds in feces.^[11] Adjusting diet sources with correct fractionation values will improve the accuracy and precision of diet estimates especially in arctic ecosystems where diet resolution may be limited by small isotopic differences between forage plant classes.

Our objectives in this study were to examine seasonal, regional, and species-specific differences in the stable isotope ratios of C and N in arctic forages to determine which of these

* Correspondence to: P. S. Barboza, Department of Wildlife and Fisheries Sciences, Texas A&M University, 210 Nagle Hall, 2258 TAMU, College Station, TX 77843, USA.
E-mail: psbarboza@tamu.edu

† Present address: Department of Wildlife and Fisheries Sciences, Texas A&M University, 210 Nagle Hall, 2258 TAMU, College Station, TX 77843, USA.

‡ Present address: Grand Teton National Park, Moose, WY 83012, USA.

elements would be more useful for estimating the diet of caribou. The stable isotope ratios of C and N in forage plants were measured before and after simulated digestion to determine whether the indigestible residue sampled from the feces or the assimilated fraction sampled from the tissues were biased in their isotopic ratios. We used total phenols as an index of plant secondary metabolites to examine the effect of these anti-nutrients on any fractionation during simulated digestion.

EXPERIMENTAL

Study area and sampling design

This study was conducted in the summer range of the Central Arctic caribou herd on the North Slope of Alaska^[12,13] from 2011 to 2012 (Fig. 1). We sampled nine sites spread evenly along the Dalton Highway from the Kuparuk River to Prudhoe Bay (Fig. 1). Sites were classified into three ecoregions: Brooks Range, Arctic Foothills, and Coastal Plain.^[14]

Samples of six preferred forage species (*Carex aquatilis*, *C. bigelowii*, *Eriophorum vaginatum*, *Pedicularis* spp., *Salix pulchra*, and *S. richardsonii*)^[15] were collected, when present, every 2 weeks from late May to late September. In addition, we collected samples of *Betula nana* in 2012, because, although this species does not make up a large part of North Slope caribou diets at present,^[16] it may be eaten by caribou more frequently in the future because it is increasing in abundance throughout the Arctic.^[17–19] Indeed, another shrub birch, *B. glandulosa*, makes up a significant part of caribou diets in Quebec.^[20] Forage plants were sampled to mimic caribou browsing and grazing – i.e., for woody browse, easily accessible leaves and twigs were stripped off, while forbs and graminoids were clipped at ground level.

Forage samples were transferred to paper bags and air-dried at ambient temperature (0–22°C) in the field, then air-dried to constant mass in a forced-air oven at 50–55°C when the samples were returned to the laboratory, within 2–6 days of collection. A small subsample (approximately 70 g) of woody browse was immediately frozen in the field




Figure 1. Location of study sites within the range of the Central Arctic caribou herd. Sites were located in three ecoregions (Coastal Plain, Arctic Foothills, and Brooks Range) along the Dalton Highway. The distribution of the Arctic ecoregions in Alaska is noted in the inset. [Color figure can be viewed at wileyonlinelibrary.com]

and freeze-dried upon return to the laboratory (model 7755044; Labconco, Kansas City, MO, USA) to test for the presence of plant secondary metabolites (PSM). Dried samples were ground through a #20 mesh (1.27 mm) in a Wiley mill (Thomas Scientific, Swedesboro, NJ, USA) or a centrifugal mill (ZM 200; Retsch, Haan, Germany).

Laboratory analyses

We measured the total N content of 771 forage samples with an elemental analyzer (CNS2000; LECO, St. Joseph, MI, USA) to select two subsets of samples for the start and end of the phenological curve for each forage at each site and year.^[21] The early-season subset ($n = 175$) were samples with the highest N content whereas the last sample collected at each location comprised the late-season subset ($n = 157$). Samples from the early- and late-season subsets were analyzed for digestibility of C and N (g digested/g whole) by analyzing the nutrient content of undigested residues after *in vitro* digestion with purified enzymes by a method that had previously been validated for caribou.^[22]

The $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of forage samples (‰) from the early- and late-season subsets in 2011 (all species) and in 2012 (woody browse species only) were determined with a Europa Scientific 20-20 continuous flow isotope ratio mass spectrometer (Sercon Ltd, Crewe, UK) at the Alaska Stable Isotope Facility, University of Alaska, Fairbanks (Fairbanks, AK, USA). In addition, residues from the *in vitro* digestibility method were analyzed for their $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values after extraction with hot water.^[11] We used peptone (P7750; Sigma-Aldrich, Milwaukee, WI, USA) as a reference standard to monitor machine drift every 10 assays within each analytical run. Fractionation between the diet-indigestible fraction ($F_{\text{diet-indigestible}}$) for the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values was calculated as:

$$F_{\text{diet-indigestible}} = \delta X_{\text{residue}} - \delta X_{\text{whole plant}} \quad (1)$$

Fractionation of the diet-digested fraction for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values was calculated by converting each isotope measurement into mass ratios (g isotope/g element) and then multiplying this value by the N and C content (g element/g dry matter), respectively, of whole plants and indigestible residues to obtain the content of ^{15}N , ^{14}N , ^{13}C , and ^{12}C on a dry matter basis.^[23] Differences in the content of each isotope between indigestible residues and whole plants were used to calculate the mass ratios and $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of digested fractions, which were then converted into delta notation.

Statistical analyses

All analyses were conducted in Stata 12.0 (StataCorp, College Station, TX, USA). Forage species were grouped to facilitate comparisons among plant functional groups: graminoid (*C. aquatilis*, *C. bigelowii*, and *E. vaginatum*), woody browse (*B. nana*, *S. pulchra*, and *S. richardsonii*), and forb (*Pedicularis* sp.). Groups of data are summarized as mean \pm standard deviation (SD) where indicated. We used $P < 0.05$ as the criterion for significance of α in all comparisons. Bonferroni corrections were applied to determine the significance of multiple post-hoc comparisons.

We used analysis of variance (ANOVA) models to examine spatial, temporal, and species-specific changes in values for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, with species, ecoregion, and subset (early season or late season) as fixed factors. We tested values of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ for interactions of species \times subset, species \times ecoregion, and ecoregion \times subset. Values of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ were log transformed using the Inskew procedure in Stata^[24] to meet assumptions for normality and tested using the Shapiro-Wilk procedure. The significance of each parameter was assessed with Wald tests.

Linear ordinary least-squares regressions were used to examine the effect of N and C digestibility on isotopic fractionation. Paired *t*-tests were used to determine the significance of fractionation between diet, indigestible residues, and digested fractions for each forage species.

RESULTS

Reference assays for N were not significantly different among analytical runs on the mass spectrometer ($n = 5$; $F_{4,30} = 0.16$) and were not significantly different from the expected $\delta^{15}\text{N}$ value of 7.00‰ ($n = 35$; 7.03 ± 0.28 ; $t_{34} = 0.72$). Similarly, reference assays for C were not significantly different among mass spectrometer runs ($n = 5$; $F_{4,30} = 0.68$) and were not significantly different from the expected $\delta^{13}\text{C}$ value of -15.80 ‰ ($n = 35$; -15.77 ± 0.23 ; $t_{34} = 0.78$).

Spatiotemporal variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values

An ANOVA model that included plant group and ecoregion without season provided the greatest inference for explaining variation in $\delta^{15}\text{N}$ values of forages. Plant groups differed in $\delta^{15}\text{N}$ values (range = -9.47 to $+3.82$ ‰; $F_{2,97} = 140.56$; $P < 0.01$; Table 1; Fig. 2), because graminoids were enriched in ^{15}N (1.87 ± 1.02 ‰), whereas woody browse (-4.31 ± 2.06 ‰) and *Pedicularis* spp. (-2.87 ± 2.93 ‰) were depleted in ^{15}N . Values of $\delta^{15}\text{N}$ also varied among species of woody browse (Kruskal-Wallis test; $P < 0.01$): *B. nana* was most depleted in ^{15}N (-7.38 ± 1.60 ‰), *S. richardsonii* was intermediate in $\delta^{15}\text{N}$ values (-4.54 ± 1.15 ‰), and *S. pulchra* was most enriched in ^{15}N (-2.87 ± 0.95 ‰). Although ecoregion contributed to the variation in $\delta^{15}\text{N}$ values, those values were not significantly different among ecoregions ($F_{2,97} = 2.65$; $P > 0.05$; Table 1; Fig. 3). Values of $\delta^{15}\text{N}$ were also not significantly different between early and late season although the total N concentration was significantly greater in early- than late-season forages ($F_{1,320} = 608.11$; $P > 0.001$): graminoids (2.66 ± 0.43 vs 1.32 ± 0.57 gN \cdot 100 g $^{-1}$ DM); woody browse (3.12 ± 0.56 vs 1.34 ± 0.35 gN \cdot 100 g $^{-1}$ DM); *Pedicularis* spp. (2.91 ± 0.66 vs 1.71 ± 0.82 gN \cdot 100 g $^{-1}$ DM).

The range of values across plant groups was smaller for $\delta^{13}\text{C}$ than for $\delta^{15}\text{N}$ (5.60 vs 13.29‰, respectively) but variation among species within plant groups was greater for $\delta^{13}\text{C}$ values than for $\delta^{15}\text{N}$ values. Plant groups differed in $\delta^{13}\text{C}$ values ($F_{2,96} = 23.33$; $P < 0.01$; Fig. 2): values for $\delta^{13}\text{C}$ were highest for graminoids (-26.57 ± 1.01 ‰), intermediate in woody browse (-27.61 ± 1.25 ‰), and lowest in *Pedicularis* spp. (-28.27 ± 1.02 ‰). Values of $\delta^{13}\text{C}$ varied among graminoids (Kruskal-Wallis test; $P < 0.01$), as *C. aquatilis* was slightly more depleted in ^{13}C (-27.39 ± 1.16 ‰) than *C. bigelowii*

Table 1. Stable isotope ratios in the summer range of the Central Arctic caribou herd, 2011–2013. *Betula nana* and *Carex bigelowii* were not found at sampling sites on the Coastal Plain

| | | $\delta^{15}\text{N}$ values | | | | | |
|-------------|-----------------------------|------------------------------|---|------------------|----|------------------|---|
| | | Ecoregion | | | | | |
| | | Brooks Range | | Arctic Foothills | | Coastal Plain | |
| Plant group | Forage species | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD | N |
| Browse | <i>Betula nana</i> | -5.89 \pm 0.74 | 4 | -8.37 \pm 1.14 | 6 | | |
| Graminoid | <i>Carex aquatilis</i> | 2.04 \pm 0.60 | 2 | 1.91 \pm 1.37 | 7 | 0.74 \pm 0.84 | 6 |
| Graminoid | <i>Carex bigelowii</i> | 2.38 \pm 0.68 | 4 | 1.32 \pm 0.45 | 6 | | |
| Graminoid | <i>Eriophorum vaginatum</i> | 2.44 \pm 0.31 | 4 | 2.06 \pm 0.76 | 8 | 2.73 \pm 1.23 | 4 |
| Forb | <i>Pedicularis</i> spp. | -3.62 \pm 0.54 | 4 | -3.39 \pm 0.77 | 6 | -2.63 \pm 1.30 | 4 |
| Browse | <i>Salix pulchra</i> | -2.20 \pm 0.54 | 8 | -3.39 \pm 0.77 | 12 | -2.63 \pm 1.30 | 4 |
| Browse | <i>Salix richardsonii</i> | -2.24 \pm 0.82 | 2 | -5.45 \pm 0.57 | 6 | -4.44 \pm 0.54 | 8 |

| | | $\delta^{13}\text{C}$ values | | | | | |
|-------------|-----------------------------|------------------------------|---|-------------------|----|-------------------|---|
| | | Ecoregion | | | | | |
| | | Brooks Range | | Arctic Foothills | | Coastal Plain | |
| Plant group | Forage species | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD | N |
| Browse | <i>Betula nana</i> | -28.42 \pm 0.59 | 4 | -28.88 \pm 0.56 | 6 | | |
| Graminoid | <i>Carex aquatilis</i> | -27.73 \pm 0.11 | 2 | -26.92 \pm 0.68 | 7 | -27.82 \pm 1.62 | 6 |
| Graminoid | <i>Carex bigelowii</i> | -25.58 \pm 0.53 | 4 | -26.22 \pm 0.44 | 6 | | |
| Graminoid | <i>Eriophorum vaginatum</i> | -25.58 \pm 0.53 | 4 | -26.22 \pm 0.44 | 6 | -26.67 \pm 0.53 | 4 |
| Forb | <i>Pedicularis</i> spp. | -28.04 \pm 1.30 | 4 | -28.12 \pm 1.10 | 6 | -28.74 \pm 0.61 | 4 |
| Browse | <i>Salix pulchra</i> | -26.19 \pm 0.70 | 8 | -26.85 \pm 0.91 | 12 | -28.73 \pm 1.37 | 4 |
| Browse | <i>Salix richardsonii</i> | -26.36 \pm 0.91 | 2 | -27.65 \pm 0.63 | 6 | -28.56 \pm 0.62 | 8 |

(-25.97 \pm 0.56 ‰) and *E. vaginatum* (-26.18 \pm 0.43 ‰). Values of $\delta^{13}\text{C}$ also varied among deciduous shrub species (Kruskal-Wallis test; $P < 0.01$): *B. nana* was the most depleted in ^{13}C ($\delta^{13}\text{C}$ -28.69 \pm 0.59 ‰), *S. richardsonii* was intermediate ($\delta^{13}\text{C}$ -27.94 \pm 0.97 ‰), and *S. pulchra* was least depleted in ^{13}C ($\delta^{13}\text{C}$ -26.94 \pm 1.25 ‰). Values of $\delta^{13}\text{C}$ also differed by ecoregion ($F_{2,96} = 15.71$; $P < 0.01$) and season ($F_{1,96} = 23.97$; $P < 0.01$): $\delta^{13}\text{C}$ values declined progressively from the Brooks Range and Arctic Foothills (-26.77 \pm 1.24 ‰ and -27.14 \pm 1.14 ‰, respectively) to the Coastal Plain (-28.15 \pm 1.21 ‰; Table 1; Fig. 3) and from early (-26.85 \pm 1.11 ‰) to late season (-27.29 \pm 1.30 ‰).

Effect of nutrient availability on caribou $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values

Digestibility of N was also greater in early than late season ($F_{1,303} = 115.23$; $P < 0.001$) for graminoids (0.59 \pm 0.10 vs 0.43 \pm 0.12), woody browse (0.42 \pm 0.16 vs 0.27 \pm 0.13) and *Pedicularis* spp. (0.80 \pm 0.07 to 0.68 \pm 0.09). Across plant groups, the fractionation of $\delta^{15}\text{N}$ values between the whole plant and the indigestible fraction was negatively correlated with N digestibility ($F_{1,77} = 10.59$; $P < 0.01$; $R^2 = 0.15$; Fig. 4), with no fractionation when the N digestibility was 0.67. As the N digestibility declined, the residues became significantly more enriched in ^{15}N than the whole plant especially among species

of woody browse (*B. nana*: $\delta^{15}\text{N}$ 2.12 \pm 1.21 ‰ and *S. richardsonii*: $\delta^{15}\text{N}$ 0.75 \pm 0.44 ‰; both $P \leq 0.01$). This increase in diet-indigestible fractionation with decreasing N digestibility may have been caused by interference from phenolic compounds, because fractionation of $\delta^{15}\text{N}$ values between the diet and the indigestible fraction was correlated with the concentration of phenolic compounds in *B. nana* ($Y = 0.13x - 0.04$; $R^2 = 0.62$; $P < 0.01$) and *S. pulchra* ($Y = 0.06x - 0.16$; $R^2 = 0.55$; $P < 0.02$). In comparison with the diet, digested fractions (available for assimilation into animal tissues) were depleted in ^{15}N for *B. nana* ($\delta^{15}\text{N}$ -10.48 \pm 12.98 ‰; $P = 0.03$), *E. vaginatum* ($\delta^{15}\text{N}$ -0.73 \pm 1.31 ‰; $P = 0.04$), and *S. richardsonii* ($\delta^{15}\text{N}$ -0.82 \pm 0.56 ‰; $P = 0.03$; Fig. 2).

The digestibility of C was greater in early than late season ($F_{1,199} = 52.50$; $P < 0.001$) for graminoids (0.34 \pm 0.06 to 0.24 \pm 0.10), woody browse (0.48 \pm 0.05 to 0.40 \pm 0.12) and *Pedicularis* spp. (0.71 \pm 0.08 to 0.68 \pm 0.11). Fractionation of the $\delta^{13}\text{C}$ values between the whole plant and the digestible and indigestible fractions was also related to forage plant quality. Across plant groups, fractionation of $\delta^{13}\text{C}$ values between the diet and the indigestible fraction was negatively correlated with C digestibility ($F_{1,75} = 40.14$; $R^2 = 0.30$; $P < 0.01$; Fig. 4), with no fractionation of the $\delta^{13}\text{C}$ values at a C digestibility of 0.30. As C digestibility increased, residues were more depleted in ^{13}C than the whole plant




Figure 2. Values for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in whole plants (black dots), indigestible residues (gray dots), and the digested fraction (hollow dots) of caribou forages collected from the summer range of the Central Arctic caribou herd, 2011–2012. Lines and whiskers represent ± 1 SD from the mean isotopic values of the whole plant. Symbols (*) next to gray dots indicate significant differences between the indigestible residue and the whole plant. Symbols (#) next to hollow dots indicate significant differences between the digested fraction and the whole plant.

(*B. nana*: $\delta^{13}\text{C}$ -0.42 ± 0.28 ‰; *Pedicularis* spp.: $\delta^{13}\text{C}$ -1.59 ± 0.46 ‰; both $P < 0.01$). Consequently, the fraction available for assimilation into the animal was more enriched in ^{13}C for *B. nana* ($\delta^{13}\text{C}$ 0.62 ± 0.38 ‰; $t_9 = -5.14$; $P < 0.01$) and *Pedicularis* spp. ($\delta^{13}\text{C}$ 0.78 ± 0.35 ‰; $t_{13} = -8.21$; $P < 0.01$; Fig. 2).




Figure 3. Values for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ of three plant groups (graminoids, woody browse and the forb *Pedicularis* spp.) across three ecoregions (Brooks Range, Arctic Foothills and Coastal Plain) from the summer range of the Central Arctic caribou herd, 2011–2012. Predicted means with ± 1 SE from ANOVA models with ecoregion and plant group as factors.

DISCUSSION

Although both C and N isotopes can be used to discriminate between monocot (graminoids) and dicot (woody browse and *Pedicularis* spp.) plants, several factors make $\delta^{15}\text{N}$ values a more reliable indicator of diet than $\delta^{13}\text{C}$ values for arctic herbivores. The $\delta^{15}\text{N}$ values had a 42% greater range than the $\delta^{13}\text{C}$ values for monocot and dicot forage plants, which was consistent with other studies of arctic and alpine plants.^[4,25–27] The values of $\delta^{15}\text{N}$ were less affected by temporal variation than those of $\delta^{13}\text{C}$, which declined over




Figure 4. Relationships between fractionation ($F_{\text{diet-indigestible}}$; the isotopic distance from the indigestible residue to the whole plant) and digestibility for both N (top panel) and C (bottom panel) in forages from the summer range of the Central Arctic caribou herd, 2011–2012. Straight lines indicate the linear regressions with all plant groups combined; thin curved lines are the 95% confidence intervals of the regression.

the season from the Brooks Range to the Coastal Plain, probably due to differing levels of water stress^[28–31] and associated effects of seasonal plant growth on stomatal exchanges of gases.^[32] Variation in values for $\delta^{15}\text{N}$ among species of deciduous shrub probably reflected differences in mycorrhizal associations.^[33]

Although differences in growing conditions can create useful distinctions between forage groups, it is still important to consider how those signatures may change during digestion because digestible and indigestible

fractions can display differing stable isotope ratios that would then be incorporated into animal tissues and feces. In particular, we found that fractionation varied according to the nutrient digestibility of forage plants. The digestibilities of both C and N showed a negative relationship with fractionation, but significant fractionation between diet and indigestible residues was only observed when the digestibilities were less than 0.67 for N and greater than 0.30 for C. Forage plants of differing digestibility may also be represented in animal tissues at different rates and in different proportions, even if they are consumed at the same rate. For example, Codron *et al.*^[9] found that the incorporation rates of ^{13}C were fastest when animals consumed highly digestible diets, and that isotopic compositions of various herbivore tissues were skewed according to the digestibility of the diet. Forbs such as *Pedicularis* spp. are avidly sought by caribou probably because their digestibility of C and N is high (Fig. 4). The C available for absorption from this forb would be enriched in ^{13}C compared with the whole plant and thus its $\delta^{13}\text{C}$ values would appear more similar to those obtained from digestion of graminoids. Fractionation of highly digestible forbs may therefore enhance estimates of the contribution of graminoids to the diet when considering the $\delta^{13}\text{C}$ values of tissues such as hair.^[4] Fractionation of ^{15}N during digestion is more likely to affect estimates of diet from indigestible residues especially when the digestibility of N is low (Fig. 3). Estimates of the diet from fecal $\delta^{15}\text{N}$ values may be skewed towards graminoids by enrichment of ^{15}N in indigestible residues of woody browse species (Fig. 2). These differences in fractionation among plant groups have the potential to bias estimates of consumption when using isotopic analyses of tissues and feces to reconstruct diets over large scales of space or time.^[34,35]

Isotopic fractionation is affected by nutrient digestibility through a variety of mechanisms. Waxes on the surface of leaves and lignin in the plant cell wall matrix are resistant to digestion by acid and enzymes.^[36] The lower $\delta^{13}\text{C}$ values in indigestible residues of forages are consistent with the $\delta^{13}\text{C}$ values of lipids and lignin that are typically lower than those of whole leaves.^[32] Digestion of N is affected by physical access of enzymes to substrates, inhibition of the enzyme, and the affinity of the enzyme for the substrate, all of which can influence N fractionation.^[36] Enzyme affinity for substrate proteins probably has little effect on fractionation because protease activities are high for a wide variety of dietary proteins and because most of the plant protein is present in the form of a single photosynthetic protein, Rubisco.^[37] Physical access and enzyme inhibition probably account for most of the fractionation due to the actions of PSMs such as tannins that can both limit physical access of dietary enzymes to protein and inhibit the dietary enzymes themselves^[38,39] depending on the binding affinity between tannins and proteins in the digestive tract.^[40] Fractionation of N in woody browse may therefore depend upon the suite of PSMs, which changes according to species, ecoregion, and season. Enrichment of ^{15}N in indigestible residues of woody browse would enhance the $\delta^{15}\text{N}$ values of fecal residues and the effect of microbial colonization of fibrous residues on fecal $\delta^{15}\text{N}$ values.^[11]

CONCLUSIONS

Fractionation between diet and indigestible residues reduces isotopic separation between browse and graminoids (Fig. 2) and thus decreases the estimated contribution of heavily defended shrubs such as *B. nana* to the diet of arctic caribou and other arctic ungulates. Mixing models that estimate diet from the isotopic ratios of herbivore tissues would be improved by including the range of fractionation values when the diet includes highly digestible items (e.g. forbs) or forages rich in PSM (e.g. woody browse).

Acknowledgements

We thank S. Aguilar, K. Iles, J. Lawlor, K. Oster, G. Roffler, R. Ruffner, B. Streever, E. Wald, R. Wilson, J. Welch, and N. Wolf for assistance in the laboratory, logistics, and with forage collections in the field. T. Howe, L. Oliver, and R. Shively assisted with laboratory analyses. This study was funded in part by the Department of Biology and Wildlife (B&W) and the Institute of Arctic Biology at University of Alaska Fairbanks. This work was part of the U.S. Geological Survey's (USGS) Changing Arctic Ecosystem Initiative and was supported by funding from the Wildlife Program of the USGS Ecosystem Mission Area. Support for LVS was provided by the B&W Department, the Calvin J. Lensink Graduate Fellowship, and the USGS through the Alaska Cooperative Fish and Wildlife Research Unit. Use of any trade names in this manuscript does not imply endorsement by the United States government.

REFERENCES

- [1] D. Codron, J. Codron. Reliability of delta C-13 and delta N-15 in faeces for reconstructing savanna herbivore diet. *Mammal Biol.* **2009**, *74*, 36.
- [2] H. Öpik, S. Rolfe. *The Physiology of Flowering Plants*. Cambridge University Press, Cambridge, UK, **2005**.
- [3] G. L. Finstad, K. Kielland. Landscape variation in the diet and productivity of reindeer in Alaska based on stable isotope analyses. *Arctic, Antarctic, Alpine Res.* **2011**, *43*, 543.
- [4] D. K. Kristensen, E. Kristensen, M. C. Forchhammer, A. Michelsen, N. M. Schmidt. Arctic herbivore diet can be inferred from stable carbon and nitrogen isotopes in C3 plants, faeces, and wool. *Can. J. Zool.* **2011**, *89*, 892.
- [5] M. H. O. M. Wittmer, K. Auerswald, P. Schönbach, R. Schäufele, K. Müller, H. Yang, Y. F. Bai, A. Susenbeth, F. Taube, H. Schnyder. Do grazer hair and faeces reflect the carbon isotope composition of semi-arid C3/C4 grassland? *Basic Appl. Ecol.* **2010**, *11*, 83.
- [6] D. Codron, J. Hull, J. S. Brink, J. Codron, D. Ward, M. Clauss. Effect of competition on niche dynamics of syntopic grazing ungulates: contrasting the predictions of habitat selection models using stable isotope analysis. *Evolut. Ecol. Res.* **2011**, *13*, 217.
- [7] C. A. Dennis, M. A. Macneil, J. Y. Rosati, T. E. Pitcher, A. T. Fisk. Diet discrimination factors are inversely related to delta15 N and delta13C values of food for fish under controlled conditions. *Rapid Commun. Mass Spectrom.* **2010**, *24*, 3515.
- [8] B. Fry. *Stable Isotope Ecology*. Springer, New York, **2006**.
- [9] D. Codron, J. Codron, M. Sponheimer, S. M. Bernasconi, M. Clauss. When animals are not quite what they eat: diet digestibility influences ^{13}C -incorporation rates and apparent discrimination in a mixed-feeding herbivore. *Can. J. Zool.* **2011**, *89*, 453.
- [10] S. T. Florin, L. A. Felicetti, C. T. Robbins. The biological basis for understanding and predicting dietary-induced variation in nitrogen and sulphur isotope ratio discrimination. *Funct. Ecol.* **2011**, *25*, 519.
- [11] D. D. Gustine, P. S. Barboza, J. Addison, R. Shively, L. Oliver. Isotopic nitrogen in fecal fiber as an indicator of winter diet in caribou and muskoxen. *Rapid Commun. Mass Spectrom.* **2014**, *28*, 625.
- [12] S. M. Arthur, P. A. Del Vecchio. *Effects of oilfield development on calf production and survival in the Central Arctic Herd, in Research Technical Report*, Juneau, AK, **2009**.
- [13] K. R. Whitten, R. D. Cameron. Nutrient dynamics of caribou forage on Alaska's arctic slope, in *Proc. 2nd International Reindeer/Caribou Symposium*, 17–21 Sept 1979 (Eds: E. Reimers, E. Gaare, S. Skjennberg), Direktoratet for vilt og ferskvannsfisk, Trondheim, **1980**, p. 159.
- [14] A. L. Gallant, E. F. Binnian, J. M. Omernik, M. B. Shasby. *Ecoregions of Alaska*. Report 1567, U.S. Geological Survey, Washington DC, **1995**.
- [15] D. C. Thompson, K. H. McCourt. Seasonal diets of the Porcupine caribou herd. *Am. Midland Naturalist* **1981**, *105*, 70.
- [16] D. E. Russell, A. M. Martell, W. A. C. Nixon. Range ecology of the Porcupine caribou herd in Canada. *Rangifer* **1993**, *13*, 1.
- [17] M. Sturm, C. Racine, K. Tape. Increases in shrub abundance in the arctic. *Nature* **2001**, *411*, 546.
- [18] I. H. Myers-Smith, B. C. Forbes, M. Wilmking, M. Hallinger, T. Lantz, D. Blok, K. D. Tape, M. Macias-Fauria, U. Sass-Klaassen, E. Lévesque, S. Boudreau, P. Ropars, L. Hermanutz, A. Trant, L. S. Collier, S. Weijers, J. Rozema, S. A. Rayback, N. M. Schmidt, G. Schaepman-Strub, S. Wipf, C. Rixen, C. B. Ménard, S. Venn, S. Goetz, L. Andreu-Hayles, S. Elmendorf, V. Ravolainen, J. Welker, P. Grogan, H. E. Epstein, D. S. Hik. Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environ. Res. Lett.* **2011**, *6*, 045509.
- [19] E. Kaarlejarvi, R. Baxter, A. Hofgaard, H. Hytteborn, O. Khitun, U. Molau, S. Sjogersten, P. Wookey, J. Olofsson. Effects of warming on shrub abundance and chemistry drive ecosystem-level changes in a forest-tundra ecotone. *Ecosystems* **2012**, *15*, 1219.
- [20] M. Crete, J. Huot, L. Gauthier. Food selection during early lactation by caribou calving on the tundra in Quebec. *Arctic* **1990**, *43*, 60.
- [21] L. L. VanSomeren. Monitoring energy and nitrogen availability for Arctic caribou (*Rangifer tarandus*). *Unpublished MS thesis*, University of Alaska, Fairbanks (Fairbanks, Alaska), **2014**.
- [22] L. L. VanSomeren, P. S. Barboza, D. P. Thompson, D. D. Gustine. Monitoring digestibility of forages for herbivores: a new application for an old approach. *Can. J. Zool.* **2015**, *93*, 187.
- [23] P. S. Barboza, K. L. Parker. Body protein stores and isotopic indicators of N balance in female reindeer (*Rangifer tarandus*) during winter. *Physiol. Biochem. Zool.* **2006**, *79*, 628.
- [24] J. H. Zar. *Biostatistical Analysis*, (4th edn.). Prentice Hall, Upper Saddle River, NJ, **1999**.
- [25] B. A. Barnett. Carbon and nitrogen isotope ratios of caribou tissues, vascular plants, and lichens from northern Alaska. *Unpublished PhD thesis*, University of Alaska, Fairbanks, **1994**.
- [26] R. R. Pattison, J. M. Welker. Differential ecophysiological response of deciduous shrubs and a graminoid to long-term experimental snow reductions and additions in moist acidic tundra, Northern Alaska. *Oecologia* **2014**, *174*, 339.

- [27] C. Körner, S. Leuzinger, S. Riedl, R. T. Siegwolf, L. Streule. Carbon and nitrogen stable isotope signals for an entire alpine flora, based on herbarium samples. *Alpine Bot.* **2016**, *126*, 153.
- [28] S. M. P. Cahoon, P. F. Sullivan, E. Post. Carbon and water relations of contrasting Arctic plants: implications for shrub expansion in West Greenland. *Ecosphere* **2016**, *7*, 1.
- [29] F. S. Chapin, P. A. Matson, H. A. Mooney. *Principles of Terrestrial Ecosystem Ecology*. Springer, New York, **2002**.
- [30] J. M. G. Hudson, G. H. R. Henry, W. K. Cornwell. Taller and larger: shifts in Arctic tundra leaf traits after 16 years of experimental warming. *Global Change Biol.* **2011**, *17*, 1013.
- [31] M. J. Wooller, G. D. Zazula, M. Edwards, D. G. Froese, R. D. Boone, C. Parker, B. Bennett. Stable carbon isotope compositions of eastern Beringian grasses and sedges: investigating their potential as paleoenvironmental indicators. *Arctic, Antarctic, and Alpine Res.* **2007**, *39*, 318.
- [32] D. R. Bowling, D. E. Pataki, J. T. Randerson. Carbon isotopes in terrestrial ecosystem pools and CO₂ fluxes. *New Phytologist* **2008**, *178*, 24.
- [33] K. Nadelhoffer, G. Shaver, B. Fry, A. Giblin, L. Johnson, R. McKane. N-15 natural abundances and N use by tundra plants. *Oecologia* **1996**, *107*, 386.
- [34] D. D. Gustine, P. S. Barboza, J. P. Lawler, L. G. Adams, K. L. Parker, S. M. Arthur, B. S. Shults. Diversity of nitrogen isotopes and protein status in caribou: implications for monitoring northern ungulates. *J. Mammal.* **2012**, *93*, 778.
- [35] D. H. Mann, P. Groves, M. L. Kunz, R. E. Reaner, B. V. Gaglioti. Ice-age megafauna in Arctic Alaska: extinction, invasion, survival. *Quat. Sci. Rev.* **2013**, *70*, 91.
- [36] P. S. Barboza, K. L. Parker, I. D. Hume. *Integrative Wildlife Nutrition*. Springer, Heidelberg, **2009**.
- [37] J. S. Sedinger. Protein and amino acid composition of tundra vegetation in relation to nutritional requirements of geese. *J. Wildlife Manage.* **1984**, *48*, 1128.
- [38] C. T. Robbins, T. A. Hanley, A. E. Hagerman, O. Hjeljord, D. L. Baker, C. C. Schwartz, W. W. Mautz. Role of tannins in defending plants against ruminants: reduction in protein availability. *Ecology* **1987**, *68*, 98.
- [39] T. N. Barry, W. C. McNabb. The implications of condensed tannins on the nutritive value of temperate forages fed to ruminants. *Br. J. Nutr.* **1999**, *81*, 263.
- [40] M. M. Lorenz, L. Alkhafadji, S. Stringano, S. Nilsson, I. Mueller-Harvey, P. Uden. Relationship between condensed tannin structures and their ability to precipitate feed proteins in the rumen. *J. Sci. Food Agric.* **2014**, *94*, 963.