

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Department of Entomology: Faculty  
Publications

Entomology, Department of

---

3-28-2024

## Corn Rootworm: Biology, Ecology, Behavior, and Integrated Management

Lance J. Meinke

Joseph L. Spencer

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Department of Entomology: Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Editorial

# Corn Rootworm: Biology, Ecology, Behavior, and Integrated Management

Lance J. Meinke <sup>1,\*</sup> and Joseph L. Spencer <sup>2</sup><sup>1</sup> Department of Entomology, University of Nebraska, Lincoln, NE 68583, USA<sup>2</sup> Illinois Natural History Survey, Prairie Research Institute, University of Illinois, Champaign, IL 61820, USA; spencer1@illinois.edu

\* Correspondence: lmeinke1@unl.edu

Species of the beetle genus *Diabrotica* (Coleoptera: Chrysomelidae) are native to North and South America, with their greatest diversity occurring in neotropical areas [1]. Little is known about the biology and ecology of many of the 400 described species, with current knowledge primarily limited to the small number of species that are pests in agricultural systems [1,2]. This Special Issue focuses primarily on key economically important pest species, i.e., *D. virgifera virgifera* LeConte, *D. speciosa* (Germar), *D. balteata* (LeConte), and *D. viridula* (F.).

The western corn rootworm, *D. v. virgifera*, is a key pest of grain maize in North America and the specific focus of many contributions to this Special Issue. The costs of managing this pest and the value of lost production annually exceed USD 2 billion in the U.S.A. This species was also accidentally introduced into Europe, where it is now an established maize pest [3]. The highly adaptable nature of this species has made management an ongoing challenge. Over time, this species has evolved resistance to active ingredients in four insecticide classes, annual crop rotation, and all commercially available rootworm-active Cry toxins (derived from the soil microbe *Bacillus thuringiensis*) expressed in Bt-maize hybrids in the U.S.A. [4–6].

The future success of *Diabrotica* pest management may depend on a more holistic view of management than that implemented in the past. This requires movement away from single-tactic approaches to a combination of tactics deployed within an integrated pest management framework [5]. This will include the conceptualization and development of new tactics that are based on an increased understanding of *Diabrotica* biology, physiology, ecology, and population dynamics [6–10].

This Special Issue provides original research and comprehensive reviews that summarize the current knowledge in key areas of *Diabrotica* biology, ecology, behavior, and management. The contributions include the following:

- An overview of the evolutionary history and host relationships of *Diabrotica* species, plus natural enemies of *Diabrotica* [1];
- The biology and management of *D. v. virgifera* in Europe [3] and *D. speciosa* (Germar), *D. balteata* (LeConte), and *D. viridula* (F.) in South America [2];
- Host-microbe relationships and aspects of chemical ecology that influence *D. v. virgifera* behavior and host plant resistance [9];
- The movement ecology of *D. v. virgifera* and its relation to management [6];
- The potential of RNAi technologies as components of *D. v. virgifera* management strategies [10];
- An overview of *D. v. virgifera* resistance to insecticides and plant-incorporated Bt traits in maize [4,5];
- Advances in *D. v. virgifera* monitoring/sampling technologies [7];
- An overview of the available computer models and modeling approaches to support research on the biology and management of *Diabrotica* species [8].



**Citation:** Meinke, L.J.; Spencer, J.L. Corn Rootworm: Biology, Ecology, Behavior, and Integrated Management. *Insects* **2024**, *15*, 235. <https://doi.org/10.3390/insects15040235>

Received: 21 March 2024

Accepted: 22 March 2024

Published: 28 March 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The history of field-evolved resistance is marked by repeated failures to appreciate the capabilities of pest insects. The western corn rootworm is a good example of this. Therefore, it is our hope that these publications, which refresh our current understanding of *Diabrotica*, will inform future research and efforts to develop novel and sustainable management tactics/strategies.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Eben, A. Ecology and evolutionary history of *Diabrotica* beetles—Overview and update. *Insects* **2022**, *13*, 156. [[CrossRef](#)] [[PubMed](#)]
2. Cabrera Walsh, G.; Ávila, C.; Cabrera, N.; Nava, D.; de Sene Pinto, A.; Weber, D. Biology and management of pest *Diabrotica* species in South America. *Insects* **2020**, *11*, 421. [[CrossRef](#)] [[PubMed](#)]
3. Bažok, R.; Lemić, D.; Chiarini, F.; Furlan, L. Western corn rootworm (*Diabrotica virgifera virgifera* LeConte) in Europe: Current status and sustainable pest management. *Insects* **2021**, *12*, 195. [[CrossRef](#)] [[PubMed](#)]
4. Gassmann, A. Resistance to Bt maize by western corn rootworm: Effects of pest biology, the pest–crop interaction and the agricultural landscape on resistance. *Insects* **2021**, *12*, 136. [[CrossRef](#)] [[PubMed](#)]
5. Meinke, L.; Souza, D.; Siegfried, B. The use of insecticides to manage the western corn rootworm, *Diabrotica virgifera virgifera*, LeConte: History, field-evolved resistance, and associated mechanisms. *Insects* **2021**, *12*, 112. [[CrossRef](#)]
6. Sappington, T.; Spencer, J. Movement ecology of adult western corn rootworm: Implications for management. *Insects* **2023**, *14*, 922. [[CrossRef](#)]
7. Tóth, Z.; Tóth, M.; Jósmai, J.; Tóth, F.; Flórián, N.; Gergőcs, V.; Dombos, M. Automatic field detection of western corn rootworm (*Diabrotica virgifera virgifera*; Coleoptera: Chrysomelidae) with a new probe. *Insects* **2020**, *11*, 486. [[CrossRef](#)] [[PubMed](#)]
8. Onstad, D.; Caprio, M.; Pan, Z. Models of *Diabrotica* populations: Demography, population genetics, geographic spread, and management. *Insects* **2020**, *11*, 712. [[CrossRef](#)] [[PubMed](#)]
9. Paddock, K.; Robert, C.; Erb, M.; Hibbard, B. Western corn rootworm, plant and microbe interactions: A review and prospects for new management tools. *Insects* **2021**, *12*, 171. [[CrossRef](#)] [[PubMed](#)]
10. Darlington, M.; Reinders, J.; Sethi, A.; Lu, A.; Ramaseshadri, P.; Fischer, J.; Boeckman, C.; Petrick, J.; Roper, J.; Narva, K.; et al. RNAi for western corn rootworm management: Lessons learned, challenges, and future directions. *Insects* **2022**, *13*, 57. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.