

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of
Nebraska-Lincoln Extension

Extension

1993

G93-1136 Potato Leafhopper Management in Alfalfa

Stephen D. Danielson

University of Nebraska - Lincoln, sdanielson1@unl.edu

Keith J. Jarvi

University of Nebraska - Lincoln, kjarvi1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>



Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

Danielson, Stephen D. and Jarvi, Keith J., "G93-1136 Potato Leafhopper Management in Alfalfa" (1993).
Historical Materials from University of Nebraska-Lincoln Extension. 1103.
<https://digitalcommons.unl.edu/extensionhist/1103>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Potato Leafhopper Management in Alfalfa

This NebGuide explains how to detect and manage potato leafhoppers to minimize alfalfa damage.

Steve Danielson, Extension Entomology Specialist
Keith Jarvi, Extension Integrated Pest Management Assistant

- [Appearance](#)
- [Migration Habits](#)
- [Damage Symptoms](#)
- [Early Detection](#)
- [Management](#)
- [Insecticides for Potato Leafhopper Control](#)

The potato leafhopper is capable of causing serious damage to alfalfa in Nebraska. This insect overwinters in the gulf states and migrates northward in the spring, usually in April or early May. During the summer months, several generations develop while feeding primarily on legumes such as alfalfa and clover. Although an occasional host, soybeans rarely suffer economic damage, particularly with the pubescent (i.e. hairy) soybean varieties that are commonly grown today. Infested potatoes also can sustain economic damage from this insect. Generally, the potato leafhopper is an economic pest in only the eastern one-third of the state but it will occasionally spread and cause major problems in other portions of the state.

Appearance

The adult and immature (i.e. nymph) stages of the potato leafhopper are very similar in appearance (*Figure 1*), with only the adults having wings. They are light green in color and slender in shape.




Figure 1. Adult (bottom) and nymph potato leafhopper.

The adults are approximately 1/8 inch long (*Figure 2*). Eggs are laid in plant tissue and are difficult to observe in the field. Both adults and nymphs have piercing-sucking mouthparts on the underside of the head that they insert into the plant to obtain liquid nourishment. The wingless nymphs often move backwards and sideways in a crablike fashion on the leaf surfaces.

Migration Habits

The potato leafhopper overwinters in the gulf states of the United States and migrates northward in the spring. It is capable of flying short distances of up to a few miles on its own. It is believed that long distance

movement is only possible when large scale weather systems are present to assist them. They travel with the same winds that move gulf moisture northward in the spring. Since these small insects are at the mercy of weather systems, their distribution patterns vary tremendously from year to year. This variability accounts for the fact that Nebraska often escapes serious infestations of the potato leafhopper. Eastern states experience heavier and more frequent rains as a result of gulf moisture moving in their direction with regularity, and therefore have serious infestations of this pest nearly every year.




Figure 2. Adult potato leafhopper.

Adults are capable of short distance movement from one field to another. This commonly occurs when alfalfa fields are harvested. Nymphs are not capable of such movement because they lack wings. As a result, they often die of starvation when fields are harvested.

Damage Symptoms

Feeding by leafhoppers disrupts the flow of water and nutrients inside the plant and the resulting symptoms of damage are often mistaken as the results of drought or disease (*Figure 3*). The first sign of damage is a yellow discoloration (i.e. chlorosis) of the leaflets that begins at the tip and progresses toward the base in a v-shaped pattern. This is often referred to as "hopperburn" or "tipburn." As damage continues to develop, the chlorotic areas become purplish in color and, if severe enough, the leaflet will turn brown and die. Severely affected plants become stunted and entire stems or plants may eventually die.



Figure 3. Symptoms of potato leafhopper damage on alfalfa.

Newly planted alfalfa stands are not very tolerant of this feeding damage and may be completely destroyed by the leafhopper. Established stands are more tolerant of the damage and will often recover once the leafhopper feeding ceases. Unfortunately, yields will be reduced as a result of leafhopper feeding even if the stand recovers.

Early Detection

Proper management of the potato leafhopper requires early detection in late May, June and early July before symptoms of damage appear. This insect is quite small and is not easy to see when casually walking through a field or even when down on your knees at plant height. Occasionally, you will see the leafhoppers hopping or flying ahead as you walk through an infested field, but this method does not provide accurate estimates of leafhopper abundance.

The use of a sweep net will allow you to obtain counts of leafhoppers. These estimates can be compared to treatment guidelines to assist in making management decisions. Sweep nets with 15-inch diameter openings are used for collecting the potato leafhopper in alfalfa. The sweep net is moved vigorously through the foliage in 180o arcs at several locations in each field. To obtain meaningful, representative estimates of leafhopper abundance, at least 25 sweeps should be taken from each of four locations in each field. The number of leafhoppers per sweep is obtained by counting the leafhoppers in the net and dividing by the number of sweeps (or arcs) taken in the field. Sweep nets can be obtained from various suppliers (two are listed below) or homemade nets can be assembled from a broomstick, some rigid wire and a pillow case.

BioQuip Products

17803 LaSalle Avenue
Gardena, CA 90248-3602
Telephone (310) 324-0620

Pest Management Supply, Inc.

P.O. Box 938

Amherst, MA 01004

Telephone (413) 253-3747

Management

Growers are encouraged to plant alfalfa varieties that are known to be resistant to the potato leafhopper whenever possible. This type of resistance is fairly reliable and will reduce the amount of damage that occurs from a given infestation level of the leafhopper. In severe cases, insecticide treatment may be required even with resistant varieties.

Research has shown that economic damage has already occurred once the chlorotic symptoms are visible. Economic damage is that which equals or exceeds the cost of an insecticide treatment. Therefore, alfalfa growers are urged to begin sampling their fields in late May and June and to continue on at least a weekly schedule through the remainder of the season. It is not a wise plan to wait until the damage appears before making a decision to manage this insect.

The following tables provide guidance for managing the potato leafhopper in alfalfa that is up to 12 inches tall. They provide a range of hay values and treatment costs. If leafhopper numbers equal or exceed the numbers in the table for the appropriate alfalfa height, hay value and treatment cost, an insecticide application should be considered to reduce the leafhopper infestation and protect the crop from further damage. An example using *Table I* involves an insecticide application cost of \$12 per acre and hay value of \$100 per ton with 1-4 inch alfalfa. In this case, the treatment threshold is 0.4 potato leafhoppers per sweep.

If leafhopper numbers have reached or exceeded the treatment threshold and the alfalfa is more than 12 inches tall, immediate harvest is suggested rather than insecticide treatment. Harvesting will cause the adult leafhoppers to move out of the field and most of the nymphs will die due to starvation and exposure. Fields can be reinfested from other nearby fields after having been treated with insecticides or harvested, so weekly scouting through the season is suggested. Growers should be aware that severely damaged alfalfa will need to be cut to allow recovery and regrowth to occur.

Value of hay (per ton)	Cost of insecticide application (per acre)					
	\$8	\$10	\$12	\$14	\$16	\$20
\$ 60	0.4	0.5	0.6	0.7	0.8	1.0
\$ 80	0.3	0.4	0.5	0.5	0.6	0.75
\$100	0.25	0.3	0.4	0.4	0.5	0.6
\$120	0.2	0.25	0.3	0.35	0.4	0.5
\$140	0.2	0.2	0.25	0.3	0.3	0.4
\$160	0.15	0.2	0.3	0.3	0.3	0.4

Table II. Dynamic Treatment Thresholds for Potato Leafhoppers (average number per sweep) on Alfalfa that is 4 to 8 inches tall.

Value of hay (per ton)	Cost of insecticide application (per acre)					
	\$8	\$10	\$12	\$14	\$16	\$20
\$ 60	0.7	0.8	1.0	1.0	1.3	1.7
\$ 80	0.6	0.6	0.75	0.9	1.0	1.3
\$100	0.4	0.5	0.6	0.7	0.8	1.0
\$120	0.3	0.4	0.5	0.6	0.7	0.8
\$140	0.3	0.35	0.4	0.5	0.6	0.7
\$160	0.25	0.3	0.4	0.4	0.5	0.6

Table III. Dynamic Treatment Thresholds for Potato Leafhoppers (average number per sweep) on Alfalfa that is 8 to 12 inches tall.

Value of hay (per ton)	Cost of insecticide application (per acre)					
	\$8	\$10	\$12	\$14	\$16	\$20
\$ 60	2.0	2.4	2.8	3.0	3.9	5.0
\$ 80	1.8	1.9	2.2	2.7	3.0	4.0
\$100	1.2	1.5	1.8	2.1	2.4	3.0
\$120	0.9	1.2	1.5	1.8	2.1	2.4
\$140	0.9	1.0	1.2	1.5	1.8	2.0
\$160	0.8	0.9	1.0	1.2	1.5	1.8

Insecticides for Potato Leafhopper Control

A number of insecticides are registered for the control of this insect on alfalfa. Refer to the current version of the *Insect Management Guide for Nebraska Alfalfa, Soybeans, Wheat, Range, and Pasture (EC-1511)* for additional information about potato leafhopper management and for a listing of currently registered insecticides.

File G1136 under: INSECTS AND PESTS

C-34, Field Crops

Paper version issued March 1993; 7,000 printed.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.

University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.