

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Papers and Publications in Animal
Science

Animal Science Department

4-9-2018

Performance and carcass characteristics of steers fed with two levels of metabolizable energy intake during summer and winter season

R A. Arias

J P. Keim

M Gandarillas

A Velasquez

C Alvarado-Gilis

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/animalscifacpub>



Part of the [Genetics and Genomics Commons](#), and the [Meat Science Commons](#)

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

R A. Arias, J P. Keim, M Gandarillas, A Velasquez, C Alvarado-Gilis, and T L. Mader

Performance and carcass characteristics of steers fed with two levels of metabolizable energy intake during summer and winter season

R. A. Arias^{1,2†}, J. P. Keim¹, M. Gandarillas¹, A. Velásquez^{3,4}, C. Alvarado-Gilis¹ and T. L. Mader⁵

¹Instituto de Producción Animal, Universidad Austral de Chile, Valdivia 5090000, Chile; ²Centro de Investigación en Suelos Volcánicos, Universidad Austral de Chile, Valdivia 5090000, Chile; ³Escuela de Agronomía, Facultad de Recursos Naturales, Universidad Católica de Temuco, Temuco 4781312, Chile; ⁴Núcleo de Investigación en Producción Agroalimentaria, Universidad Católica de Temuco, Temuco 4781312, Chile; ⁵Animal Science Department, University of Nebraska-Lincoln, NE 68028, USA

(Received 7 June 2017; Accepted 9 April 2018; First published online 22 May 2018)

Climate change is producing an increase on extreme weather events around the world such as flooding, drought and extreme ambient temperatures impacting animal production and animal welfare. At present, there is a lack of studies addressing the effects of climatic conditions associated with energy intake in finishing cattle in South American feed yards. Therefore, two experiments were conducted to assess the effects of environmental variables and level of metabolizable energy intake above maintenance requirements (MEI) on performance and carcass quality of steers. In each experiment (winter and summer), steers were fed with 1.85 or 2.72 times of their requirements of metabolizable energy of maintenance. A total of 24 crossbred steers per experiment were used and located in four pens (26.25 m²/head) equipped with a Calan Broadbent Feeding System. Animals were fed with the same diet within each season, varying the amount offered to adjust the MEI treatments. Mud depth, mud scores, tympanic temperature (TT), environmental variables, average daily gain, respiration rates and carcass characteristics plus three thermal comfort indices were collected. Data analysis considered a factorial arrangement (Season and MEI). In addition, a repeated measures analysis was performed for TT and respiration rate. Mean values of ambient temperature, solar radiation and comfort thermal indices were greater in the summer experiment as expected ($P < 0.005$). The mean values of TT were higher in steers fed with higher MEI and also in the summer season. The average daily gain was greater during summer v. winter (1.10 ± 0.11 v. 0.36 ± 0.06) kg/day, also when steers were fed 2.72 v. 1.85 MEI level (0.89 ± 0.12 v. 0.57 ± 0.10) kg/day. In summer, respiration rate increased in 41.2% in the afternoon. In winter, muddy conditions increased with time of feeding, whereas wind speed and rainfall had significant effects on TT and average daily gain. We conclude that MEI and environmental variables have direct effects on the physiology and performance of steers, including TT and average daily gain, particularly during the winter. In addition, carcass characteristics were affected by season but not by the level of MEI. Finally, due to the high variability of data as well as the small number of animals assessed in these experiments, more studies on carcass characteristics under similar conditions are required.

Keywords: tympanic temperature, muddy conditions, respiration rate, panting scores, climatic factors

Implications

Beef production systems of temperate regions are experiencing important challenges due to the changes in the environmental conditions as well as the major concern of consumers by animal well-being. This study was conducted to assess the effects of metabolizable energy intake on the physiology, performance and carcass characteristics of finished steers during the summer and winter seasons. Results indicate that weather conditions

affected animal performance, particularly wind speed and rainfall in the winter. Thus, cattlemen must consider weather conditions of their regions in their feed yard design as well as their feeding management, particularly the energy intake level.

Introduction

Climate conditions have a great importance on animal production because it impacts the surrounding environment in which animals are raised and reproduced (Mader and Gaughan, 2011). In fact, two of three major welfare issues

† E-mail: rodrigo.arias@uach.cl

involve environmental subjects, muddy conditions and heat stress (Grandin, 2016). Animals are continuously attempting to achieve homeothermic status, consequently the thermo-regulation is a very dynamic process. In addition, it has been established that the environment is a potent factor influencing the rate of energy loss and thereby playing a significant role in the energetic efficiency (National Research Council, 1981). Therefore, season of year affects animal's maintenance energy requirement, which in turn impacts animal performance (Birkelo *et al.*, 1991), particularly in average daily gain (ADG) and feed to gain ratio (F:G) (Birkelo and Johnson, 1993). Previous research has demonstrated that cattle in cold wet muddy conditions require more energy and more effort to walk, resulting in lower performance (Dijkman and Lawrence, 1997). In addition, many feed yards in South America have not been properly designed nor has pen surface been properly prepared. Grandin (2016) pointed out recently, the lack of peer reviewed scientific studies on the effects of mud on beef cattle. Therefore, a need to study the effects of environmental variables under these particular conditions can provide valuable information to producers and cattle industry in these regions. In addition, there is limited information about the effects of climate on beef production in temperate regions of southern South America. The objective of the present study was to assess the effects of environmental conditions and level of metabolizable energy intake on performance and carcass quality of finished steers.

Material and methods

Two experiments were conducted at the Experimental Research Station of the Universidad Católica de Temuco in Temuco City, Chile, with the approval of the ethics committee for research.

Facilities and animals

It consisted of four pens (six head/pen; 26.25 m²/head) equipped with a Calan Broadbent Feeding System (American CALAN Inc., Northwood, NH, USA), which in turn had a shed protecting the feeding area to prevent equipment and feed-stuffs from being affected by rainfall. The facility was located

15 km northeast of Temuco city. Water tanks were shared between fences of pens one-two and three-four. In both experiments, steers were provided by a private company (Ferias Araucanía S.A.) Once steers arrived at the facilities, they were sorted by BW, assigned randomly to two blocks (light or heavy) and then allocated into one of two treatments: T1 = 1.85 times or T2 = 2.72 times metabolizable energy intake above maintenance (MEI), according to requirements estimated on BW by using the software of the National Academies of Sciences Engineering and Medicine (2016).

Experiment 1 (Winter). The experiment had 91 days on feed, beginning on 4 July 2013 and finishing on October 03 of 2013. A total of 24 crossbred steers (Angus × Hereford; BW = 420 kg ± 4.4 kg) arrived on 13 June 2013. There was a pre-experimental period (21 days) that allowed steers to get used and trained with the feeding system and adapted to the facilities. Throughout this period steers were fed *ad libitum* (100% haylage, Table 1) with free access to water. Subsequently, in the experimental period, steers were fed once per day with the experimental diet (Table 1). The amount of feed offered by group was different in order to adjust the MEI as was established for the treatments.

Individual animal BW were recorded by using an electronic scale (Gallagher W300, Hamilton, New Zealand), and then used to calculate average daily gain (ADG, g/day). Likewise, mud depth was recorded twice during the experimental period (23 July 2013 and 21 September 2013) in three different locations of each pen and then averaged. In addition, the amount of mud in the animal's body was also estimated by using a simple mud score (1 = clean; 2 = dirty legs; 3 = dirty legs and belly; 4 = dirty legs, belly and partial sides of the animal; and 5 = dirty legs, belly and sides of the animal). Animals received a veterinary treatment (Triclabendazole 10%, 10 ml/100 kg and Ivermectin 1 cc/50 kg), and a growth promoting implant (140 mg Trenbolone Acetate + 20 mg estradiol; Intervet, Schering-Plough Animal Health, Kenilworth, NJ, USA) before the experimental period on 3 July 2013.

Table 1 Amount of ingredients (dry matter (DM) basis, kg/day) at the beginning of the experimental periods for the two studies of finishing steers

Ingredients ¹	Winter ²		Summer ²		DM content (%)		Energy density (Mcal ME/kg DM)		CP content (%)	
	1.85 ×	2.72 ×	1.85 ×	2.72 ×	Winter	Summer	Winter	Summer	Winter	Summer
Silage	4.0	5.8	4.1	6.0	46.3	40.0	2.2	2.5	11.8	11.5
Lupine	0.5	0.8	–	–	89.7	–	2.8	–	29.5	–
Triticale	1.9	2.8	–	–	87.9	–	3.3	–	11.5	–
Canola meal	–	–	0.4	0.6	–	89.9	–	2.7	–	36.7
Oat	–	–	2.0	2.9	–	87.9	–	2.8	–	11.0
Minerals	0.1	0.1	0.1	0.1	100.0	100.0	–	–	–	–

ME = metabolizable energy.

¹Diet composition: Experiment 1 (winter): 62% haylage, 8% lupine whole grain and 29% triticale whole grain; Experiment 2 (summer): 64% haylage, 5% canola meal and 30% oat whole grain. Diet energy density in Exp 1 = 2.69 EM/kg DM and 13.03% CP; diet energy density in Exp 2 = 2.51 EM/kg DM and 13.04% CP. Salt minerals included a ionophore (Lasalocid sodium 3 mg/g).

²1.82 × and 2.72 × = 1.82 or 2.72 times metabolizable energy intake above maintenance.

Tympanic temperature. A total of 16 steers (eight per treatment, four by pen) were randomly chosen to receive a iButton device to collect TT (Maxim Integrated Products Inc., San José, CA, USA). This device was fitted manually into the tympanic canal in the ear allowing the collection of data continuously. After the device was placed, this was covered with absorbent material (Tampax Palitex Edgewell), and subsequently with pipe foam (Isoplast) to give firmness to the structure of the ear pinna, which was wrapped with an elastic bandage (Coban Nexcare 3M, St. Paul, MN, USA). Finally, the elastic bandage was wrapped with adhesive tape for medical use (5 cm wide Leukoplast, BSN Medical, Hamburg, Germany). The devices were programmed to collect TT at intervals of 10 min and installed on 31 July 2013. Subsequently, data were compiled into hourly readings. The criterion to install the devices was made considering the weather forecast. The retrieving of the devices was made 10 days later. During this experiment, a total of six devices were lost (three in each treatment group), finishing with five devices per treatment.

Environmental data collection and thermal comfort indices. Ambient temperature (AT, °C), wind speed (WS, m/s), relative humidity (RH, %), total solar radiation (SR, W/m²) and precipitation (PP, mm/day) were collected continuously at 15 min intervals by using a weather station (Campbell Scientific CR1000, Utah, USA) located at 5 km southeast from the research site. Later, these data were compiled into hourly values to match TT dataset. In addition, these climatic data were used to calculate three thermal comfort indices: (a) Comprehensive climate index, CCI (Mader *et al.*, 2010); (b) temperature–humidity index, THI (Hahn *et al.*, 2009); and (c) the THI adjusted by wind speed and solar radiation, THI_{adj} (Mader *et al.*, 2006), according to equations given below:

$$\text{THI} = 0.8 \times \text{AT} + ((\text{RH}/100) \times (\text{AT} - 14.4)) + 46.4 \quad (1)$$

$$\text{THI}_{\text{adj}} = \text{THI} + 4.51 - (1.992 \times \text{WS}) + (0.068 \times \text{SR}) \quad (2)$$

$$\text{CCI} = \text{AT} + \text{FRH} + \text{FWS} + \text{FSR} \quad (3)$$

Where:

FRH corresponds to the relative humidity correction factor;

$$e^{(0.00182 \times \text{RH} + 1.8 \times 10^{-5} \times \text{AT} \times \text{RH})} \times 0.000054 \times \text{AT}^2 + 0.00192 \times \text{AT} - 0.0246 \times \text{RH} - 30$$

FWS corresponds to the wind speed correction factor;

$$\frac{-6.56}{e^{\left\{ \frac{1}{[2.26 \times \text{WS} + 0.23^{0.45}]} \times [2.9 + 1.14 \times 10^{-6} \times \text{WS}^{2.5} - \log_{0.3}(2.26 \times \text{WS} + 0.33)^{-2}] \right\}}} - 0.0556 \times \text{WS}^2 + 3.33$$

and FSR corresponds to the solar radiation correction factor;

$$0.0076 \times \text{SR} - 0.00002 \times \text{AT} + 0.00005 \times \text{AT}^2 \times \sqrt{\text{SR}} + 0.1 \times \text{AT} - 2$$

Experiment 2 (Summer). The experiment had 95 days on feed, beginning on 23 January 2014 and finishing on 28 April 2014. A total of 24 crossbred (Angus × Hereford) steers (BW = 431 kg ± 10.4 kg) arrived on 4 January 2014. There was a pre-experimental period of 19 days with a diet similar than in Exp 1. In order to fit the MEI supply for treatments T1 and T2 feed intake was different, with a similar approach than in Exp 1. Tympanic temperatures, environmental data and thermal comfort indices were also collected in a similar manner as described in Exp 1. In addition, panting scores (PS) and respiration rates (RR) were collected during 3 days, in the hottest week of the summer (Table 1), from 13 February 2014 to 15 February 2014, according to procedure described by Mader *et al.* (2006). A total of 22 iButton devices were installed in steers (T1 = 12 and T2 = 10) on 6 February 2014 and retrieved 10 days later. Four devices were lost during this experiment (three in T2 and one in T1), ending up with nine devices for T1 and T2. In addition, insecticide-impregnated cattle ear tags were applied for control of horn flies (Moskimat Drag pharma/Expert Plus Intervet Schering-Plough Animal Health, 11 January 2014) and the same growth promoting implant described in Exp 1 on 23 January 2014. During the summer mud depth was not measured, because no mud was present.

Respiration rates (breaths per minute = bpm) were collected twice by counting 20 movements on each animal's flank and timing the required time during those days, in the morning starting at 0900 h and in the afternoon at 1500 h. The criterion to install the devices and measure RR was done based on weather forecast.

Carcass data collection. Steers were slaughtered in Temuco city after 91 and 95 days on feed for experiments 1 and 2 respectively. In both experiments, the goal was to reach a minimum of 90 days on feeding. Carcass data included: cold carcass weight, rib eye area, longissimus muscle pH, fat coverage, kidney–pelvic–heart fat (KPH), back fat and marbling. After slaughtering, carcasses were kept in cold chambers at a temperature of 3°C by 48 h. Later the carcasses were opened between the 12th and 13th (Exp 1) and 9th and 10th rib (Exp 2) from where data were collected. The complete external fat coverage was estimated by the official certification personnel of the slaughterhouse (Tecno-Carnes Ltda.), based on the Chilean Standard of Classification (INN, 2002), as well as the muscle pH values. In Exp 1 data of 23 animals were collected because one animal from 2.72 EMm diet was removed from the experiment 50 days before the slaughter date due to a minor leg injury not related with the treatments, which affected its normal displacement in the pen.

Statistical analysis. Only days with full TT and weather data were considered, for the analysis. The analysis was performed considering a randomized complete block design (blocking factor = BW) with a factorial arrangement of 2 × 2, with season and level of MEI as main factors. Each animal was considered as an observational and experimental unit, the level of significance was 0.05. All numerical variables were analysed by ANOVA test. The statistical model used for

the analysis was: $Y_{ijk} = \mu + \alpha_i + \tau_j + \beta_k + (\alpha\tau)_{ij} + \epsilon_{ijk}$, where μ is the general mean, α_i the level of MEI effect, τ_j the season effect, $(\alpha\tau)_{ij}$ the interaction of season and MEI, β_k the block effect and ϵ_{ijk} the experimental error. The statistical package used for the analysis was JMP 11 (SAS Institute, Cary, NC, USA). Likewise, TT and RR were modelled using a repeated measurements analysis (MIXED procedure of SAS 9.1; SAS Institute) with TT as the dependent variable with day, season, hour and MEI levels and their interaction as independent variables in the model, being the hour the repeated measured. The random effect was animal (MEI). In addition, RR was also a dependent variable with day, hour and MEI levels and their interaction as independent variables in the model. The random effect was also animal (MEI). Finally, a principal component analysis was performed for the following variables: cold carcass weight (CCW, kg); back fat (BF, mm); rib eye area (RA, cm²); dressing (%); beef pH; final body weight (FBW, kg); tympanic temperature (TT, °C); and the mean values of the climatic variables (AT, WS, SR, RH, PP) for those days in which TT was also collected. The analysis was performed in JMP statistical package (JMP 12.0, SAS Institute) based on the correlation matrix given the scale differences in the variables included. The correlations were estimated by using the REML method.

Results

Weather variables and tympanic temperatures

Table 2 shows a summary of daily mean values (\pm standard error of the mean) for meteorological variables and TT. As

expected, mean values of AT, SR, THI, THI_{adj} and CCI were greater in the summer by $\Delta = 9.6^\circ\text{C}$, $\Delta = 196.0 \text{ W/m}^2$, $\Delta = 15.0$, $\Delta = 30.9$ and $\Delta = 15.8^\circ\text{C}$, respectively. Daily mean values of RH, WS and PP were greater in the winter experiment by $\Delta = 15.1\%$, $\Delta = 1.3 \text{ m/s}$ and $\Delta = 11.6 \text{ mm/day}$, respectively. During the summer, daily mean TT presented higher values and also a more homogeneous pattern than during the winter (Figure 1) independent of the level of MEI. By contrast, values of TT were much more variable in the winter, presenting a strong drop on day 2. This coincides with an increase in the values of WS and PP (Table 2 and Figure 1). After the fifth day in the winter study, TT decreased again but in this case in a lower magnitude, in that day PP was the highest but WS was moderate.

There was a trend for an interaction season \times MEI on mean daily TT ($P = 0.09$). There was also an effect of main factors ($P < 0.001$ and $P = 0.0063$). Steers fed with 1.85 and 2.72 times MEI averaged 38.0°C and $38.1^\circ\text{C} \pm 0.01^\circ\text{C}$ during summer experiment, whereas in the winter experiment averaged 37.4°C and $37.5^\circ\text{C} \pm 0.02^\circ\text{C}$, respectively.

No interaction of season \times MEI \times hour was found for hourly mean TT ($P = 0.9862$), but there were interactions for season \times hour ($P < 0.0001$), season \times MEI ($P < 0.0001$) and hour effect ($P < 0.0001$). Figure 2 shows the hourly mean TT values for both seasons and MEI levels fed to steers. In general, patterns of TT were similar between seasons, showing minimum TT early during the morning (0700 to 0800 h) to quickly increase later during the day, reaching the maximum TT between 1700 and 1800 h. Nevertheless, during the winter the increase in TT

Table 2 Summary of average daily values for climate and tympanic temperature (standard error of the mean) of finished steers fed with two levels of metabolizable energy intake above maintenance requirements during days of tympanic temperature data collection in both seasons

	AT (°C)	RH (%)	WS (m/s)	SR (W/m ²)	THI	THI _{adj}	CCI (°C)	PP (mm/day)	TT (°C)
Winter 2013 (date)									
1 August 2013	3.8 \pm 0.5	99.3 \pm 1.0	0.8 \pm 0.1	34.7 \pm 10.6	38.9 \pm 0.8	42.0 \pm 0.8	2.5 \pm 0.6	0.2 \pm 0.0	37.4 \pm 0.05
2 August 2013	7.8 \pm 0.5	85.9 \pm 0.5	2.2 \pm 0.3	68.0 \pm 23.8	46.7 \pm 1.0	47.2 \pm 0.7	4.4 \pm 0.4	1.8 \pm 0.1	37.9 \pm 0.04
3 August 2013	8.1 \pm 0.3	87.8 \pm 0.7	3.5 \pm 0.2	40.8 \pm 13.5	47.2 \pm 0.5	45.1 \pm 0.5	2.0 \pm 0.4	18.5 \pm 0.2	37.5 \pm 0.04
4 August 2013	9.3 \pm 0.2	86.6 \pm 0.6	6.2 \pm 0.2	14.3 \pm 4.2	49.5 \pm 0.3	41.8 \pm 0.5	1.0 \pm 0.2	19.9 \pm 0.2	37.2 \pm 0.05
5 August 2013	8.7 \pm 0.3	81.8 \pm 0.5	3.3 \pm 0.3	52.4 \pm 16.5	48.6 \pm 0.5	46.9 \pm 0.7	3.4 \pm 0.5	4.2 \pm 0.1	37.4 \pm 0.05
6 August 2013	7.3 \pm 0.2	89.5 \pm 0.5	3.8 \pm 0.5	6.0 \pm 1.8	45.8 \pm 0.3	42.7 \pm 1.0	1.2 \pm 0.7	39.9 \pm 0.4	37.3 \pm 0.04
7 August 2013	6.1 \pm 0.4	87.9 \pm 0.3	2.6 \pm 0.2	56.9 \pm 15.9	43.9 \pm 0.8	43.6 \pm 0.7	1.1 \pm 0.5	6.2 \pm 0.1	37.4 \pm 0.04
8 August 2013	4.6 \pm 0.6	89.6 \pm 0.5	1.1 \pm 0.1	88.7 \pm 26.4	41.0 \pm 1.1	43.9 \pm 1.1	3.0 \pm 0.7	2.2 \pm 0.6	37.6 \pm 0.04
Summer 2014 (date)									
7 February 2014	15.7 \pm 0.7	79.1 \pm 1.6	1.8 \pm 0.2	144.8 \pm 40.2	59.7 \pm 1.0	61.6 \pm 1.2	15.8 \pm 1.1	0 \pm 0.0	38.1 \pm 0.03
8 February 2014	16.2 \pm 0.6	81.2 \pm 2.6	1.4 \pm 0.2	192.5 \pm 46.7	60.6 \pm 0.8	63.6 \pm 0.8	18.2 \pm 0.6	0 \pm 0.0	38.2 \pm 0.02
9 February 2014	15.2 \pm 1.1	76.4 \pm 4.2	1.6 \pm 0.2	269.6 \pm 70.5	58.1 \pm 1.7	61.1 \pm 1.7	16.3 \pm 1.4	0 \pm 0.0	37.9 \pm 0.03
10 February 2014	14.8 \pm 1.3	79.2 \pm 4.3	1.4 \pm 0.3	258.4 \pm 68.9	57.3 \pm 2.0	60.9 \pm 1.9	16.8 \pm 1.5	0 \pm 0.0	38.1 \pm 0.02
11 February 2014	15.7 \pm 0.7	80.6 \pm 2.0	1.8 \pm 0.2	195.3 \pm 55.5	59.7 \pm 1.0	62.0 \pm 0.9	16.5 \pm 0.8	0 \pm 0.0	38.1 \pm 0.02
12 February 2014	15.4 \pm 1.0	71.4 \pm 4.2	2.1 \pm 0.3	287.6 \pm 70.1	58.4 \pm 1.5	60.7 \pm 1.4	16.1 \pm 1.1	0 \pm 0.0	38.0 \pm 0.02
13 February 2014	18.6 \pm 1.8	58.2 \pm 4.9	1.7 \pm 0.3	277.2 \pm 67.7	61.7 \pm 2.3	64.8 \pm 2.2	20.4 \pm 1.9	0 \pm 0.0	38.1 \pm 0.03
14 February 2014	19.5 \pm 1.6	60.8 \pm 5.0	1.2 \pm 0.2	273.9 \pm 67.2	63.3 \pm 2.0	67.3 \pm 2.0	22.5 \pm 1.7	0 \pm 0.0	38.1 \pm 0.02
15 February 2014	18.6 \pm 1.5	68.4 \pm 4.5	1.5 \pm 0.3	272.1 \pm 67.1	62.6 \pm 2.0	65.9 \pm 2.0	21.1 \pm 1.6	0 \pm 0.0	38.1 \pm 0.03
16 February 2014	16.3 \pm 1.1	79.6 \pm 3.4	1.8 \pm 0.3	240.8 \pm 65.0	60.2 \pm 1.6	62.7 \pm 1.6	17.6 \pm 1.4	0 \pm 0.0	38.0 \pm 0.03
Winter week mean	7.1 \pm 0.1	73.5 \pm 0.2	3.0 \pm 0.1	45.2 \pm 1.8	45.5 \pm 0.1	40.9 \pm 0.1	2.4 \pm 0.1	0.5 \pm 0.0	37.6 \pm 0.02
Summer week mean	16.6 \pm 0.1	88.3 \pm 0.3	1.6 \pm 0.1	241.3 \pm 4.6	60.2 \pm 0.1	45.0 \pm 0.3	18.0 \pm 0.1	0 \pm 0.0	38.1 \pm 0.01

AT = air ambient temperature; RH = relative humidity; WS = wind speed; SR = solar radiation; THI = temperature–humidity index (dimensionless); THI_{adj} = temperature–humidity index (dimensionless) adjusted by WS and SR; CCI = comprehensive climate index (°C); PP = rainfall; TT = tympanic temperature.

was less sharp than during the summer. In addition, steers showed a lower TT during the winter season ($\Delta = 0.62^\circ\text{C}$ less), through hours of the day, independent of the MEI level fed. Only in the winter season, there were some hours of the day when steers fed with higher MEI showed different TT (or tended to be different), than steers fed 1.85 times MEI. These differences were observed at 1200; 1500 and 2100 h ($P = 0.07$; $P = 0.04$; and $P = 0.08$, respectively).

Animal performance and carcass quality

No interaction between season \times MEI level was observed for ADG ($P = 0.7859$), but there was an effect of season




Figure 1 Mean daily tympanic temperatures of steers fed with two levels of metabolizable energy intake above maintenance requirements and by season for those periods of data collection (Sum = summer season and Win = Winter season; 1.85 MEI = 1.85 times metabolizable energy intake and 2.72 MEI = 2.72 times metabolizable energy intake).




Figure 2 Mean hourly tympanic temperatures of steers fed with two levels of metabolizable energy intake above maintenance requirements during the summer and winter experiments (1.85 MEI = 1.85 times metabolizable energy intake and 2.72 MEI = 2.72 times metabolizable energy intake).

($P < 0.0001$) and MEI level ($P = 0.0088$). The performance of steers was better during the summer period in comparison to the winter period (1.10 ± 0.11 v. 0.36 ± 0.06) kg/day, as presented in Figure 3. Similarly, animals fed with the higher MEI level across seasons, showed a greater ADG (0.89 ± 0.12 v. 0.57 ± 0.10) kg/day.

A summary of carcasses variables across seasons and MEI levels are presented in Table 3. Only cold and hot carcass weight showed an interaction between season \times MEI level. However, for all the variables there was a season effect ($P < 0.0001$), with the exception of Back fat, that ranged from 2.93 to 4.46 mm. Values of muscle pH were lower and less variables during the summer season, with only two samples showing values above 5.8 but below 6.0, whereas 13 samples were observed in the winter study, with four of them with pH values above 6.0. The rib eye area was 2.1 times greater in the summer experiment, but the KPH was 1.8 times greater in the winter experiment. Fat coverage was greater in the summer time, but the dressing percentage was lower by $\Delta = 6\%$.

Environmental stressors

During Exp 1, mud accumulated in the pens increased ($P = 0.0261$) with days of feeding, averaging 11.2 and 16.1 ± 1.7 cm for 23 July 2013 and 21 September 2013, respectively. No differences were observed for mud depth between pens for each date. In addition, no differences in mud scores (MS) were observed for 23 July 2013 ($P = 0.46$), $MS = 3.37$ v. 3.95 ± 0.22 for 1.85 and 2.72 times MEI, respectively, neither for 21 September 2013 ($P = 0.38$), $MS = 4.16$ v. 4.40 ± 0.20 for 1.85 and 2.72 times MEI. However, there was a difference between the two dates

($P=0.0036$), $MS=3.82$ v. 4.28 ± 0.15 for 23 July 2013 and 21 September 2013, respectively.

In Exp 2, none of the daily mean values of thermal comfort indices assessed were above the critical ones: 68 for THI and 25°C for CCI. However, during the 10 days of TT collection, a total of 18.33% of hours (44 h) had CCI values $\geq 25^{\circ}\text{C}$. In addition, in those days where RR were collected (13 to 15 February), the number of hours with CCI $\geq 25^{\circ}\text{C}$ had a mean of 9.3 h/day, concentrated between 0900 and 1800 h. Respiration rates were greater at the third day of collection ($P=0.005$). In addition, RR increased after midday from 49.8 to 70.3 ± 2.5 bpm ($P<0.0001$), whereas TT reached mean values of $37.90^{\circ}\text{C} \pm 0.04^{\circ}\text{C}$ and $38.45^{\circ}\text{C} \pm 0.02^{\circ}\text{C}$, AM and PM respectively ($P<0.0001$). In fact, the CCI values of these hours in days where RR were collected, averaged 30.6°C ; 31.05°C ; and 30.2°C , respectively. These uncomfortable thermal conditions were reflected in an increase of RR in the afternoon measurements (Figure 4), but during the mornings no differences were observed across days of evaluation. Finally, CCI showed a great fluctuation during the day, as




Figure 3 Least squares mean values (standard error of the mean) for the average daily gain of steers fed with two levels of metabolizable energy and by season (MEI=1.85 or 2.72 times metabolizable energy intake above maintenance requirements).

shown in Figure 5, with one mean daily amplitude of 25.6°C between the maximum and minimum CCI.

The first two components (Figure 6) explain 80.0% of the observed variance. Adding a third component improves it to an 88.7%. The first component separates clearly the summer and winter experiments and is positively affected by CCW, dressing, pH, RH, PP and WS variables, whereas is negatively affected by the SR, AT, RA, TT variables. On the other hand, the second component is positively affected by BF and FBW variables, which in turns are highly correlated. Steers fed during the summer had higher RA and TT. On the contrary, the steers finished during winter showed a higher pH, which makes sense because they were exposed to more rainfall, WS, RH and less AT (environmental stressors).

Discussion

Most of the previous studies, regarding environmental effects, have been focused on heat stress in dairy and beef cattle including: performance (Mader and Davis, 2004,




Figure 4 Least squares mean values (standard error of the mean) for respiration rates (bpm = breaths per minute) per day and moment of the day (grey columns = before noon and black columns = afternoon) of steers fed with two levels of metabolizable energy intake above maintenance requirements. Different letters in the bar represent statistical differences (Tukey's test, $P<0.05$).

Table 3 Least square means (standard error of the mean) of carcasses variables of finished steers fed with two levels of metabolizable energy intake above maintenance requirements and by season

Variable*	Winter		Summer		SEM	P-value		
	1.85 MEI	2.72 MEI	1.85 MEI	2.72 MEI		Season	Energy	Interaction
Hot carcass weight (kg)	293.3	313.6	280.5	270.4	4.72	<0.0001	0.286	0.003
Cold carcass weight (kg)	285.6	306.3	274.1	263.1	4.65	<0.0001	0.307	0.002
Back fat (mm)	2.9	4.5	3.9	3.9	0.65	0.7553	0.256	0.232
Fat coverage† (0 to 4)	1.0	1.0	1.7	1.6	0.10	<0.0001	0.694	0.694
KPH (1 a 5)	2.1	2.0	1.1	1.2	0.14	<0.0001	1.000	0.557
pH	5.87	5.82	5.57	5.52	0.04	<0.0001	0.242	0.953
Rib eye area (cm ²)	45.9	45.9	98.0	96.2	3.50	<0.0001	0.789	0.806
Dressing (%)	57.4	57.7	52.4	50.6	0.006	<0.0001	0.221	0.105

MEI = metabolizable energy intake above energy maintenance requirements; SEM = standard error of the mean; KPH = kidney–pelvic–heart fat.

*Back fat, pH, and rib eye area were measured between the 12th/13th in the winter, and between 9th/10th rib in the summer. This change was done by slaughter plant personnel.
†Adipose tissue covering the outer face of the carcass (from 0 = no adipose tissue to 3 = abundant adipose tissue, without being excessive and its distribution) Official Chilean Standard NCh 1306, of 2002.




Figure 5 Hourly least square mean values of the comprehensive climate index (CCI) and the tympanic temperature (TT) of steers fed with two levels of metabolizable energy intake above maintenance requirements for each day of respiration rate data collection of the summer experiment.




Figure 6 Bi-plot (score plot and the loading plot) of the two first principal components of steers finished with two levels of metabolizable energy intake during the summer and winter seasons. SR=solar radiation (W/m^2); TT=tympanic temperature ($^{\circ}C$); RA=Rib eye area (cm^2); FBW=final live BW (kg); BF=back fat (mm); CCW=cold carcass weight (kg); WS=wind speed (m/s); dressing=dressing percentage; and pH=degree of acidity of the beef carcass (●=summer 2.72 times metabolizable energy intake above maintenance requirements (MEI); ○=summer 1.85 times MEI; ■=winter 2.72 times MEI; □=winter 1.85 times MEI).

Bernabucci *et al.*, 2014), animal metabolism (Baumgard and Rhoads, 2013), fertility (Ferreira *et al.*, 2011), gene expression (Rhoads *et al.*, 2011, Howard *et al.*, 2014) and nutrition (Mader and Davis, 2002, Smith *et al.*, 2006) among others.

On the contrary, cold stress has received less attention (Young *et al.*, 1989), with an emphasis on the lower critical temperature and its impact on the same variables before mentioned, probably, because cattle naturally has a greater level of adaptation to it.

In many regions of North America (Canada and United States) muddy conditions are limited to 4 months, mainly due to snow melting and ground thawing. However, rainfall seems to be less important (Grandin, 2016). On the contrary, in regions with humid-temperate climate, winter season is characterized by abundant rainfall (up to 800 mm/year), creating important muddy conditions as well as almost a lingering wet hair coat condition. In fact, at the experimental site location rainfall season can extent to at least 5 to 7 months. Indeed, during the last 35 years there was an average of 134 days with rainfall and an accumulated precipitation of 1010 mm/year. Thus, mud depth and mud in animal's body depends on the amount of precipitations and cattle management (density, BW, soil properties). Few studies have been conducted in humid-temperate regions that compare the effects of environmental factors on animal welfare and animal performance. However, there exists an increasing concern by producers and government agencies about these topics, which in turn may also be associated with climate change.

The differences in TT observed between seasons seem not to be biologically important. In addition, they are in agreement with the results previously reported by Arias *et al.* (2011), who reported a greater TT during the summer without differences due to the level of energy intake. In addition, they also reported a difference in TT due to energy intake during the wintertime, coinciding with our results. These differences can be explained due to a higher amount of direct

solar radiation being received by the animals in the summer season (Arias, 2008), but also because different genes have been associated with body temperature regulation during winter and summer seasons (Howard *et al.*, 2014). In fact, it has been established that animal performance varies between breeds (Pesonen *et al.*, 2012). This could be explained in part by the different energy requirements for maintenance as well as per the composition of the gain (NRC, 1981; NASEM, 2016). As consequence, the present results may differ according to the genotype. During the winter experiment, unlike that observed in the summer, mean values of TT dropped sharply after the second day of data collection. This could be explained by changes in the climatic conditions with an increase of both precipitations and wind speed (Table 2). Thus, PP increased almost 10 times (from <2 to almost 20 mm/day) on days three and four; whereas WS increased 1.6 and 2.8 times, respectively. Similarly, on days 6 and 7 there was also an increase in PP, even greater than days three and four, but the increase in WS was moderate. The previous conditions favoured mud formation which in turns affect animal welfare and animal performance by increasing energy of maintenance requirements and feed intake (Mader and Gaughan, 2011). In addition, mud is of great concern regarding contamination of the carcass (Garcia *et al.*, 2008). Thus, the combination of WS and PP have an important role in the thermal balance during the wintertime (Mader and Griffin, 2015), even though the values of AT were above 0°C, as such pointed out by Brownson and Ames (1980), who indicate that a steer may experience cold stress at 15.6°C if its hair coat is wet. The relevance of WS was also discussed by Angrecka and Herbut (2015) in a study of dairy cattle in a free stall barn. Thus, changes in the winter weather conditions implies adjustments in cattle behaviour (Graunke *et al.*, 2011).

Energy intake has been also reported as a factor affecting carcass quality (McCarthy *et al.*, 1985; Sami *et al.*, 2006), with animals fed with higher energy diets showing greater hot carcass weight and better quality grades (Berthiaume *et al.*, 2006). In addition, other researchers have reported changes on digestibility of organic matter and ether extract as well as changes in blood metabolite concentrations (Fiems *et al.*, 2007). There is also an effect of season of the year on percentage choice cattle (Birkelo and Johnson, 1993).

During the summer experiment, the mean AT for the week of TT collection was 16.6°C. However, 50% of those days presented maximum AT above 25.0°C, averaging 28.5°C. Nevertheless, during night time the AT averaged 7.5°C, that is a drop of 21.04°C. This great difference in AT between day and night allowed for the dissipation of heat load accumulated during the day. In addition, none of the daily mean values of thermal comfort indices exceeded the thresholds established in the literature, that is, THI = 68 (Zimbelman *et al.*, 2009) and CCI = 25.0°C (Mader *et al.*, 2010). Nevertheless, daily maximum values for those days in THI, THI_{adj} and CCI were 75.5, 79.9 and 34.2°C, respectively. The previous was in agreement with the signs of thermal discomfort showed by the animals (increase in RR), that showed an

average of 20 breaths per minute more during the afternoon when compared with the morning. Similarly, Gaughan and Mader (2014) in a study conducted in Queensland Australia, reported greater RR and panting scores during the afternoon, which in turns resulted in higher body temperature. In addition, these authors concluded that there is a close relationship between RR and body temperature. This discomfort observed in the animals (higher RR), could be associated with the largest solar radiation received between 0900 h and 1800 h that averaged $683.2 \pm 26.8 \text{ W/m}^2$. In contrast, the rest of the hours of the day averaged $142.0 \pm 16.4 \text{ W/m}^2$.

On the other hand, ADG observed in the summer season were in line with those projected by the NRC Beef model (National Academies of Sciences Engineering and Medicine, 2016), but were lower than projected under thermoneutral conditions by the model in the winter time, where the expected ADG were 0.4 and 1.07 kg/day for treatments 1.85 times and 2.72 times MEI, respectively. However, observed ADG was reduced in 50.5% and 42.5% each. The combination of low temperatures, hair wet coat, mud in the body and mud depth in the pen could explain this difference in ADG (Mader, 2011). Rayburn and Fox (1990), indicated that the ADG decreased more as the AT continues decreasing under 20°C. In addition, they point out that there exists a great sensibility in ADG due to WS and hair wet coat. For example, animals with hair wet coat decreased ADG in 8.2% and 46.4% when AT dropped from 20°C to 10°C and 0°C, respectively. Similarly, animals with wet hair coats and exposed to WS of 1.78 m/s showed a decrease of 44.3% in ADG. In the same context, Morrison *et al.* (1970) reported that ADG decreased by 14.69% when cattle were exposed to 10 min/h of artificial rainfall and also a 30.17% due to muddy conditions in the pen. In our study, mean AT and WS for all the feeding period of the winter experiment were $8.09^\circ \text{C} \pm 0.09^\circ \text{C}$ and $2.02 \pm 0.03 \text{ m/s}$, whereas there were only 44 days without PP. Thus, steers had its hair coat wet at least 69.7% of days of feeding. Finally, using a similar methodology to assess mud in the animals' body, Honeyman *et al.* (2012) found no difference in mud scores between winter and summer seasons. We did not record mud depth in the summer experiment.

In the winter study, the first measurement of mud depth was done 41 days after steers occupied the pens. Accumulated rainfall to that date was 210.4 mm, whereas the second measurement was done after 102 days of pen occupation with another 278.1 mm of accumulated rainfall. During the last 45 days, accumulated precipitation declined to 40.2 mm. Thus, mean rainfall was (5.13; 4.71; and 0.89) mm/day for each period. The accumulation of mud negatively affected animal performance, achieving lower ADG, when compared summer v. winter. On the contrary, summer performance was 3.1 times better than winter, whereas steers fed with higher MEI diet had 1.6 times better performance than steers fed with a low MEI level. The last value is in accordance with the differences in MEI provided between treatments which was a ratio of 1.5 times. This demonstrates the strong effect of environmental conditions on animal performance.

From a productive and economic point of view, the winter period proved to be the most critical. There was a great inefficiency during the winter period, as F:G ratio was 3.8 times greater in the winter than the summer. That is, steers consumed 28.8 kg of DM to produce one kg of BW, whereas in the summer they needed only 7.6 kg of DM. These differences increased when the energy level of the diet was considered. Thus, during the winter, steers fed with 1.85 times the MEI consumed 39.6 kg of DM for each kg of BW gained. Meanwhile, those fed 2.72 times MEI required 18.0 kg DM (2.2 times less feed). However, in the summer experiment both values were similar, requiring 7.14 and 8.01 kg of DM for each kg of BW gained for 1.85 and 2.72 times the MEI, respectively.

The MEI had a direct effect on the TT and ADG of steers finished during the winter period in an open feedlot. In addition, both variables were directly affected by weather conditions, particularly WS and PP (rainfall), as well as feed conversion. On the contrary, during summer period MEI level fed to steers did not affect TT neither ADG.

Acknowledgements

This work was supported by a grant from the Chilean National Fund for Science and Technology (FONDECYT, Project 11121320; Santiago, Chile).

Declaration of interest

The authors of the manuscript declare that there is no conflict of interest including financial, non-financial and relationships.

Ethics statement

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional guides on the care and use of farm animals.

Software and data repository resources

The datasets generated in the current study are available from the corresponding author on reasonable request.

References

Angrecka S and Herbut P 2015. Conditions for cold stress development in dairy cattle kept in free stall barn during severe frosts. *Czech Journal of Animal Science* 2, 81–87.

Arias RA 2008. Modeling the effects of environmental factors on finished cattle. PhD thesis, University of Nebraska, Lincoln, NE, USA.

Arias RA, Mader TL and Parkhurst AM 2011. Effects of diet type and metabolizable energy intake on tympanic temperature of steers fed during summer and winter seasons. *Journal of Animal Science* 89, 1574–1580.

Baumgard L and Rhoads RP 2013. Effects of heat stress on postabsorptive metabolism and energetics. *Annual Review of Animal Biosciences* 1, 311–337.

Bernabucci U, Biffani S, Buggiotti L, Vitali A, Lacetera N and Nardone A 2014. The effects of heat stress in Italian Holstein dairy cattle. *Journal of Dairy Science* 97, 471–486.

Berthiaume R, Mandell I, Faucitano L and Lafreniere C 2006. Comparison of alternative beef production systems based on forage finishing or grain-forage diets with or without growth promotants: 1. Feedlot performance, carcass quality, and production costs. *Journal of Animal Science* 84, 2168–2177.

Birkelo CP and Johnson DE 1993. Seasonal environment, performance and energy metabolism of feedlot cattle in northern Colorado (USA). In *Livestock Environment IV International Symposium* (ed. E Collins and C Boon), pp. 1117–1124. ASAE, University of Warwick, Coventry, UK.

Birkelo CP, Johnson DE and Phetteplace HP 1991. Maintenance requirements of beef cattle as affected by season on different planes of nutrition. *Journal of Animal Science* 69, 1214–1222.

Brownson R and Ames D 1980. Winter stress in beef cattle. In *Great Plains beef cattle handbook cooperative extension service*, Great Plains States, USA.

Dijkman J and Lawrence P 1997. The energy expenditure of cattle and buffaloes walking and working in different soil conditions. *The Journal of Agricultural Science* 128, 95–103.

Ferreira RM, Ayres H, Chiaratti MR, Ferraz ML, Araujo AB, Rodrigues CA, Watanabe YF, Vireque AA, Joaquim DC, Smith LC, Meirelles FV and Baruselli PS 2011. The low fertility of repeat-breeder cows during summer heat stress is related to a low oocyte competence to develop into blastocysts. *Journal of Dairy Science* 94, 2383–2392.

Fiems LO, Vanacker JM, De Boever JL, van Caelenbergh W, Aerts JM and De Brabander DL 2007. Effect of energy restriction and re-alimentation in Belgian Blue double-musled beef cows on digestibility and metabolites. *Journal of Animal Physiology and Animal Nutrition* 91, 54–61.

Garcia LG, Nicholson KL, Hoffman TW, Lawrence TE, Hale DS, Griffin DB, Savell JW, VanOverbeke DL, Morgan JB, Belk KE, Field TG, Scanga JA, Tatum JD and Smith GC 2008. National beef quality audit-2005: survey of targeted cattle and carcass characteristics related to quality, quantity, and value of fed steers and heifers. *Journal of Animal Science* 86, 3533–3543.

Gaughan JB and Mader TL 2014. Body temperature and respiratory dynamics in un-shaded beef cattle. *International Journal of Biometeorology* 58, 1443–1450.

Grandin T 2016. Evaluation of the welfare of cattle housed in outdoor feedlot pens. *Veterinary and Animal Science* 1–2, 23–28.

Graunke KL, Schuster T and Lidfors LM 2011. Influence of weather on the behaviour of outdoor-wintered beef cattle in Scandinavia. *Livestock Science* 136, 247–255.

Hahn GL, Gaughan JB, Mader TL and Eigenberg RA 2009. Thermal indices and their applications for livestock environments. In *Livestock energetics and thermal environmental management* (ed. JA DeShazer), pp. 113–130. ASABE, St. Joseph, MI, USA.

Honeyman MS, Maxwell D and Busby WD 2012. Effects of stocking density on steer performance and carcass characteristics in bedded hoop barns. *Animal Industry Report*, p. 4. Iowa, USA.

Howard JT, Kachman SD, Snelling WM, Pollak EJ, Ciobanu DC, Kuehn LA and Spangler ML 2014. Beef cattle body temperature during climatic stress: a genome-wide association study. *International Journal of Biometeorology* 58, 1665–1672.

INN 2002. Bovine carcasses – definitions and grading. In (ed. INd Normalización), p. 10. INN, Santiago, Chile.

Mader TL 2011. Mud effects on feedlot cattle. *Nebraska Beef Report*, pp. 82–83. University of Nebraska-Lincoln, USA.

Mader TL and Davis MS 2002. Wind speed and solar radiation corrections for the temperature–humidity index. In *Proceedings of the 15th Conference on Biometeorology and Aerobiology joint with the 16th International Congress on Biometeorology*, Boston, MA, USA, pp. 154–157.

Mader TL and Davis MS 2004. Effect of management strategies on reducing heat stress of feedlot cattle: feed and water intake. *Journal of Animal Science* 82, 3077–3087.

Mader TL and Gaughan JB 2011. Effects of climate variability on domestic livestock. In *Handbook on climate change and agriculture* (ed. ADAR and Mendelsohn), pp. 32–48. Edward Elgar Publishing Limited, MA, USA.

Mader TL, Davis MS and Brown-Brandl T 2006. Environmental factors influencing heat stress in feedlot cattle. *Journal of Animal Science* 84, 712–719.

Mader TL and Griffin D 2015. Management of cattle exposed to adverse environmental conditions. *Veterinary Clinics of North America: Food Animal Practice* 31, 247–258.

Mader TL, Johnson LJ and Gaughan JB 2010. A comprehensive index for assessing environmental stress in animals. *Journal of Animal Science* 88, 2153–2165.

McCarthy FD, Hawkins DR and Bergen WG 1985. Dietary energy density and frame size effects on composition of gain in feedlot cattle. *Journal of Animal Science* 60, 781–790.

- Morrison SR, Givens RL, Garrett WN and Bond TE 1970. Effects of mud-wind-rain on beef cattle performance in feed lot. *California Agriculture* 24, 6–7.
- National Academies of Sciences Engineering and Medicine 2016. Nutrient requirements of beef cattle, 8th revised edition. The National Academies Press, Washington, DC, USA.
- National Research Council 1981. Effect of environment on nutrient requirements of domestic animals. National Academy Press, Washington, DC, USA.
- Pesonen M, Honkavaara M and Huuskonen AK 2012. Effect of breed on production, carcass traits and meat quality of Aberdeen Angus, Limousin and Aberdeen Angus × Limousin bulls offered a grass silage-grain-based diet. *Agricultural and Food Science* 361–369.
- Rayburn EB and Fox DG 1990. Predicting growth and performance of Holstein steers. *Journal of Animal Science* 68, 788–798.
- Rhoads RP, La Noce AJ, Wheelock JB and Baumgard LH 2011. Short communication: alterations in expression of gluconeogenic genes during heat stress and exogenous bovine somatotropin administration. *Journal of Dairy Science* 94, 1917–1921.
- Sami AS, Koegel J and Eichinger H 2006. Effects of the dietary energy source on meat quality and eating quality attributes and fatty acid profile of Simmental bulls. *Animal Research* 55, 287–299.
- Smith TR, Chapa A, Willard S, Hemdon C Jr., Williams RJ, Crouch J, Riley T and Pogue D 2006. Evaporative tunnel cooling of dairy cows in the southeast. II: impact on lactation performance. *Journal of Dairy Science* 89, 3915–3923.
- Young BA, Walker B, Dixon AE and Walker VA 1989. Physiological adaptation to the environment. *Journal of Animal Science* 67, 2426–2432.
- Zimbelman RB, Rhoads RP, Rhoads ML, Duff GC, Baumgard LH and Collier JL 2009. A re-evaluation of the impact of temperature humidity index (THI) and black globe humidity index (BGHI) on milk production in high producing dairy cows. In *Proceedings of the 24th Southwest Nutrition and Management Conference*, Tempe, AZ, USA, pp. 158–169.