

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA Wildlife Services - Staff Publications

U.S. Department of Agriculture: Animal and
Plant Health Inspection Service

2012

Detection of prion protein in the cerebrospinal fluid of elk (*Cervus canadensis nelsoni*) with chronic wasting disease using protein misfolding cyclic amplification

Tracy A. Nichols

U.S. Department of Agriculture, tracy.a.nichols@aphis.usda.gov

Terry R. Spraker

Colorado State University - Fort Collins, terry.spraker@colostate.edu

Tom Gidlewski

National Wildlife Disease Program, Thomas.Gidlewski@aphis.usda.gov

Jenny G. Powers

National Park Service

Glenn C. Telling

Colorado State University - Fort Collins, glenn.telling@colostate.edu

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc

Nichols, Tracy A.; Spraker, Terry R.; Gidlewski, Tom; Powers, Jenny G.; Telling, Glenn C.; VerCauteren, Kurt C.; and Zabel, Mark D., "Detection of prion protein in the cerebrospinal fluid of elk (*Cervus canadensis nelsoni*) with chronic wasting disease using protein misfolding cyclic amplification" (2012). *USDA Wildlife Services - Staff Publications*. 1172.

https://digitalcommons.unl.edu/icwdm_usdanwrc/1172

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA Wildlife Services - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Tracy A. Nichols, Terry R. Spraker, Tom Gidlewski, Jenny G. Powers, Glenn C. Telling, Kurt C. VerCauteren, and Mark D. Zabel

Detection of prion protein in the cerebrospinal fluid of elk (*Cervus canadensis nelsoni*) with chronic wasting disease using protein misfolding cyclic amplification

Journal of Veterinary Diagnostic Investigation
24(4) 746–749
© 2012 The Author(s)
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1040638712448060
<http://jvdi.sagepub.com>

Tracy A. Nichols,¹ Terry R. Spraker, Tom Gidlewski, Jenny G. Powers,
Glenn C. Telling, Kurt C. VerCauteren, Mark D. Zabel

Abstract. Cerebrospinal fluid (CSF) has been examined as a possible source for preclinical diagnosis of prion diseases in hamsters and sheep. The present report describes the detection of chronic wasting disease (CWD) in the CSF of elk and evaluates its usefulness as an antemortem test for CWD. The CSF from 6 captive and 31 free-ranging adult elk was collected at necropsy and evaluated for the presence of the abnormal isoform of the prion protein that has been associated with CWD (PrP^{CWD}) via protein misfolding cyclic amplification. Additionally, the obex from each animal was examined by immunohistochemistry (IHC). Four out of 6 captive animals were CWD-positive and euthanized due to signs of terminal CWD. The remaining 2 were CWD negative. None of the 31 free-range animals showed overt signs of CWD, but 12 out of 31 tested positive for CWD by IHC. Protein misfolding cyclic amplification detected PrP^{CWD} from 3 of the 4 captive animals showing clinical signs of CWD and none of the nonclinical animals that were CWD positive by IHC. The data suggests that CWD prions can be detected in the CSF of elk, but only relatively late in the course of the disease.

Key words: Cerebrospinal fluid; *Cervus canadensis nelsoni*; chronic wasting disease; protein misfolding cyclic amplification; prion protein; Rocky Mountain elk.

Chronic wasting disease (CWD) is a fatal, infectious neurodegenerative disease that affects wild and captive cervids. It is a transmissible spongiform encephalopathy, or prion disease. Development of an antemortem test for detecting the misfolded prion protein associated with CWD (PrP^{CWD}) in nonclinical animals would be useful for wildlife and captive population management strategies. To date, preclinical testing for PrP^{CWD} utilizes immunohistochemistry (IHC) of the palatine tonsils or rectal lymphoid tissues in cervids.^{18,21,24} However, IHC does not routinely detect very early cases of CWD in these tissues.^{17,20} Various fluids, such as saliva, blood, urine, and cerebrospinal fluid (CSF) have been evaluated in animals for their use in preclinical antemortem detection of abnormal prions.^{4,6,10,22} Due to the minute concentrations of infectious proteins in saliva, blood, and urine, detection of the misfolded infectious prion protein (PrP^{res}) has been limited or unsuccessful without the use of protein misfolding cyclic amplification (PMCA), which increases the concentration to a detectable level, and is orders of magnitude more sensitive for detecting PrP^{CWD} than commonly used methods such as Western blotting and IHC,⁷ enabling detection of minute amounts of infectious prions in animal tissues and environmental samples.⁹

In the current study, the usefulness of PMCA to detect PrP^{CWD} in CSF from wild and captive elk (*Cervus canadensis*

nelsoni) was evaluated as a potential tool for early detection of PrP^{CWD}. The PMCA in vitro technique amplifies minute amounts of infectious prions by rapidly converting normal cellular prion protein (PrP^c) to PrP^{res}.^{11,16} Cerebrospinal fluid (5–15 ml) was collected at necropsy from 6 captive and 31 free-ranging, adult elk. Four of the 6 captive elk were euthanized due to positive CWD rectal biopsy results and terminal clinical signs. Cerebrospinal fluid was collected during the postmortem examination just prior to removal of the head. The head was flexed dorsally, the ventral muscles severed, the ventral aspect of the foramen magnum was exposed, and CSF collected with an 18-gauge needle attached to a 12-cc syringe. Great care was taken to ensure that the extraction site

From the U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services, National Wildlife Research Center (Nichols, VerCauteren) and National Wildlife Disease Program (Gidlewski), Fort Collins, CO; Colorado State University Diagnostic Laboratory (Spraker) and the Prion Research Program (Spraker, Telling, Zabel), Department of Microbiology, Immunology and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO; and the National Park Service, Biological Resources Management Division, Fort Collins, CO (Powers).

¹Corresponding Author: Tracy A. Nichols, National Wildlife Research Center, U.S. Department of Agriculture, 4101 La Porte Avenue, Fort Collins, CO 80521. tracy.a.nichols@aphis.usda.gov

of the CSF was clean and without blood contamination. Cerebrospinal fluid was frozen at -80°C until samples were analyzed using PMCA.

The CWD status of each elk was determined by examination of the obex at the level of the dorsal motor nucleus of the vagus nerve by using previously described IHC techniques.¹⁸ The obex from all CWD-positive elk was scored as previously reported.¹⁹ Briefly, nuclear regions and axonal tracts were evaluated and scored 0–10 for the severity of spongiform degeneration and accumulation of PrP^{CWD} immunoreactivity (IR). Tissue samples showing no detectable IR received a score of zero, and samples showing heavy immunoreactivity in all nuclei and axons received a score of 10.

Brain homogenate from transgenic Tg5037 mice, which express a 5-fold increase in elk PrP^{C} , were used as a source of PrP^{C} substrate in the PMCA reactions.² A 10% (wt/vol) normal brain homogenate (NBH) was prepared as previously described, utilizing a homogenizer^a at setting 6 for 5 min.⁹ Equivalent volumes of CSF and NBH (25 μl of each) were added to individual 200- μl PCR tubes sealed with plastic paraffin film with a paper backing and sonicated^b with 40-sec pulses at 37°C , power setting 7, every 30 min for 24 hr. After each 24-hr round, 25 μl of each sample was mixed with 25 μl of fresh NBH and transferred to a fresh tube. A positive amplification control of 1:100,000 CWD-positive brain homogenate was also amplified with each sample group as were NBH-negative controls. Duplicate samples were amplified in 3 independent replicate experiments for a total of 6 samples. Upon completion of 6 rounds, amplified samples were digested with proteinase K^c and immunoblotted as previously described.⁹ The proteinase K-resistant core of PrP^{res} appears on Western blots as bands shifted below those seen for full-length PrP . Only samples in which signal was detected in 2 of the 3 replicate experiments were considered positive.

The misfolded prion protein associated with CWD was detected in 3 out of 4 captive, IHC-positive clinical elk (Fig. 1, Table 1). However, the nonclinical, wild CWD IHC-positive elk did not yield detectable CWD in the CSF (Table 1). These animals were in generally good health, as indicated by body and coat condition.

Although it is unknown how long the 3 PMCA-positive elk had been infected, their obex scores ranged from 9 to 10 (Table 1), whereas the obex scores of 12 IHC-positive but PMCA-negative nonclinical elk ranged from 2 to 9. Negative captive elk had no detectable PrP^{CWD} and therefore had scores of 0. Previous studies involving sheep and hamsters demonstrated that PrP^{res} could be detected in the CSF.^{3,10} However, the present findings are not readily compared to those previous results, which were derived exclusively from terminal, clinical animals.^{3,10} Transport of prions to the brain from distal regions such as the gut has been shown to occur via movement along peripheral nerves and via the blood.¹⁴ Studies conducted in naturally and orally exposed sheep to scrapie suggest that prions migrate from the gut-associated




Figure 1. Western blots of captive elk cerebrospinal fluid after 6 rounds of protein misfolding cyclic amplification. NBH = normal brain homogenate negative controls. Positive control 1:100,000 dilution of chronic wasting disease-positive brain homogenate. Chronic wasting disease was detected in captive elk 1–3. Representative of 3 replicates.

Table 1. Source, and protein misfolding cyclic amplification (PMCA) and immunohistochemistry (IHC) results for 6 captive and 31 free-ranging adult elk (*Cervus canadensis nelsoni*).

Elk ID	Source	PMCA results	IHC obex score
1	Captive	–	0
2	Captive	–	0
1	Captive	+	9–10
2	Captive	+	9–10
3	Captive	+	9–10
4	Captive	–	9–10
1	Free range	–	0
2	Free range	–	9
3	Free range	–	4
4	Free range	–	8
5	Free range	–	7
6	Free range	–	4
7	Free range	–	7
8	Free range	–	7
9	Free range	–	2
10	Free range	–	2
11	Free range	–	6
12	Free range	–	5
13	Free range	–	6
14–31	Free range	–	0

lymph tissue to the central nervous system along nerve tracts to the spinal cord and brain.¹ A similar pattern of PrP^{CWD} deposition has been observed in deer.^{8,12,13,23} Chronic wasting

disease has been detected in blood,⁶ and studies done in sheep demonstrate that intravenous inoculation with scrapie results in neuroinvasion and disease indistinguishable from other routes of infection.¹⁵ It has been postulated in the literature that the circumventricular organs (CVO) in the brain could be a site of entry for prions from the blood into the brain.¹⁵ The CVO are specialized structures that line the third and fourth ventricles of the brain and are considered to be absent of the “blood-brain barrier.”⁵ As has been seen with scrapie in sheep,¹⁵ PrP^{CWD} can be detected in the CVO, with the exception of the choroid plexus, in nonclinical deer and elk with obex scores greater than 9 (with the obex scoring system, clinical signs are usually not observed until obex scores of 9 or 10 are achieved (unpublished data, 2005). Early accumulation of scrapie in the CVO has prompted some researchers to propose that because the CVO is devoid of the blood-brain barrier, prions can cross unhampered from the blood to the brain and then on to regional lymph nodes.¹⁵ The data suggests that in the earlier stages of disease, the intact ependymal and CVO appear to sequester PrP^{CWD} and prevent infiltration of PrP^{CWD} into the CSF. It is therefore reasonable to propose that in the latter stages of disease, with obex scores greater than 9, the PrP^{CWD} begins to leak through the ependyma, allowing PrP^{CWD} access to the CSF. Leakage is not thought to be caused by autolysis at necropsy, as the postmortem interval of the positive wild elk was 3–4 hr and only 1 hr for the captive elk. In conclusion, the present data suggests that PrP^{CWD} infiltrates the CSF at relatively late stages of CWD and that PMCA of CSF is not useful as a diagnostic tool in nonclinical elk. Work needs to continue to investigate other tissues such as saliva and lymph node biopsies as PMCA antemortem diagnostic specimens.

Acknowledgements

The authors thank the North American Deer Farmers Association for research support.

Sources and manufacturers

- a. Bullet Blender[®] homogenizer, Next Advance Inc., Averill Park, NY.
- b. Model 3000MP, Misonix Inc., Farmingdale, NY.
- c. Roche, Madison, WI.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Andréoletti O, Berthon P, Marc D, et al.: 2000, Early accumulation of PrP^(Sc) in gut-associated lymphoid and nervous tissues

- of susceptible sheep from a Romanov flock with natural scrapie. *J Gen Virol* 81:3115–3126.
2. Angers RC, Seward TS, Napier D, et al.: 2009, Chronic wasting disease prions in elk antler velvet. *Emerg Infect Dis* 15:696–703.
3. Atarashi R, Moore RA, Sim VL, et al.: 2007, Ultrasensitive detection of scrapie prion protein using seeded conversion of recombinant prion protein. *Nat Methods* 4:645–650.
4. Haley NJ, Seelig DM, Zabel MD, et al.: 2009, Detection of CWD prions in urine and saliva of deer by transgenic mouse bioassay. *PLoS One* 4:e4848.
5. Johnson AK, Gross PM: 1993, Sensory circumventricular organs and brain homeostatic pathways. *FASEB J* 7:678–686.
6. Mathiason CK, Powers JG, Dahmes SJ, et al.: 2006, Infectious prions in the saliva and blood of deer with chronic wasting disease. *Science* 314:133–136.
7. Meyerett C, Michel B, Pulford B, et al.: 2008, In vitro strain adaptation of CWD prions by serial protein misfolding cyclic amplification. *Virology* 382:267–276.
8. Miller MW, Williams ES: 2002, Detection of PrP^(CWD) in mule deer by immunohistochemistry of lymphoid tissues. *Vet Rec* 151:610–612.
9. Nichols TA, Pulford B, Wyckoff AC, et al.: 2009, Detection of protease-resistant cervid prion protein in water from a CWD-endemic area. *Prion* 3:171–183.
10. Orrú CD, Wilham JM, Hughson AG, et al.: 2009, Human variant Creutzfeldt-Jakob disease and sheep scrapie PrP^(res) detection using seeded conversion of recombinant prion protein. *Protein Eng Des Sel* 22:515–521.
11. Saá P, Castilla J, Soto C: 2005, Cyclic amplification of protein misfolding and aggregation. *Methods Mol Biol* 299:53–65.
12. Sigurdson CJ, Spraker TR, Miller MW, et al.: 2001, PrP^(CWD) in the myenteric plexus, vagosympathetic trunk and endocrine glands of deer with chronic wasting disease. *J Gen Virol* 82:2327–2334.
13. Sigurdson CJ, Williams ES, Miller MW, et al.: 1999, Oral transmission and early lymphoid tropism of chronic wasting disease PrP^(res) in mule deer fawns (*Odocoileus hemionus*). *J Gen Virol* 80:2757–2764.
14. Sisó S, González L, Jeffrey M: 2010, Neuroinvasion in prion diseases: the roles of ascending neural infection and blood dissemination. *Interdiscip Perspect Infect Dis* 2010:747892.
15. Sisó S, Jeffrey M, González L: 2009, Neuroinvasion in sheep transmissible spongiform encephalopathies: the role of the haematogenous route. *Neuropathol Appl Neurobiol* 35:232–246.
16. Soto C, Anderes L, Suardi S, et al.: 2005, Pre-symptomatic detection of prions by cyclic amplification of protein misfolding. *FEBS Lett* 579:638–642.
17. Spraker TR, Gidlewski TL, Balachandran A, et al.: 2006, Detection of PrP^(CWD) in postmortem rectal lymphoid tissues in Rocky Mountain elk (*Cervus elaphus nelsoni*) infected with chronic wasting disease. *J Vet Diagn Invest* 18:553–557.
18. Spraker TR, O'Rourke KI, Balachandran A, et al.: 2002, Validation of monoclonal antibody F99/97.6.1 for immunohistochemical staining of brain and tonsil in mule deer (*Odocoileus*

- hemionus*) with chronic wasting disease. J Vet Diagn Invest 14:3–7.
19. Spraker TR, O'Rourke KI, Gidlewski T, et al.: 2010, Detection of the abnormal isoform of the prion protein associated with chronic wasting disease in the optic pathways of the brain and retina of Rocky Mountain elk (*Cervus elaphus nelsoni*). Vet Pathol 47:536–546.
 20. Spraker TR, VerCauteren KC, Gidlewski T, et al.: 2009, Antemortem detection of PrP^{CWD} in preclinical, ranch-raised Rocky Mountain elk (*Cervus elaphus nelsoni*) by biopsy of the rectal mucosa. J Vet Diagn Invest 21:15–24.
 21. Spraker TR, VerCauteren KC, Gidlewski TL, et al.: 2009, Impact of age and sex of Rocky Mountain elk (*Cervus elaphus nelsoni*) on follicle counts from rectal mucosal biopsies for preclinical detection of chronic wasting disease. J Vet Diagn Invest 21:868–870.
 22. Tamgüney G, Miller MW, Wolfe LL, et al.: 2009, Asymptomatic deer excrete infectious prions in faeces. Nature 461:529–532.
 23. Williams ES: 2005, Chronic wasting disease. Vet Pathol 42: 530–549.
 24. Wolfe LL, Spraker TR, Gonzalez L, et al.: 2007, PrP^{CWD} in rectal lymphoid tissue of deer (*Odocoileus* spp.). J Gen Virol 88:2078–2082.