

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Papers in Natural Resources

Natural Resources, School of

---

2011

## Soil Moisture: A Central and Unifying Theme in Physical Geography

Rezaul Mahmood

*University of Nebraska - Lincoln*

Follow this and additional works at: <https://digitalcommons.unl.edu/natrespapers>



Part of the [Natural Resources and Conservation Commons](#), [Natural Resources Management and Policy Commons](#), and the [Other Environmental Sciences Commons](#)

---

Mahmood, Rezaul, "Soil Moisture: A Central and Unifying Theme in Physical Geography" (2011). *Papers in Natural Resources*. 1258.

<https://digitalcommons.unl.edu/natrespapers/1258>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/260592274>

# Soil Moisture: A central and unifying theme in Physical Geography

Article in *Progress in Physical Geography* · February 2010

DOI: 10.1177/0309133310386514

---

CITATIONS

165

READS

853

Some of the authors of this publication are also working on these related projects:



Desertification and dune activity along the U.S. - Mexico Border [View project](#)



The Holocene geology of South Texas [View project](#)



# Soil moisture: A central and unifying theme in physical geography

**David R. Legates**

University of Delaware, USA

**Rezaul Mahmood**

Western Kentucky University, USA

**Delphis F. Levia**

University of Delaware, USA

**Tracy L. DeLiberty**

University of Delaware, USA

**Steven M. Quiring**

Texas A&M University, USA

**Chris Houser**

Texas A&M University, USA

**Frederick E. Nelson**

University of Delaware, USA

## Abstract

Soil moisture is a critical component of the earth system and plays an integrative role among the various subfields of physical geography. This paper highlights not just how soil moisture affects atmospheric, geomorphic, hydrologic, and biologic processes but that it lies at the intersection of these areas of scientific inquiry. Soil moisture impacts earth surface processes in such a way that it creates an obvious synergistic relationship among the various subfields of physical geography. The dispersive and cohesive properties of soil moisture also make it an important variable in regional and microclimatic analyses, landscape denudation and change through weathering, runoff generation and partitioning, mass wasting, and sediment transport. Thus, this paper serves as a call to use research in soil moisture as an integrative and unifying theme in physical geography.

## Keywords

biogeography, climatology, geomorphology, hydrology, soil moisture

## 1 Introduction

Soil moisture or the available soil water content – the water contained in the root zone of the soil and available for plant utilization – is an

---

### Corresponding author:

David R. Legates, Department of Geography, 216 Pearson Hall, University of Delaware, Newark, DE 19716-2541, USA

Email: [legates@udel.edu](mailto:legates@udel.edu)

essential component of the hydrologic cycle (Mahmood, 1996). In the fundamental equation that describes the hydrologic cycle:

$$\begin{aligned} dS/dt = & (P_r + M) - (E + T) \\ & - (R_o + R_s + R_g) \end{aligned} \quad (1)$$

the time rate of change of soil moisture ( $dS/dt$ ) is balanced by the moisture input from rainfall ( $P_r$ ) and snowmelt ( $M$ ) and the moisture loss through soil evaporation ( $E$ ), plant transpiration ( $T$ ), overland runoff ( $R_o$ ) through either saturation or Hortonian flow, lateral subsurface flow ( $R_s$ ), and percolation to groundwater ( $R_g$ ). At the basic level, soil moisture content describes the temporal condition of water available to plants as well as providing an integrated assessment of the relative state of water supply versus water demand.


In addition to its obvious importance to hydrology, soil moisture is a variable whose importance extends to all aspects of physical geography. It provides the reservoir of water through which the fluxes of energy and moisture between the land surface and the atmosphere interact; it establishes the ground conditions upon which water movement from saturated, parched, or frozen soils can occur; and it supplies the necessary ingredients for the existence and development of plant life through both transport (lateral and vertical) and solubility. By its very nature, therefore, soil moisture serves to integrate physical geography. Figure 1 highlights these linkages and discusses impacts of various levels of soil moisture and climate, biogeography, and geomorphology.

Soil moisture is estimated by either in situ measurements, remote sensing techniques, or by atmospheric or hydrologic modeling. In the absence of a well-integrated program of sampling and scaling, the limited areal coverage over large land surface areas makes it difficult to use in situ measurements for studies more extensive than field-scale (Robinson et al., 2008; Robock et al., 2000; Verstraeten et al.,

2008; Wu and Li, 2009). By contrast, remote sensing of soil moisture can be advantageous owing to its continuous temporal and spatial coverage. A combined data assimilation approach with remotely sensed estimates and hydrologic modeling, calibrated with in situ measurements to quantify errors and uncertainties, is the most promising approach for soil moisture estimation involving large areas, although it too is likely to have deficiencies. A complete discussion of soil moisture measurements is beyond the scope of this topical review, but for a more thorough discussion of methods to measure soil moisture, the reader is encouraged to consult the following references. Observations – Francesca et al. (2010); Robinson et al. (2008); Vera et al. (2009); Vereecken et al. (2008). Models – Choi and Liang (2010); Fulakeza et al. (2002); Steiner et al. (2005). Remote sensing – Anderson and Croft (2009); Kerr (2007); Kidd et al. (2009); Schmugge et al. (2002); Tang et al. (2009); Wagner et al. (2007a, 2007b). Data assimilation – Balsamo et al. (2007); Dorigo et al. (2007); Ni-Meister (2008); Reichle et al. (2004).

## II Soil moisture and climate

In its most basic role, climatology (and climate science in general) seeks to describe the spatial and temporal fluxes of energy, moisture, and momentum between the atmosphere and the land surface, for which soil moisture plays an integral part (Thorntwaite, 1961; McCumber and Pielke, 1981; Chen and Avissar, 1994; Mahmood, 1996; Mahmood and Hubbard, 2002; 2004; 2007; LeMone et al., 2007). Water, in general, and soil moisture in particular are of fundamental importance to the ‘topoclimatology’ espoused by C.W. Thorntwaite (1953; 1961) and forms the basis of much of the sub-field of hydroclimatology (Mather, 1991). *Topoclimatology* was defined by Thorntwaite as the study of those interactions between climate and the earth’s surface that lead to geographical



**Figure 1.** Schematic showing simplified interactions between soil moisture, weather and climate, geomorphology and soils, and biogeography for high (upper left), normal (upper right), and low (lower left) soil moisture conditions. Bold arrows imply stronger linkage. In particular, soil moisture affects climate during dry and wet conditions (albedo change and partitioning between latent and sensible heat) while during normal conditions, climate variability drives fluctuations in soil moisture. Soil moisture is affected by fluvial processes during dry conditions and aeolian processes during wet conditions whereas geomorphology and soil type dictate the water-holding properties of the soil. Soil moisture affects biogeography during both wet (root rot and waterlogging), normal, and dry (wilting) conditions whereas the influence of biogeography on soil moisture is strongest during normal conditions. Interactions among the three subdisciplines also are shown.

variability of climate. Soil moisture is an integral component of the topoclimate.

Indeed, the concept of climatic classification has often focused on the interplay between moisture availability (precipitation) and moisture demand (potential evapotranspiration, with air temperature often used as a surrogate). It is no surprise then that moisture indices derived from climate classifications are reflective of the temporal characteristics of the soil moisture condition (Feddema, 2007; Mather, 1985; Thornthwaite, 1948; Willmott and Feddema,

1992). Indeed, the temporal variability of soil moisture in a given region is fundamental to the definition of its climate.

Far from being simply a resultant statistic, soil moisture itself is an active variable. It has been suggested that local evapotranspiration (i.e. latent heat flux) is the source of 10 to 30% of atmospheric water vapor (Brubaker et al., 1993). Obviously, soil moisture plays an important role in regulating the rate of the evapotranspiration. Moreover, Chen and Avissar (1994) noted that the development of clouds and

precipitation in a dry atmosphere is critically dependent on these local sources of moisture. Research has shown that changes to soil moisture modify the surface Bowen ratio (Rosenberg et al., 1983; Thom, 1972), the convective available potential energy (Clark and Arritt, 1995; Pielke, 2001), development of cloud (Ek and Holtslag, 2004; Findell and Eltahir, 2003), precipitation (Ookouchi et al., 1984; Pal and Eltahir, 2001; Pan et al., 1996; Quintanar et al., 2008; 2009), and the daytime evolution of the planetary boundary layer (Zhang and Anthes, 1982), particularly its wind field (Segal and Arritt, 1992). For example, Rabin et al. (1990) found convective clouds formed first over mesoscale-size areas of harvest wheat in Oklahoma with high sensible heat flux, in comparison to adjoining areas dominated by growing vegetation. Clouds were suppressed over relatively long bands downwind of small lakes and heavy tree cover represented by high latent heat flux. The more recent empirical and modeling study of Santanello et al. (2007) examined interactions between planetary boundary level height, initial atmospheric stability, and soil moisture using observed radiosonde data and a one-dimensional model. Both positive and negative feedbacks were identified that explained the land surface-atmospheric interactions; for example, a positive feedback resulted for dry soils with entrainment on soil drying, surface heating, and residual boundary layer growth.

Additional studies have focused specifically on the interactions of soil moisture and patterns of precipitation. Dong et al. (2007) noted a positive relationship between soil moisture and precipitation over grassland while Chang et al. (2009) found that antecedent soil moisture (wetter and warmer) further intensifies land-falling monsoonal low pressures. In a detailed regional modeling study, Koster et al. (2004) found areas of strong coupling, or 'hot spots', between soil moisture and precipitation. These so-called 'hot spots' are generally located in transition zones between wet and dry climates

such as those found in the central Great Plains of North America, the Sahel, equatorial Africa, and India. In these regions, potential evapotranspiration is consistently high while actual evapotranspiration is sensitive to soil moisture availability. Koster et al. (2004) further found that the response of the atmosphere to soil moisture is non-linear and not unidirectional and, moreover, wet soils can both increase and decrease the possibility of convection and precipitation, depending upon the initial state of the atmosphere (cf. Findell and Eltahir, 2003).

Owing to the lack of high-density networks over much of the globe, proper initiation of soil moisture in numerical models is often difficult. Studies that have reasonably prescribed soil conditions in atmospheric models have noted improvements in simulated forecasts (Dirmeyer, 2000; Douville et al., 2001; Hong and Kalnay, 2000; Huang et al., 1996; Schlosser and Milly, 2002). Dirmeyer (2000), in particular, noted a reduction in root-mean-square error in modeled near-surface temperature and improved rainfall patterns with reasonable specification of root zone soil moisture. Similarly, both Huang et al. (1996) and Schlosser and Milly (2002) found a strong correlation between subsurface water storage and modeled near-surface temperature and precipitation. Grasso (2000) has also noted an improvement in modeled forecasts with proper specification of both soil moisture and overlying vegetation.

Latent and sensible heat fluxes are affected by variations in soil moisture, both spatially and temporally, which alter the near-surface air temperature and humidity. This, in turn, can alter precipitation (Clark et al., 2004; Jones and Brunzell, 2009; Walker and Rowntree, 1977), temperature (Brown and Wax, 2007; De Laat and Maurellis, 2006), atmospheric circulation (Namias, 1959; Ookouchi et al., 1984; Pinty et al., 1989; Walker and Rowntree, 1977), and the climate in general (Rind, 1982; Shukla and Mintz, 1982; Yeh et al., 1984) which often accentuates and extends anomalous conditions

(Andreadis and Lettenmaier, 2006; DeLiberty and Legates, 2003, 2008; Entin et al., 2000; Koster and Suarez, 2003; Oglesby and Erickson, 1989; Rasmusson and Arkin, 1993). As a consequence, weather conditions may be affected by abnormal soil moisture conditions, provided the anomaly is of sufficient spatial and temporal size and intensity.

Because droughts are directly related to soil moisture conditions, much effort has been put into drought monitoring (e.g. Narasimhan and Srinivasan, 2005; Quiring, 2004), drought mitigation (Wilhite, 1997), and insurance protection (Abbaspour et al., 1992). However, flood events too can be exacerbated by high antecedent soil moisture conditions (Gamble and Meentemeyer, 1997; Hossain, 2006; Lacava et al., 2005). Gamble and Meentemeyer (1997) showed, for example, that high soil moisture with moderate rains often produced unseasonable floods in the southeastern United States. Indeed, the impact of too little as well as too much soil moisture can adversely affect crop yields such that accurate forecasts are needed to monitor the situation (e.g. Quiring and Legates, 2008). Soil moisture can be both a symptom of drought as well as a contributing factor. Although remote forcings such as sea-surface temperature (SST) anomalies in the Pacific and Atlantic Oceans are often the primary cause of North American droughts (e.g. McCabe et al., 2004, 2010; Quiring and Goodrich, 2008), Wu and Kinter (2009) distinguished the timescales of droughts and found the roles of SST forcing and local soil moisture differ significantly for long-term versus short-term droughts. Moreover, it has recently been demonstrated that soil moisture can play a significant role in modifying summer precipitation during the periods when SST anomalies are weak (Meng and Quiring, 2010). Local precipitation recycling can contribute to the intensity and duration of droughts and, therefore, an understanding of local land-atmosphere interactions (e.g. soil moisture) is a key to understanding and predicting the occurrence of droughts.

### **III Soil moisture in geomorphology and hydrology**

The dispersive and cohesive properties of soil moisture makes it an important variable in landscape denudation through weathering (e.g. Anderson, 2005; Richards and Kump, 2003; Stockwell et al., 2006), runoff generation and erosion (e.g. McDowell and Sharpley, 2002; McDowell et al., 2001; Seeger et al., 2004), and mass wasting (e.g. Acharya et al., 2009; Korup et al., 2004; Schuerch et al., 2006). It is also responsible for increasing soil moisture capacity through soil development (Phillips et al., 2008) and promoting the development of vegetation that can limit erosion (Harden and Scruggs, 2003). Specifically, the spatial and temporal variation in soil moisture in a drainage basin has implications for understanding environmental processes such as runoff generation and continuity as well as the erosion and sedimentation driven by overland flow. Overland runoff is generated through either the saturation of the soil to the surface (i.e. saturation overland flow) or through precipitation exceeding infiltration rates (i.e. Hortonian overland flow). The latter is the dominant mechanism in arid and semi-arid environments and depends on the precipitation rate relative to the infiltration capacity of the soil; it is therefore dependent on how much water is already in the soil. Infiltration capacity is greatest at the start of a storm and decreases rapidly until a constant infiltration rate is reached (Horton, 1933).

#### ***I Flow partitioning and mass wasting***

In humid-temperate regions, saturation overland flow is generated when the soil becomes totally saturated and is unable to receive any more water (Kirkby and Chorley, 1967). These variable source areas are responsible for storm runoff (e.g. Dunne and Black, 1970; Hewlett and Hibbert, 1967; Tsukamoto, 1963) and first develop where the water table is relatively

shallow. Regardless of the environment, antecedent moisture in the soil promotes the development of overland flow earlier in a precipitation event, meaning that a significant volume of water is contributed to the main channel network in a short time. In this respect, Vertessy and Elsenbeer (1999) argued that a poor understanding of soil moisture remains one of the most significant weaknesses in process-based storm flow models and the ability to predict runoff generation and erosion, while Grayson et al. (1992) argued that antecedent moisture is the parameter most likely to undermine model predictions (see also Mahanama et al., 2008).

Through its role in flow partitioning, antecedent soil moisture is also an important physical control on nutrient and sediment loss by overland flow (Casenave and Valentin, 1992; McDowell and Sharpley, 2002; McDowell et al., 2001) and has been demonstrated in several studies (Ceballos and Schnabel, 1998; Fitzjohn et al., 1998; Karnieli and Ben-Asher, 1993). The degree to which soil moisture is a primary control on runoff generation and erosion depends on basin size and the characteristics of the precipitation event. Castillo et al. (2003) have shown that the peak discharge and runoff during high-intensity, low-frequency storms is independent of initial soil water content, but is important in controlling runoff during medium- and low-intensity storms that are primarily responsible for erosion in semi-arid environments (see also Merz and Bardossy, 1998; Poesen and Hooke, 1997). Soil moisture is more important where vegetation increases the spatial variability of soil characteristics and produces a range of runoff and infiltration sites (Castillo et al., 2003). Similarly, Fitzjohn et al. (1998) have shown that spatial heterogeneity of surface moisture can reduce widespread catchment runoff and erosion by promoting discontinuity in hydrological pathways through the isolation of runoff-producing areas and the reabsorption of runoff generated upstream. By contrast, antecedent moisture can reduce surface crusting

through dissolution and disturbance, which limits runoff and interrill erosion (Le Bissonnais and Singer, 1992). Higher soil moisture contents are also responsible for limiting the development of hydrophobic compounds that coat particles and create a water-repellent layer following forest fires (Robichaud, 2000; Robichaud and Hungerford, 2000), similarly limiting post-fire runoff and erosion.

Spatial and temporal variations in soil moisture and other physical properties of the soil (e.g. clay composition, buried organic matter) are partly responsible for significant variation in the generation of overland flow (Godsey et al., 2004; Sharma et al., 1980; Sidle et al., 2000; Uchida et al., 1999; Ziegler et al., 2001). Soil moisture variation depends in part on the distribution of vegetation (Harden and Scruggs, 2003), topography (Julien and Moglen, 1990), and surface characteristics (Katra et al., 2008; Poesen et al., 1999); variables that are themselves dependent on the distribution of soil moisture. This spatial variation affects the amount and source of sediment delivered to the main channel. Seeger et al. (2004) found that variations in suspended sediment concentration during flood events are associated with different levels of humidity and rainfall, indicating different mechanisms for the development of runoff and sediment transport. A high level of antecedent soil moisture promotes runoff generation and sediment supply to the nearby channel, leading to an increase in suspended sediment concentration over the course of a storm as compared to 'normal' moisture levels that limit runoff and the sediment source to near the channel. The impact of antecedent soil moisture is much greater in small catchments (Seeger et al., 2004). More sediment is available to overland flow from dry soils that break down faster through slaking caused by the compression of entrapped air, while moist soils prevent slaking and limit the ability of the soil to be disaggregated (McDowell and Trudgill, 2000).

Sediment delivery to channels and the long-term shaping of a watershed can also occur

through mass movements triggered by soil at or close to saturation (e.g. Acharya et al., 2009; Benda and Dunne, 1997; Korup et al., 2004; Schuerch et al., 2006; Schwab et al., 2008). The frequency of mass movements is correlated with an increase in pore pressure resulting from heavy precipitation (Johnson and Sitar, 1990) and wetter climates (Brooks and Richards, 1994; Grove, 1972; Innes, 1983; Pitts, 1983). These events exhibit a threshold dependence on soil moisture since pore pressure, internal friction, and cohesion are primarily dependent on soil moisture content (Iverson, 2000; Pelletier et al., 1997; Ray and Jacobs, 2007; Van Asch et al., 1999). Numerous field studies have shown that antecedent soil moisture and surface runoff from upslope are important controls on the frequency and size of mass movements (e.g. Iverson, 2000; Larsen and Simon, 1993; Van Asch and Sukmantalya, 1993; Van Asch et al., 2009; Wiczczonek and Glade, 2005). The importance of soil moisture in controlling mass wasting is highlighted by the strong correlation between the location and frequency of landslides with soil drainage and the storage properties of soil moisture (Pelletier et al., 1997; Ray and Jacobs, 2007).

## 2 Aeolian processes

In addition to flow partitioning and mass wasting, soil moisture is also of particular importance in controlling sediment mobility in wind-blown environments; indeed, the role of moisture in surface deflation has been the focus of several field and laboratory studies (e.g. Azizov, 1977; Bisal and Hsieh, 1966; Brazel et al., 1986; Chepil, 1956; Davidson-Arnott et al., 2005; Fecan et al., 1999; Hotta et al., 1984; Jackson and Nordstrom, 1997; Logie, 1982; McKenna-Neuman and Maljaars, 1997; 1998; Sarre, 1987; Smalley, 1970; Wiggs et al., 2004). Several field studies have shown that the role of soil moisture is especially important in controlling the exchange of sediment between beach

and dune (e.g. Arens, 1996; Bauer et al., 2009; Hotta et al., 1984; Sherman et al., 1998; Svasek and Terwindt, 1974; Wiggs et al., 2004; Yang and Davidson-Arnott, 2005) where spatial and temporal variations in soil moisture can degrade the ability of currently available models to predict sediment transport (Sherman et al., 1998). Similarly, empirical relationships between dust events and climate tend to be inaccurate for prediction unless soil moisture is explicitly modeled (McTainsh et al., 1998). These relationships are complicated by the growth of vegetation in response to surface moisture. Vegetation can increase the threshold shear velocity by directly covering part of the surface and absorbing part of the wind momentum that would otherwise go to the sediment surface (Lancaster and Baas, 1998; Wolfe and Nickling, 1993). The role of vegetation in controlling dust emissions is most difficult to quantify because the relationships between soil moisture, vegetation cover, and turbulent shear stresses are poorly understood.

Low soil moisture combined with a lack of surface cover, strong winds and low humidity is an important factor associated with dust events (Nickling and Brazel, 1984; Lee et al., 1994; Stout, 2001). Interannual variations in dust emissions appear to be closely associated with antecedent precipitation (Nickling and Brazel, 1984) with dust emissions increasing during regionally wet El Niño events as a result of the erosion of vegetation and the deposition of new sediment (Reynolds et al., 2007). Nickling and Brazel (1984) observed that the occurrence of large-scale dust storms followed rainstorm activity but quickly dissipated. The resealing of clay crusts, which occurred after these rainstorms, was limited to the surface and consequently peel and curl to expose thin pieces of crust that are easily abraded by saltating grains. As the weaker surface crust was abraded, a strong stable clay/silt crust was left behind that was more difficult to abrade (Houser and Nickling, 2001). More recently, Stout (2001) described the annual dust cycle of the Southern

High Plains of Texas as a reflection of seasonal change in environmental factors, including surface moisture. Relationships among El Niño-Southern Oscillation (ENSO) events, dust sources, and dust composition also have recently been described by Okin and Reheis (2002) and Reheis (2006).

McKenna-Neuman and Nickling (1989) have shown that gravimetric water contents of  $\sim 1\%$  create sufficient interparticle cohesion that entrainment by aerodynamic forces is all but impossible, although there is much disagreement. Furthermore, Wiggs et al. (2004) have shown that soil moisture contents of up to 1.68% do not act as a barrier to sediment flux. Field studies have shown that soil moisture contents over which transport events can occur tend to be greater than those measured in the laboratory or predicted theoretically. Hotta et al. (1984) suggests that the threshold shear velocity increases by  $7.5 \text{ cm s}^{-1}$  for each 1% increase in soil moisture up to a moisture content of 8%, at which entrainment ceases. If saltating particles are present, then sediment entrainment is dominated by impact forces. Saltation is not inhibited until soil moisture content reaches 14% (Sarre and Chancey, 1990) and Jackson and Nordstrom (1998) observed transport at surface moistures of  $>7\%$  following light rains. The rate of transport increased following the rain event as the surface dried but remained greater than the transport rates measured on dry beaches. An earlier field study by Jackson and Nordstrom (1997) and a laboratory study by McKenna-Neuman and Maljaars (1997; 1998) indicated that transport rates are greater on damp surfaces due to the creation of a relatively elastic surface compared to dry, cohesion-less sediment. Drying of soil moisture is promoted by wind and is an important influence in environments where winds are competent to entrain dry but not moist sediments (Hotta et al., 1984; Jackson and Nordstrom, 1997; Logie, 1982; Sherman, 1990). Spatial variation in soil moisture and drying can lead to intermittent transport over short time periods

(Davidson-Arnott et al., 2005) that can vary over an order of magnitude in the presence of a steady wind (McKenna-Neuman and Langston, 2006). Spatial and temporal variations in surface moisture can lead to the organization of the saltation cloud into streamers (Baas and Sherman, 2005). The short time required to dry a thin layer of sediment during strong winds can lead to high transport within 10–30 minutes following a soaking rain (Gillette, 1999).

The moisture content in the top sediment layer and the ability of the wind to dry the surface is dependent (to varying degrees) on the humidity of the near-surface boundary layer. Low humidity and stronger winds in the afternoon extract moisture from the surface, creating the potential for dust emissions (Stout, 2001) – most dust events on the Texas High Plains are associated with humidity levels below 30%. The humidity of the near-surface boundary layer affects particle entrainment through changes in both the kinematic viscosity (i.e. the absolute viscosity of a fluid divided by its mass density) and the density of the air (McKenna-Neuman, 2003) which, in turn, affect the critical shear velocity required for the entrainment of sediment (Belly, 1964). An increase in density reduces the aerodynamic drag force on the bed, while an increase in the kinematic viscosity increases the drag force but limits turbulence generation at both the particle and boundary-layer scale. Vapor pressures at or close to saturation can maintain high soil moisture (and vice versa), although there is a decrease in the matric potential that reduces the ability of the water to be adsorbed to the charged sediment surface (Hillel, 1998; Jury et al., 1991). Despite the change in matric potential, interparticle cohesion in moist soils is the dominant control on sediment entrainment for particles with diameters  $<75 \mu\text{m}$  (Greeley and Iversen, 1985). McKenna-Neuman (2003) has shown that the role of surface moisture in aeolian transport is temperature dependent. The entrainment threshold tends to be lower in cold environments due to

a reduction in both the vapor pressure and matric potential, while in warm environments the high vapor pressure and matric potential leads to greater entrainment thresholds.

### 3 Periglacial processes

As noted by Washburn (1980), soil moisture is the sine qua non of frost action, in both seasonally and perennially frozen ground. The very definition of *permafrost* – subsurface earth materials remaining continuously at or below 0°C (i.e. the freezing point of water) for two or more years – is testament to the dominant role soil moisture plays in the geomorphic, ecological, and pedogenic processes of cold regions. A fundamental distinction in permafrost studies lies between *ice-rich* and *dry* permafrost (Bockheim and Tarnocai, 1998). Thaw of permafrost with abundant excess ice has potential for profound disturbance to ecological communities and human infrastructure at the surface through thaw consolidation and development of thermokarst terrain. In contrast, thaw of permafrost lacking appreciable water content holds little potential for disturbance at the surface.

Soil moisture plays a critical role in the long-term evolution and morphology of permafrost landscapes. Water redistributed in the soil column through freeze-thaw action creates impressive suites of landforms, including large and small frost mounds (pingos and palsas), networks of ice-wedge polygons that encompass many hectares and extend far below the ground surface, and solifluction lobes and terraces conveying vivid impressions of mass movement on hillslopes.

In medium-textured soils experiencing relatively slow freezing, moisture is attracted to freezing fronts, creating lenses of *segregated ice*. Locations at which the bottom of the seasonally thawed layer (the ‘active layer’) is coincident with the top of permafrost experience ‘two-side freezing’, a process that involves refreezing of the active layer both downward

from the ground surface and upward from the permafrost table. Two-sided freezing draws moisture from the central part of the active layer to feed growing ice lenses at the upper and lower freezing fronts. Under such conditions, refreezing of the active layer can be very slow, with isothermal conditions persisting for weeks or even months, owing to latent heat effects. This ‘zero curtain effect’ (Outcalt et al., 1990) is characteristic of regions with ice-rich permafrost. Repeated annually over long periods (decades to millennia) these processes create thick accretions of segregated ice in the upper permafrost, just below the base of the active layer. The large amounts of energy required to thaw this extremely ice-rich layer, referred to as the ‘transient layer’ (Shur et al., 2005), impart buffering qualities to it that resist the deep penetration of thaw that might otherwise occur rapidly under climatic warming.

Another unusual aspect of permafrost regions lies in the key role frozen water in the soil column provides for the interpretation of Quaternary history. The continuity, arrangement, structure, crystallography, and chemistry of the various forms of underground ice (Mackay, 1972) – in soil pores, as veins and wedges, in segregated lenses, as massive bodies of injection ice, and in other forms – has given rise to a sub-discipline of permafrost science known as *cryostratigraphy* (e.g. French, 1998). Under favorable circumstances cryostratigraphic analysis can provide detailed information about local and regional climatic and geomorphic conditions extending back thousands of years.

### IV Soil moisture and biogeography

Because plants are almost exclusively dependent on soil moisture to acquire needed water for photosynthesis, soil moisture and its spatial and temporal variability represent an indispensable quantity for evaluating and understanding vegetation patterns. Soil moisture recharge and utilization directly affect the distribution of

vegetation and its overarching canopy structure. Vegetation directly affects soil moisture inputs at the base of plant canopies through precipitation partitioning. Precipitation partitioning is the process whereby incident rainfall and snowfall is divided into canopy interception, throughfall, or stemflow (Hewlett, 1982). The extent to which the incident precipitation is intercepted, entrained as throughfall, or routed to the subcanopy as stemflow is dependent on both biotic (canopy) and abiotic (climatological) factors, such as aboveground surface area and storm event characteristics (Levia and Frost, 2003, 2006). Precipitation intercepted by plant canopies does not contribute to soil moisture recharge. Throughfall and stemflow inputs directly affect the spatiotemporal variability of soil moisture recharge (Durocher, 1990). In fact, stemflow in temperate forests has been observed to cause the water table beneath individual tree boles to reach the soil surface, thereby accounting for drastic differences in soil moisture beneath and between trees over space and time (Durocher, 1990). More recent work by Liang et al. (2009) documents the effect of stemflow on hillslope-scale soil moisture dynamics, noting the very different influence of stemflow on vertical soil water content compared to infiltration of rainfall. In short, soil moisture controls the establishment of plant communities which, in turn, influence soil moisture recharge and its usage. This eventually further affects the establishment and expansion of plant communities as well as land-surface atmosphere interactions and variety of geomorphic processes in general.

Soil moisture extraction by vegetation is dependent upon phenological stage, stomatal resistance, vegetation type, and vegetation density. It is also known that vegetation type and fractional vegetation control transpiration rates and thus availability and variability in soil moisture (Avissar and Pielke, 1989; Barlage and Zeng, 2004; Lyons, 2002; McPherson, 2007; Mahmood and Hubbard, 2004; Narisma and Pitman, 2003; Pielke, 2001). Soil moisture

availability, in turn, affects plant transpiration and related physiological activities.

Soil moisture exerts a substantial influence on vegetation, acting as a key control on stem water dynamics, stomatal regulation, and transpiration loss (Bréda and Granier, 1996; Kozłowski, 1958; Milburn, 1979). Recent work by David et al. (2004) clearly demonstrates the interplay among soil moisture, vapor pressure deficit, and sapflow for an evergreen oak in the Portuguese *montado*, sparse savannah-like ecosystems composed of evergreen green oak species (although this is a single tree, David et al. measured neighboring trees to verify their numbers were realistic). The sample tree in this study had access to deeper groundwater. Thus, soil moisture was not a limiting factor in transpiration for this particular tree (David et al., 2004). Their work demonstrated an inverse relationship between leaf water potential and sapflow. Sapflow reached a maximum of approximately  $0.2 \text{ mm h}^{-1}$  with a vapor pressure deficit of 2.0 kPa and remained at the upper limit of  $0.2 \text{ mm h}^{-1}$  with vapor pressure deficits to approximately 6.0 kPa (David et al., 2004). The upper limit of sapflow was controlled via stomatal regulation to prevent cavitation (David et al., 2004). In contrast to trees in well-watered environments, where transpiration loss is limited via stomatal control, the mismatch in water demand and supply (that vary as a function of soil type and texture) limit transpiration loss for vegetation in most soils that exhibit diurnal and seasonal fluctuations in soil moisture supply (Llorens et al., 2003). It also is important to note that interannual variations in leaf area index will have a detectable impact on transpiration loss (Bréda and Granier, 1996). Both flooding and cold soils have been found to reduce leaf conductance and transpiration for some species (Kreuzwieser et al., 2002; Teskey et al., 1984).

Research has demonstrated that transpiration differs greatly throughout the vertical profile of a forest canopy (Roberts, 2000). Although the upper portions of the forest canopy have lower proportions of leaf area compared to the mid-

portions of the canopy, the upper canopy contributes a disproportionately large share to total transpiration (Roberts, 2000), despite generally smaller leaves and closer coupling with the atmosphere than larger leaves in the lower canopy. The greater rates and amounts of transpiration from leaves in the upper canopy can be explained, in part, by the differing leaf morphology of sun versus shade leaves (Horn, 1971).

Canopy structural characteristics, including branching patterns and bark microrelief, also affects the distribution of soil moisture by affecting stemflow inputs to the forest floor (Levia and Herwitz, 2005; Van Stan and Levia, 2010). As branching patterns change with stand age, it has been found that stemflow production can change with stand age (Murakami, 2009). Murakami (2009) reported a drastic decrease in stemflow production from ages 9 to 10 in Japanese cypress and attributed the reduction to changes in tree architecture. Therefore, it is possible that soil moisture heterogeneity may be partly accounted for by change in canopy structure and corresponding alterations in stemflow production.

As briefly indicated above, soil moisture plays a critical role in determining development and evolution of a vegetation type in a region and subsequently impacts climate. The importance of land cover (particularly vegetation) and atmospheric interactions and associated feedbacks have been well documented (Adegoke et al., 2006; Fu, 2003; Narisma and Pitman, 2003; Pielke et al., 1999, 2007; Schneider and Eugster, 2005). Carleton et al. (2008) noted changes in local atmospheric circulation by vegetation boundary discontinuity which could be determined by availability of moisture. McPherson and Stensrud (2005) have shown changes in vegetation cover impact meso-scale atmospheric circulation and development of boundary layer atmosphere. Again, establishment of a particular type of vegetation is determined by soil moisture. Adegoke et al. (2003) have also shown changes in vegetation cover

along with soil moisture affect near-surface atmospheric moisture content and variety of other processes.

Whereas interception generally leads to a decrease in the input of incident precipitation to the forest floor, fog interception leads to fog drip and an increase in net precipitation inputs to forest soils (Cavelier et al., 1996; Holder, 2006). In fact, the thinning of forest in areas where fog is prevalent actually decreases fog precipitation inputs to the forest floor and, thus, soil moisture (Aboal et al., 2000). In the Canary Islands, it was found that decreases in leaf area index and basal area have led to lower inputs of moisture via fog drip (Aboal et al., 2000).

## V The integrative nature of soil moisture

Soil moisture is not just a process that is integral to climate, geomorphology, and biogeography – it truly lies at the intersection of all three branches of physical geography. A complete understanding of soil moisture and its spatial and temporal variability and impact draws upon interactions among and expertise gained from all three subdivisions. Soil moisture lies at the intersection of climatology, geomorphology, biogeography, and hydrology, thereby providing true integration of the subdisciplines rather than just supplying a common theme. While the interaction of climate, soils, vegetation, and hydrology have long been recognized and analyzed (see, for example, Mather, 1978), a cross-section of some of these interactions is worth noting.

While plant growth and development is clearly directly related to soil moisture, plant responses also can be indirectly associated with the effects of soil moisture. For example, Dyer (2002) found that soil moisture, as modeled by the climatic water balance, was an excellent predictor of the occurrence of American beech (*Fagus grandifolia*). Medler et al. (2002) also evaluated the interaction between soil moisture and the spatial and temporal patterns of snowfall

which, in turn, affect plant growth in the spring and the development of summer wildfires. They concluded that increases in severe wildfire resulting from decreased soil moisture are only moderately related to lowered snowfall in the previous winter. Land surface-atmosphere interactions also were found to be important in the lower Mississippi River basin by Brown and Wax (2007). In particular, they observed that differences in soil types which led to spatially variable soil moisture regimes caused significant differences in seasonal maximum and minimum air temperatures within the alluvial valley.

The highly varied spatial and temporal character of soil moisture easily lends itself to the study of its impacts on other environmental variables. Walsh et al. (1998), for example, used scale, pattern, and process with remote sensing and geographic information systems to couple geomorphic variables – including soil moisture – to biogeographic and landscape ecological processes. Bridge and Johnson (2000) found significant relationships between geomorphic principles and vegetation gradients, particularly as they relate to soil moisture and water availability. Cammeraat (2002) also focused on scale issues in geomorphology to examine how hydrological variables, such as soil moisture, are influenced by biological and climatological processes. Similarly, Western and Bloschl (1999) and Western et al. (2002) evaluated this variability in soil moisture and showed the importance of its non-linear response in environmental analyses. More recently, research has focused on a more holistic treatment of ecosystems wherein the climate, geomorphology, vegetation and soils are integrated to provide a better assessment of the overall interactions between the various components of the environment (see, for example, Abella, 2003; Botter et al., 2007; Hughes, 1997). The take-away message associated with these studies is that the entire physical system as represented by the biosphere, the atmosphere, the hydrosphere, and the lithosphere must be considered as a collective whole

– not a collection of unassociated parts – if the environment is to be properly represented. Soil moisture, since its spatial and temporal characteristics are derived from the climate, land surface, and vegetative characteristics, is therefore an integrative component which summarizes and affects all components of the environment.

## **VI Summary and recommendations for future research**

As Mather (1993) suggested in his Presidential Address to the *Association of American Geographers*, geographers should ‘emphasize those aspects that bind us together rather than separate us’. Soil moisture is truly a variable that binds together the various branches of physical geography. From its impact on and the influence on it by weather and climate, geomorphology and soils, biogeography, and hydrology, soil moisture integrates all aspects of physical geography. Thus, we call on physical geographers to use soil moisture as a unifying theme in physical geography.

With that in mind, we suggest several areas of research that illustrate the integrative nature of soil moisture. For example, the WMO (2008) report on ‘Future Climate Change Research and Observations’ recommends that soil moisture data should be assembled, quality controlled, and harmonized because of its importance in (1) providing an improved understanding of land-atmosphere interactions, (2) the development of seasonal-to-decadal climate forecasting tools, (3) calibration, validation, and improvements in the physical parameterizations in regional and global land surface models, (4) developing and validating algorithms for determining estimates of soil moisture using satellite-based techniques, and (5) monitoring and detecting climate variability and change. Although much of this work is already underway, we anticipate and hope that more research will be focused on understanding the nature of

land-atmosphere interactions and the role that soil moisture and vegetation play in influencing climate on seasonal to decadal timescales. We also foresee continued efforts to improve the accuracy of soil moisture models and the representation of soil moisture in general circulation models (GCMs) to improve the accuracy of soil energy and water fluxes to the atmosphere. But, most importantly, efforts will be focused on the development of new modeling and observational technologies, including both remotely sensed surface soil wetness products and additional in situ measurements of soil moisture. These data acquisition platforms will provide new opportunities for land surface model calibration and for developing and validating satellite-derived soil moisture algorithms.

Patch-scale temporal and spatial heterogeneity of soil moisture in forested and agricultural ecosystems is also a key element that remains inadequately understood. How do the interrelationships between and among canopy structural components, microclimates, and soil characteristics influence the timing and spatial patterns of soil moisture? A detailed understanding of soil moisture dynamics in relation to vegetative canopies will yield important insights into the exploitation of environmental heterogeneity by plants. Such knowledge at the patch scale could allow evapotranspiration models to better predict water use by plants and, ultimately, improve crop yields from precision agriculture.

From the perspective of process geomorphology, soil moisture is an important but by no means simple control on wind-blown sediment entrainment and transport, flow partitioning, mass wasting, and weathering. Contemporary research tends to be focused on spatial and temporal variations in soil moisture at a range of scale and the feedbacks therein to improve the ability of currently available models to predict sediment transport in the prototype, such as the exchange of sediment between beach and dune critical to dune recovery following storms. Moreover, the ability to discern, at high spatial

resolution, variations in soil moisture and the effects of freeze and thaw over broad areas and in three dimensions is urgently needed. Methodologies for creating detailed maps of surficial heave and subsidence are under development and will prove instrumental for creating maps of hazard potential at local scales.

## References

- Abbaspour KC, Hall JW, and Moon DE (1992) A yield model for use in determining crop insurance premiums. *Agricultural and Forest Meteorology* 60(1–2): 33–51.
- Abella SR (2003) Quantifying ecosystem geomorphology on the southern Appalachian Mountains. *Physical Geography* 24(6): 488–501.
- Aboal JR, Jiménez MS, Morales D, and Gil P (2000) Effects of thinning on throughfall in Canary Islands pine forest- the role of fog. *Journal of Hydrology* 238: 218–230.
- Acharya G, Cochrane TA, Davies T and Bowman E (2009) The influence of shallow landslides on sediment supply: A flume-based investigation using sandy soil. *Engineering Geology* 109(3–4): 161–169.
- Adegoke JO, Pielke RA, and Carleton AM (2006) Observational and modeling studies of the impact of agriculture-related land use change on climate in the central U.S. *Agricultural and Forest Meteorology* 142: 203–215.
- Adegoke JO, Pielke RA, Eastman J, Mahmood R, and Hubbard GK (2003) Impact of irrigation of midsummer surface fluxes and temperature under dry synoptic conditions: A regional atmospheric model study of the U.S. High Plains. *Monthly Weather Review* 131: 556–564.
- Anderson K and Croft H (2009) Remote sensing of soil surface properties. *Progress in Physical Geography* 33: 457–473.
- Anderson SP (2005) Glaciers show direct linkage between erosion rate and chemical weathering fluxes. *Geomorphology* 67(1–2): 147–157.
- Andreadis KM and Lettenmaier DP (2006) Trends in 20th century drought over the continental United States. *Geophysical Research Letters* 33(L10403). doi: 10.1029/2006GL025711.
- Arens SM (1996) Rates of aeolian transport on a beach in a temperate humid climate. *Geomorphology* 17: 3–18.
- Avissar R and Pielke RA (1989) A parameterization of heterogeneous land surfaces for atmospheric numerical

- models and its impact on regional meteorology. *Monthly Weather Review* 117: 2113–2136.
- Azizov A (1977) Influence of soil moisture on the resistance of soil to wind erosion. *Soviet Soil Science* 9: 105–108.
- Baas ACW and Sherman DJ (2005) The formation and behavior of aeolian streamers. *Journal of Geophysical Research* 110: F03011. doi: 10.1029/2004JF000270.
- Balsamo G, Mahfouf JF, Belair S, and DeBlonde G (2007) A land data assimilation system for soil moisture and temperature: An information content study. *Journal of Hydrometeorology* 8: 1225–1242.
- Barlage M and Zeng X (2004) The effects of observed fractional vegetation cover on the land surface climatology of the community land model. *Journal of Hydro-meteorology* 5: 823–830.
- Bauer BO, Davidson-Arnott RGD, Hesp PA, Namikas SL, Ollerhead J, and Walker IJ (2009) Aeolian sediment transport on a beach: Surface moisture, wind fetch, and mean transport. *Geomorphology* 105(1–2): 106–116.
- Belly Y (1964) *Sand movement by wind*. U.S. Army Corps of Engineers, Coastal Engineering Research Center, Technical Memo 1, Addendum III, 24 pp.
- Benda L and Dunne T (1997) Stochastic forcing of sediment supply to channel networks from landsliding and debris flow. *Water Resources Research* 33: 2849–2863.
- Bisal F and Hsieh J (1966) Influence of moisture on erodibility of soil by wind. *Soil Science* 3: 143–146.
- Bockheim JG and Tarnocai C (1998) Nature, occurrence and origin of dry permafrost. In: Lewkowicz AG (ed.) *Proceedings of the Seventh International Conference on Permafrost*. Québec: Centre d'études nordiques, Université Laval, Publication 57, 57–63.
- Botter G, Porporato A, Daly E, Rodriguez-Iturbe I, and Rinaldo A (2007) Probabilistic characterization of base flows in river basins: Roles of soil, vegetation, and geomorphology. *Water Resources Research* 43: W06404. doi: 10.1029/2006WR005397.
- Brazel AJ and Nickling WG (1987) Dust storms and their relation to moisture in the Sonoran-Mojave Desert region of the Southwestern United States. *Journal of Environmental Management* 24(3): 279–291.
- Brazel AJ, Nickling WG, and Lee J (1986) Effect of antecedent moisture conditions on dust storm generation in Arizona. In Nickling WG (ed.) *Aeolian Geomorphology*. Proceedings of the 17th Annual Binghamton Symposium, Allen and Unwin, 261–271.
- Bréda N and Granier A (1996) Intra- and interannual variations of transpiration, leaf area index and radial growth of a sessile oak stand (*Quercus patraea*). *Annals of Forest Science* 53: 521–536.
- Bridge SRJ and Johnson EA (2000) Geomorphic principles of terrain organization and vegetation gradients. *Journal of Vegetation Science* 11: 57–70.
- Brooks SM and Richards KS (1994) The significance of rainstorm variations to shallow translational hillslope failure. *Earth Surface Processes and Landforms* 19: 85–94.
- Brown ME and Wax CL (2007) Temperature as an indicator of landform influence on atmospheric processes. *Physical Geography* 28(2): 148–157.
- Brubaker LK, Entekhabi D, and Eagleson SP (1993) Estimation of continental precipitation recycling. *Journal of Climate* 6: 1077–1089.
- Cammeraat LH (2002) A review of two strongly contrasting geomorphological systems within the context of scale. *Earth Surface Processes and Landforms* 27: 1201–1222.
- Carleton AM, Travis DJ, Adegoke JO, Arnold DL, and Curran S (2008) Synoptic circulation and land surface influences on convection in the Midwest U.S. 'Corn Belt' during the summers of 1999 and 2000. Part II: Role of vegetation boundaries. *Journal of Climate* 21: 3617–3641.
- Casenave A and Valentin C (1992) A runoff capability classification system based on surface features criteria in semi-arid areas of West Africa. *Journal of Hydrology* 130: 231–249.
- Castillo VM, Gomez-Plaza A, and Martinez-Mena M (2003) The role of antecedent soil water content in the runoff response of semiarid catchments: A simulation approach. *Journal of Hydrology* 284(1–4): 114–130.
- Cavelier J, Solis D, and Jaramillo MA (1996) Fog interception in montane forests across the Central Cordillera of Panama. *Journal of Tropical Ecology* 12: 357–369.
- Ceballos A and Schnabel S (1998) Hydrological behavior of a small catchment in the dehesa landuse system (Extremadura, SW Spain). *Journal of Hydrology* 210: 146–160.
- Chang H-I, Niyogi D, Kumar A, Kishtawal CM, Dudhia J, Chen F, et al. (2009) Possible relation between land surface feedback and the post-landfall structure of monsoon depressions. *Geophysical Research Letters* 36: L15826. doi: 10.1029/2009GL037781.
- Chen F and Avissar R (1994) Impact of land-surface moisture variability on local shallow convective cumulus and

- precipitation in large-scale models. *Journal of Applied Meteorology* 33: 1382–1401.
- Chepil WS (1956) Influence of moisture on erodibility of soil by wind. *Soil Science Society Proceedings* 20: 288–292.
- Choi HI and Liang X-Z (2010) Improved terrestrial hydrologic representation in mesoscale land surface models. *Journal of Hydrometeorology* 11: 797–809.
- Clark AC and Arritt RW (1995) Numerical simulations of the effect of soil moisture and vegetation cover on the development of deep convection. *Journal of Applied Meteorology* 34: 2029–2045.
- Clark DB, Taylor CM, and Thorpe AJ (2004) Feedback between the land surface and rainfall at convective length scales. *Journal of Hydrometeorology* 5(4): 625–639.
- David TS, Ferreira MI, Cohen S, Pereira JS, and David JS (2004) Constraints on transpiration from an evergreen oak tree in southern Portugal. *Agricultural and Forest Meteorology* 122: 193–205.
- Davidson-Arnott RGD, MacQuarrie K, and Aagaard T (2005) The effect of wind gusts, moisture content and fetch length on sand transport on a beach. *Geomorphology* 68(1–2): 115–129.
- De Laat ATJ and Maurellis AN (2006) Evidence for influence of anthropogenic surface processes on lower tropospheric and surface temperature trends. *International Journal of Climatology* 26: 897–913.
- DeLiberty TL and Legates DR (2003) Interannual and seasonal variability of modelled soil moisture in Oklahoma. *International Journal of Climatology* 23: 1057–1986.
- DeLiberty TL and Legates DR (2008) Spatial variability and persistence of soil moisture in Oklahoma. *Physical Geography* 29(2): 121–139.
- Dirmeyer PA (2000) Using a global soil wetness dataset to improve seasonal climate simulation. *Journal of Climate* 13: 2900–2922.
- Dong J, Ni-Meister W, and Houser PR (2007) Impacts of vegetation and cold season processes on soil moisture and climate relationships over Eurasia. *Journal of Geophysical Research* 112: 1–11.
- Dorigo WA, Zurita-Milla R, de Wit AJW, Brazile J, Singh R, and Schaepman ME (2007) A review on reflective remote sensing and data assimilation techniques for enhanced agroecosystem modeling. *International Journal of Applied Earth Observation and Geoinformation* 9: 165–193.
- Douville H, Chauvin F, and Broqua H (2001) Influence of soil moisture on the Asian and African Monsoons. Part I: Mean monsoon and daily precipitation. *Journal of Climate* 14: 2381–2403.
- Dunne T and Black RD (1970) Partial area contributions to storm runoff in a small New England watershed. *Water Resources Research* 6: 1296–1311.
- Durocher MG (1990) Monitoring of spatial variability of forest interception. *Hydrological Processes* 4: 215–229.
- Dyer JM (2002) A comparison of moisture scalars and water budget methods to assess vegetation-site relationships. *Physical Geography* 23(3): 245–258.
- Ek MB and Holtslag AAM (2004) Influence of soil moisture on boundary layer cloud development. *Journal of Hydrometeorology* 5: 86–99.
- Entin JK, Robock A, Vinnikov KY, Hollinger SE, Liu S, and Namkai A (2000) Temporal and spatial scales of observed soil moisture variations in the extratropics. *Journal of Geophysical Research* 105: 11,865–11,877.
- Fecan F, Marticorena B, and Bergametti G (1999) Parameterization of the increase of the aeolian erosion threshold wind friction velocity due to soil moisture for arid and semiarid areas. *Annales Geophysicae*, 17(1): 149–157.
- Feddema JJ (2007) A revised Thornthwaite-type global climate classification. *Physical Geography* 26(6): 442–466.
- Findell KL and Eltahir EAB (2003) Atmospheric controls on soil moisture-boundary layer interactions: Three-dimensional wind effects. *Journal of Geophysical Research* 108(D8): 8385–8405.
- Fitzjohn C, Ternan JL, and Williams AG (1998) Soil moisture variability in a semi-arid gully catchment: Implications for runoff and erosion control. *Catena* 32: 55–70.
- Francesca V, Osvaldo F, Stefano P, and Paola RP (2010) Soil moisture measurements: Comparison of instrumentation performances. *Journal of Irrigation and Drainage Engineering – ASCE* 136: 81–89.
- French HM (1998) An appraisal of cryostratigraphy in north-west Arctic Canada. *Permafrost and Periglacial Processes* 9: 297–312.
- Fu C (2003) Potential impacts of human-induced land cover change on East Asia monsoon. *Global and Planetary Change* 37: 219–229.
- Fulakeza M, Druyan LM, and Krishnamurti TN (2002) A simple soil moisture scheme for regional climate

- simulations in the tropics. *Meteorology and Atmospheric Physics* 79: 105–126.
- Gamble DW and Meentemeyer VG (1997) A synoptic climatology of extreme unseasonable floods in the southeastern United States, 1950–1990. *Physical Geography* 18(6): 496–524.
- Gillette DA (1999) A qualitative geophysical explanation for hot spot dust emitting source regions. *Contributions to Atmospheric Physics* 72: 67–77.
- Godsey S, Elsenbeer H, and Stallard R (2004) Overland flow generation in two lithologically distinct rainforest catchments. *Journal of Hydrology* 295: 276–290.
- Grasso LD (2000) A numerical simulation of dryline sensitivity to soil moisture. *Monthly Weather Review* 128: 2816–2834.
- Grayson RB, Moore ID, and McMahon TA (1992) Physically based hydrologic modelling, II. Is the concept realistic? *Water Resources Research* 28: 2659–2666.
- Greeley R and Iversen JD (1985) *Wind as a Geological Process on Earth, Mars, Venus and Titan*. New York: Cambridge University Press, 333 pp.
- Grove JM (1972) The incidence of landslides, avalanches, and floods in eastern Norway during the little ice age. *Artic Alpine Research* 4: 131–138.
- Harden CP and Scruggs PD (2003) Infiltration on mountain slopes: A comparison of three environments. *Geomorphology* 55: 5–24.
- Hewlett JD (1982) *Principles of Forest Hydrology*. Athens, GA: University of Georgia Press, 183 pp.
- Hewlett JD and Hibbert AR (1967) Factors affecting the response of small watersheds to precipitation in humid areas. In: Sopper WE and Lull HW (eds) *International Symposium on Forest Hydrology*. Oxford: Pergamon, 275–290.
- Hillel D (1998) *Environmental Soil Physics*. New York: Academic Press, 771 pp.
- Holder CD (2006) The hydrological significance of cloud forests in the Sierra de las Minas Biosphere Reserve, Guatemala. *Geoforum* 37: 82–93.
- Hong SY and Kalnay E (2000) Role of sea surface temperature and soil-moisture feedback in the 1988 Oklahoma-Texas drought. *Nature* 408: 842–844.
- Horn HS (1971): *The Adaptive Geometry of Trees*. Monographs in Population Biology 3. Princeton, NJ: Princeton University Press, 144 pp.
- Horton RE (1933) The role of infiltration in the Hydrologic Cycle. *Transactions of the American Geophysical Union* 14: 446–460.
- Hossain F (2006) Towards formulation of a space-borne system for early warning of floods: Can cost-effectiveness outweigh prediction uncertainty? *Natural Hazards* 37: 263–276.
- Hotta S, Kubota S, Katori S, and Horikawa K (1984) Sand transport by wind on a wet sand surface. *Proceedings of the 19th Coastal Engineering Conference of the ASCE*. New York: ASCE, 1265–1281.
- Houser CA and Nickling WG (2001) The factors influencing the abrasion efficiency of saltating grains on a clay-crustured playa. *Earth Surface Processes and Landforms* 48: 255–267.
- Huang J, Van Den Dool HM, and Georgakakos KP (1996) Analysis of model-calculated soil moisture over the United States (1931–1993) and applications to long-range temperature forecasts. *Journal of Climate* 9: 1350–1362.
- Hughes FMR (1997) Floodplain biogeomorphology. *Progress in Physical Geography* 21(4): 501–529.
- Innes JL (1983) Lichenometric dating of debris-flow deposits in the Scottish highlands. *Earth Surface Processes* 8: 579–588.
- Iverson RM (2000) Landslide triggering by rain infiltration. *Water Resources Research* 36: 1897–1910.
- Jackson NL and Nordstrom KF (1997) Effects of time-dependent moisture content of surface sediments on aeolian transport rates across a beach, Wildwood, New Jersey, USA. *Earth Surface Processes and Landforms* 22: 611–621.
- Jackson NL and Nordstrom KF (1998) Aeolian transport of sediment on a beach during and after rainfall, Wildwood, NJ, USA. *Geomorphology* 22(2): 151–157.
- Johnson KA and Sitar N (1990) Hydrologic conditions leading to debris-flow initiation. *Canadian Geotechnical Journal* 27(6): 789–801.
- Jones AR and Brunsell NA (2009) A scaling analysis of soil moisture–precipitation interactions in a regional climate model. *Theoretical and Applied Climatology* 98: 221–235.
- Julien PY and Moglen GE (1990) Similarity and length scale for spatially varied overland flow. *Water Resources Research* 26: 1819–1832.
- Jury WA, Gardner WR, and Gardner WH (1991) *Soil Physics*. New York: Wiley, 328 pp.
- Karnieli A and Ben-Asher J (1993) A daily runoff simulation in semi-arid watersheds based on soil water deficit calculations. *Journal of Hydrology* 149: 9–25.

- Katra I, Lavee H, and Sarah P (2008) The effect of rock fragment size and position on topsoil moisture on arid and semi-arid hillslopes. *Catena* 72: 49–55.
- Kerr YH (2007) Soil moisture from space: Where are we? *Hydrogeology Journal* 15: 117–120.
- Kidd C, Levizzani V, and Bauer P (2009) A review of satellite meteorology and climatology at the start of the twenty-first century. *Progress in Physical Geography* 33: 474–489.
- Kirkby MJ and Chorley RJ (1967) Throughflow, overland flow and erosion. *Bulletin of the International Association of Scientific Hydrology* 12: 5–21.
- Korup O, McSaveney KO, and Davies TRH (2004) Sediment generation and delivery from large historic landslides in the Southern Alps, New Zealand. *Geomorphology* 61: 189–207.
- Koster RD and Suarez MJ (2003) Impact of land surface initialization on seasonal precipitation and temperature prediction. *Journal of Hydrometeorology* 4: 408–423.
- Koster RD, Dirmeyer PA, Guo ZC, Bonan G, Chan E, Cox P, et al. (2004) Regions of strong coupling between soil moisture and precipitation. *Science* 305: 1138–1140.
- Kozlowski TT (1958): Water relations and growth of trees. *Journal of Forestry* 56: 498–501.
- Kreuzwieser J, Fürniss S, and Rennenberg H (2002) Impact of waterlogging on the N-metabolism of flood tolerant and non-tolerant tree species. *Plant, Cell and Environment* 25: 1039–1049.
- Lacava T, Greco M, Di Leo EV, Martino G, Pergola N, Sannazzaro F, et al. (2005) Monitoring soil wetness variations by means of satellite passive microwave observations: The HYDROPTIMET study cases. *Natural Hazards and Earth System Sciences* 5: 583–592.
- Lancaster N and Baas A (1998) Influence of vegetation cover on sand transport by wind: Field studies at Owens Lake, California. *Earth Surface Processes and Landforms* 23(1): 69–82.
- Larsen MC and Simon A (1993) A rainfall intensity-duration threshold for landslides in a humid-tropical environment, Puerto Rico. *Geografiska Annaler* 75: 13–23.
- Le Bissonnais YL and Singer MJ (1992) Crusting, runoff, and erosion response to soil water content and successive rainfalls. *Soil Science Society of America Journal* 56: 1898–1903.
- Lee JA, Moffett KE, Allen BL, Peterson RE, and Gregory JM (1994) Environmental controls on blowing dust direction at Lubbock, Texas, USA. *Earth Surface Processes and Landforms* 5: 437–449.
- LeMone AM, Chen F, Alfieri GJ, Tewari M, Geerts B, Miao Q, et al. (2007) Influence of land cover and soil moisture on horizontal distribution of sensible and latent heat fluxes in southeast Kansas during IHOP\_2002 and CASES-97. *Journal of Hydrometeorology* 8(1): 68–87.
- Levia DF and Frost EE (2003) A review and evaluation of stemflow literature in the hydrologic and biogeochemical cycles of forested and agricultural ecosystems. *Journal of Hydrology* 274: 1–29.
- Levia DF and Frost EE (2006) Variability of throughfall volume and solute inputs in wooded ecosystems. *Progress in Physical Geography* 30: 605–632.
- Levia DF and Herwitz SR (2005) Interspecific variation of bark water storage capacity of three deciduous tree species in relation to stemflow yield and solute flux to forest soils. *Catena* 64: 117–137.
- Liang W-L, Kosugi K, and Mizuyama T (2009) A three-dimensional model of the effect of stemflow on soil water dynamics around a tree on a hillslope. *Journal of Hydrology* 366: 62–75.
- Llorens L, Peñuelas J, and Filella I (2003) Diurnal and seasonal variations in the photosynthetic performance and water relations of two co-occurring Mediterranean shrubs, *Erica multiflorai* and *Globularia alypum*. *Physiologia Plantarum* 118: 84–95.
- Logie M (1982) Influence of roughness elements and soil moisture of sand to wind erosion. *Catena* 1: 161–173.
- Lyons TJ (2002) Clouds prefer native vegetation. *Meteorology and Atmospheric Physics* 80: 131–140.
- McCabe GJ, Palecki MA, and Betancourt JL (2004) Pacific and Atlantic Ocean influences on multidecadal drought frequency in the United States. *Proceedings of the National Academy of Sciences* 101(12): 4136–4141.
- McCabe GJ Jr, Legates DR, and Lins HF (2010) Variability and trends in dry day frequency and dry event length in the Southwestern United States. *Journal of Geophysical Research* 115: D07108. doi: 10.1029/2009JD012866.
- McCumber MC and Pielke RA (1981) Simulation of the effects of surface fluxes of heat and moisture in a mesoscale Numerical Model. *Journal of Geophysical Research* 86: 9929–9938.
- McDowell RW and Sharpley AN (2002) Approximating Phosphorus Release from Soils to Surface Runoff and Subsurface Drainage. *Journal of Environmental Quality* 30: 508–520.

- McDowell RW and Trudgill S (2000) Variation of phosphorus loss from a small catchment in south Devon, UK. *Agriculture Ecosystems and Environment* 79(2–3): 143–157.
- McDowell RW, Sharpley AN, Condron LM, Haygarth PM, and Brookes PC (2001) Processes controlling soil phosphorus release to runoff and implications for agricultural management. *Nutrient Cycling in Agroecosystems* 59: 269–284.
- Mackay JR (1972) The world of underground ice. *Annals of the Association of American Geographers* 62: 1–22.
- McKenna-Neuman C (2003) Effects of temperature and humidity upon the entrainment of sedimentary particles by wind. *Boundary-Layer Meteorology* 108: 1573–1472.
- McKenna-Neuman C and Langston G (2006) Measurement of water content as a control of particle entrainment by wind. *Earth Surface Processes and Landforms* 31: 303–317.
- McKenna-Neuman C and Maljaars M (1997) Wind tunnel measurement of boundary layer response to sediment transport. *Boundary Layer Meteorology* 84: 67–83.
- McKenna-Neuman C and Maljaars M (1998) A wind tunnel study of the influence of pore water on aeolian sediment transport. *Journal of Arid Environments* 39(3): 403–419.
- McKenna-Neuman C and Nickling WG (1989) A theoretical and wind tunnel investigation of the effect of capillary water on the entrainment of sediment by wind. *Canadian Journal of Soil Science* 69: 79–96.
- McPherson RA (2007) A review of vegetation-atmosphere interactions and their influences on mesoscale phenomena. *Progress in Physical Geography* 31(3): 261–285.
- McPherson RA and Stensrud DJ (2005) Influences of a winter wheat belt on the evolution of the boundary layer. *Monthly Weather Review* 133: 2178–2199.
- McTainsh GH, Lynch AW, and Tews EK (1998) Climatic controls upon dust storm occurrence in eastern Australia. *Journal of Arid Environments* 39(3): 457–466.
- Mahanama SPP, Koster RD, Reichle RH, and Zubair L (2008) The role of soil moisture initialization in sub-seasonal and seasonal streamflow prediction – a case study in Sri Lanka. *Advances in Water Resources* 31: 1333–1343.
- Mahmood R (1996) Scale issues in soil moisture modeling: Problems and prospects. *Progress in Physical Geography* 20(3): 273–291.
- Mahmood R and Hubbard KG (2002) Anthropogenic land-use change in the North American tall grass-short grass transition and modification of near-surface hydrologic cycle. *Climate Research* 21: 83–90.
- Mahmood R and Hubbard KG (2004) An analysis of simulated long-term soil moisture data for three land uses under contrasting hydroclimatic condition in the Northern Great Plains. *Journal of Hydrometeorology* 5: 160–180.
- Mahmood R and Hubbard KG (2007) Relationship between soil moisture of near surface and multiple depths of the root zone under heterogeneous land uses and varying hydroclimatic conditions. *Hydrological Processes* 21: 3449–3462.
- Mather JR (1978) *The Climatic Water Budget in Environmental Analysis*. Lexington, MA: Lexington Books, 239 pp.
- Mather JR (1985) The water budget and the distribution of climate, vegetation, and soils. *Publications in Climatology* 38(2).
- Mather JR (1991) A history of hydroclimatology. *Physical Geography* 12(3): 260–273.
- Mather JR (1993) A shared vision. *Annals of the Association of American Geographers* 83(4): 561–567.
- Medler MJ, Montesano P, and Robinson DA (2002) Examining the relationship between snowfall and wildfire patterns in the western United States. *Physical Geography* 23(4): 335–342.
- Meng L and Quiring SM (2010) Observational relationship of sea surface temperatures and precedent soil moisture with summer precipitation in the USA Great Plains. *International Journal of Climatology* 30(6): 884–893.
- Merz B and Bardossy A (1998) Effects of spatial variability on the rainfall runoff process in a small loess catchment. *Journal of Hydrology* 212/213: 304–317.
- Milburn JA (1979) *Water Flow in Plants*. London: Longman, 225 pp.
- Murakami S (2009) Abrupt changes in annual stemflow with growth in a young stand of Japanese cypress. *Hydrological Research Letters* 3: 32–35.
- Namias J (1959) Persistence of mid-tropospheric circulation between adjacent months and seasons. In: Bolin B (ed.) *The Atmosphere and the Sea in Motion, Rossby Memorial Volume*, New York: Rockefeller Institute Press/Oxford University Press, 240–248.
- Narasimhan B and Srinivasan R (2005) Development and evaluation of Soil Moisture Deficit Index (SMDI) and Evapotranspiration Deficit Index (ETDI) for

- agricultural drought monitoring. *Agricultural and Forest Meteorology* 133: 69–88.
- Narisma GT and Pitman AJ (2003) The impact of 200 years of land cover change on the Australian near-surface climate. *Journal of Hydrometeorology* 4: 424–436.
- Nickling WG and Brazel AJ (1984) Temporal and spatial characteristics of Arizona dust storms (1965–1980). *International Journal of Climatology* 4: 645–660.
- Ni-Meister W (2008) Recent advances on soil moisture data assimilation. *Physical Geography* 29: 19–37.
- Oglesby RJ and Erickson DJ III (1989) Soil moisture and the persistence of North American drought. *Journal of Climate* 2: 1362–1380.
- Okin GS and Reheis MC (2002) An ENSO predictor of dust emission in the southwestern United States. *Geophysical Research Letters* 29(9): 1332. doi: 10.1029/2001GL014494.
- Ookouchi Y, Segal M, Kessler RC, and Pielke RA (1984) Evaluation of soil moisture effects on the generation and modification of mesoscale circulations. *Monthly Weather Review* 112: 2281–2292.
- Outcalt SI, Nelson FE, and Hinkel KM (1990) The zero-curtain effect: Heat and mass transfer across an isothermal region in freezing soil. *Water Resources Research* 26: 1497–1509.
- Pal JS and Eltahir EAB (2001) Pathways relating soil moisture conditions to future summer rainfall within a model of the land–atmosphere system. *Journal of Climate* 14: 1227–1242.
- Pan Z, Takle E, Segal M, and Turner R (1996) Influences of model parameterization schemes on the response of rainfall to soil moisture in the Central United States. *Monthly Weather Review* 124: 1786–1802.
- Pelletier JD, Malmud BD, Blodgett T, and Turcotte DL (1997) Scale-invariance of soil moisture variability and its implications for the frequency-size distribution of landslides. *Engineering Geology* 48: 255–268.
- Phillips JD, Turkington AV, and Marion DA (2008) Weathering and vegetation effects in early stages of soil formation. *Catena* 72: 21–28.
- Pielke RA (2001) Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convection rainfall. *Reviews of Geophysics* 39: 151–177.
- Pielke RA, Adegoke J, Beltran-Przekurat A, Hiemstra CA, Lin J, Nair US, et al. (2007) An overview of regional land use and land cover impacts on rainfall. *Tellus B* 59: 587–601.
- Pielke RA, Walko RL, Steyaert LT, Vidale PL, Liston GE, Lyons WA, et al. (1999) The influence of anthropogenic landscape changes on weather in south Florida. *Monthly Weather Review* 127: 1663–1673.
- Pinty J-P, Mascart P, Richard E, and Rosset R (1989) An investigation of mesoscale flows induced by vegetation inhomogeneities using an evapotranspiration model calibrated against HAPEX-MOBILHY data. *Journal of Applied Meteorology* 28: 976–992.
- Pitts J (1983) The temporal and spatial development of landslides in the Axmouth-Lyme Regis undercliffs national nature reserve. *Earth Surface Processes and Landforms* 8(6): 589–603.
- Poesen JWA and Hooke JM (1997) Erosion, flooding and channel management in Mediterranean environments of Southern Europe. *Progress in Physical Geography* 21(2): 157–199.
- Poesen J, De Luna E, Franca A, Nachtergaele J, and Govers G (1999) Concentrated flow erosion rates as affected by rock fragment cover and initial soil moisture content. *Catena* 36: 315–329.
- Quintanar A, Mahmood R, Loughrin J, and Lovanh NC (2008) A coupled MM5-Noah land surface model-based assessment of sensitivity of planetary boundary layer variables to anomalous soil moisture conditions. *Physical Geography* 29(1): 54–78.
- Quintanar AI, Mahmood R, Motley MV, Yan J, Loughrin J, and Lovanh NC (2009) Boundary layer trajectory dispersion sensitivity to soil moisture conditions as simulated by the MM5 and the Noah Land Surface model. *Atmospheric Environment* 43: 3774–3785.
- Quiring SM (2004) Developing a real-time agricultural drought monitoring system for Delaware. *Publications in Climatology* 57(1).
- Quiring SM and Goodrich GB (2008) Nature and causes of the 2002 to 2004 drought in the southwestern United States compared with the historic 1953 to 1957 drought. *Climate Research* 36: 41–52.
- Quiring SM and Legates DR (2008) Application of CERES-Maize for within-season prediction of rainfed corn yields in Delaware, USA. *Agricultural and Forest Meteorology* 148: 964–975.
- Rabin RM, Stadler S, Wetzel PJ, Stensrud DJ, and Gregory M (1990) Observed effects of landscape variability on convective clouds. *Bulletin of the American Meteorological Society* 71: 272–280.

- Rasmusson EM and Arkin PA (1993) A global view of large-scale precipitation variability. *Journal of Climate* 6(8): 1495–1522.
- Ray RL and Jacobs JM (2007) Relationships among remotely sensed soil moisture, precipitation and landslide events. *Natural Hazards* 43: 211–222.
- Reheis MC (2006) A 16-year record of eolian dust in Southern Nevada and California, USA: Controls on dust generation and accumulation. *Journal of Arid Environments* 67: 487–520.
- Reichle RH, Koster RD, Dong J, and Berg AA (2004) Global soil moisture from satellite observations, land surface models, and ground data: Implications for data assimilation. *Journal of Hydrometeorology* 5: 430–442.
- Reynolds RL, Yount JC, Reheis M, Goldstein H, Chavez P, Fulton R, et al. (2007) Dust emission from wet and dry playas in the Mojave Desert, USA. *Earth Surface Processes and Landforms* 32: 1811–1827.
- Richards PL and Kump LR (2003) Soil pore-water distributions and the temperature feedback of weathering in soils. *Geochimica et Cosmochimica Acta* 67(20): 3803–3815.
- Rind D (1982) The influence of ground moisture conditions in North America on summer climate as modeled in the GISS GCM. *Monthly Weather Review* 110: 1487–1494.
- Roberts J (2000): The influence of physical and physiological characteristics of vegetation on their hydrological response. *Hydrological Processes* 14: 2885–2901.
- Robichaud PR (2000) Fire effects on infiltration rates after prescribed fire in Northern Rocky Mountain forests, USA. *Journal of Hydrology* 231–232: 220–229.
- Robichaud PR and Hungerford RD (2000) Water repellency by laboratory burning of four northern Rocky Mountain forest soils. *Journal of Hydrology* 231–232: 207–219.
- Robinson DA, Campbell CS, Hopmans JW, Hornbuckle BK, Jones SB, Knight R, et al. (2008) Soil moisture measurement for ecological and hydrological watershed-scale observatories: A review. *Vadose Zone Journal* 7: 358–389.
- Robock A, Vinnikov KY, Srinivasan G, Entin JK, Hollinger SE, Speranskaya NA, et al. (2000) The Global Soil Moisture Data Bank. *Bulletin of the American Meteorological Society* 81(6): 1281–1299.
- Rosenberg NJ, Blad BL, and Verma SB (1983) *Microclimate: The Biological Environment*, second edition. New York: Wiley-Interscience, 528 pp.
- Santanello JA Jr, Friedl MA, and Ek MB (2007) Convective planetary boundary layer interactions with the land surface at diurnal time scales: Diagnostics and feedbacks. *Journal of Hydrometeorology* 8(5): 1082–1097.
- Sarre RD (1987) Aeolian sand transport. *Progress in Physical Geography* 11: 157–182.
- Sarre RD and Chancey CC (1990) Size segregation during Aeolian saltation on sand dunes. *Sedimentology* 37(2): 357–365.
- Schlosser CA and Milly PCD (2002) A model-based investigation of soil moisture predictability and associated climate predictability. *Journal of Hydrometeorology* 3: 483–501.
- Schmugge TJ, Kustas WP, Ritchie JC, Jackson TJ, and Rango A (2002) Remote sensing in hydrology. *Advances in Water Resources* 25: 1367–1385.
- Schneider N and Eugster W (2005) Historical land use changes and mesoscale summer climate on the Swiss plateau. *Journal of Geophysical Research* 110: 1–18.
- Schuerch P, Densmore AL, McArdeell BW, and Molnar P (2006) The influence of landsliding on sediment supply and channel change in a steep mountain catchment. *Geomorphology* 78: 222–235.
- Schwab M, Rieke-Zapp D, Schneider H, Liniger M, and Schlunegger F (2008) Landsliding and sediment flux in the Central Swiss Alps: A photogrammetric study of the Schimbrig landslide, Entlebuch. *Geomorphology* 97: 392–406.
- Seeger M, Errea MP, Beguería S, Arnáez J, Martí C, and García-Ruiz JM (2004) Catchment soil moisture and rainfall characteristics as determinant factors for discharge/suspended sediment hysteretic loops in a small headwater catchment in the Spanish Pyrenees. *Journal of Hydrology* 288: 299–311.
- Segal M and Arritt RW (1992) Nonclassical mesoscale circulations caused by surface sensible heat-flux gradients. *Bulletin of American Meteorological Society* 73: 1593–1604.
- Sharma ML, Gander GA, and Hunt GC (1980) Spatial variability of infiltration in a watershed. *Journal of Hydrology* 45: 101–122.
- Sherman DJ (1990) Evaluation of aeolian sand transport equations using intertidal zone measurements, Staunton-Sands, England – discussion. *Sedimentology* 37(2): 385–388.
- Sherman DJ, Jackson DWT, Namikas SL, and Wang J (1998) Wind-blown sand on beaches: an evaluation of models. *Geomorphology* 22: 113–133.

- Shukla J and Mintz Y (1982) Influence of land-surface evapotranspiration on Earth's climate. *Science* 215: 1498–1501.
- Shur Y, Hinkel KM, and Nelson FE (2005) The transient layer: Implications for geocryology and global-change science. *Permafrost and Periglacial Processes* 16: 5–17.
- Sidle RC, Tsuboyama Y, Noguchi S, Hosoda I, Fujieda M, and Shimizu T (2000) Stormflow generation in steep forested headwaters: A linked hydrogeomorphic paradigm. *Hydrological Processes* 14: 369–385.
- Smalley IJ (1970) Cohesion of soil particles and the intrinsic resistance of simple soil. *Journal of Soil Science* 21(1): 154–161.
- Steiner AL, Pal JS, Giorgi F, Dickinson RE, and Chameides WL (2005) The coupling of the Common Land Model (CLM0) to a regional climate model (RegCM). *Theoretical and Applied Climatology* 82: 225–243.
- Stockwell J, Smith L, Jambor JL, and Beckie R (2006) The relationship between fluid flow and mineral weathering in heterogeneous unsaturated porous media: A physical and geochemical characterization of a waste-rock pile. *Applied Geochemistry* 21(8): 1347–1361.
- Stout JE (2001) Dust and environment in the Southern High Plains of North America. *Journal of Arid Environments* 47: 425–441.
- Svasek JN and Terwindt JHJ (1974) Measurements of sand transport by wind on a natural beach. *Sedimentology* 21: 311–322.
- Tang Q, Gao H, Lu H, and Lettenmaier DP (2009) Remote sensing: Hydrology. *Progress in Physical Geography* 33: 490–509.
- Teskey RO, Hinckley TM, and Grier CC (1984) Temperature induced changes in the water relations of *Abies amabilis* (Dougl.) forbes. *Plant Physiology* 74: 77–80.
- Thom AS (1972) Momentum, mass and heat exchange of vegetation. *Quarterly Journal of the Royal Meteorological Society* 98: 124–134.
- Thornthwaite CW (1948) An approach toward a rational classification of climate. *Geographical Review* 38(1): 55–94.
- Thornthwaite CW (1953) Topoclimatology. In: *Proceedings of the Toronto Meteorological Conference*, Toronto, Ontario, Canada, 227–232.
- Thornthwaite CW (1961) The task ahead. *Annals of the Association of American Geographers* 51(4): 345–356.
- Tsukamoto Y (1963) Storm discharge from an experimental watershed. *Journal of the Japanese Society of Forestry* 45: 186–190.
- Uchida T, Kosugi K, and Mizuyama T (1999) Runoff characteristics of pipeflow and effects of pipeflow on rainfall-runoff phenomena in a mountainous watershed. *Journal of Hydrology* 222: 18–36.
- Van Asch TWJ and Sukmantalya IN (1993) The modeling of soil slip erosion in the upper Komerang area, South Sumatra-Province, Indonesia. *Geografia Fisica e Dinamica Quaternaria* 16: 81–86.
- Van Asch TWJ, Buma J, and Van Beek LPH (1999) A view on some hydrological triggering systems in landslides. *Geomorphology* 30: 25–32.
- Van Asch TWJ, Van Beek LPH, and Bogaard TA (2009) The diversity in hydrological triggering systems of landslides. In: Picarelli L, Tommasi P, Urcinco G, and Versace P (eds) *Proceedings, 1st Italian Workshop on Landslides*, Naples, Italy, 151–156.
- Van Stan JT II and Levia DF (2010) Inter- and intraspecific variation of stemflow production from *Fagus grandifolia* Ehrh. (American beech) and *Liriodendron tulipifera* L. (yellow poplar) in relation to bark microrelief in the eastern United States. *Ecohydrology* 3: 11–19.
- Vera J, Mounzer O, Ruiz-Sanchez MC, Abrisqueta I, Tapia LM, and Abrisqueta JM (2009) Soil water balance trial involving capacitance and neutron probe measurements. *Agricultural Water Management* 96: 905–911.
- Vereecken H, Huisman JA, Bogena H, Vanderborght J, Vrugt JA, and Hopmans JW (2008) On the value of soil moisture measurements in vadose zone hydrology: A review. *Water Resources Research* 44: W00D06. doi: 10.1029/2008WR006829.
- Verstraeten WW, Veroustraete F, and Feyen J (2008) Assessment of evapotranspiration and soil moisture content across different scales of observation. *Sensors* 8: 70–117.
- Vertessy RA and Elsenbeer H (1999) Distributed modeling of storm flow generation in an Amazonian rain forest catchment: Effects of model parameterization. *Water Resources Research* 35: 2173–2187.
- Wagner W, Blöschl G, Pampaloni P, Calvet J-C, Bizzarri B, Wigneron J-P, and Kerr Y (2007a) Operational readiness of microwave remote sensing of soil moisture for hydrologic applications. *Nordic Hydrology* 38: 1–20.
- Wagner W, Naeimi V, Scipal K, de Jeu R, and Martínez-Fernández J (2007b) Soil moisture from operational

- meteorological satellites. *Hydrogeology Journal* 15: 121–131.
- Walker J and Rowntree PR (1977) The effect of soil moisture on circulation and rainfall in a tropical model. *Quarterly Journal of the Royal Meteorological Society* 103: 29–46.
- Walsh SJ, Butler DR and Malanson GP (1998) An overview of scale, pattern, process relationships in geomorphology: A remote sensing and GIS perspective. *Geomorphology* 21: 183–205.
- Washburn AL (1980) *Geocryology: A Survey of Periglacial Processes and Environments*. New York: Wiley, 406 pp.
- Western AW and Bloschl G (1999) On the spatial scaling of soil moisture. *Journal of Hydrology* 217: 203–223.
- Western AW, Grayson RB, and Bloschl G (2002) Scaling of soil moisture: A hydrologic perspective. *Annual Review of Earth and Planetary Science* 30: 149–180.
- Wieczorek GF and Glade T (2005) Climatic factors influencing occurrence of debris flows. In: Jakob M and Hungr O (eds) *Debris-flow Hazards and Related Phenomena*. Berlin: Springer-Praxis, 325–362.
- Wiggs GFS, Atherton RJ, and Baird AJ (2004) Thresholds of aeolian sand transport: Establishing suitable values. *Sedimentology* 51(1): 95–108.
- Wilhite DA (1997) Responding to drought: Common threads from the past, visions for the future. *Journal of the American Water Resources Association* 33(5): 951–959.
- Willmott CJ and Feddema JJ (1992) A more rational climate moisture index. *The Professional Geographer* 44(1): 84–87.
- Wolfe S and Nickling WG (1993) The protective role of sparse vegetation in wind erosion. *Progress in Physical Geography* 17(1): 50–68.
- World Meteorological Organization (WMO) (2008) *Future Climate Change Research and Observations: GCOS, WCRP and IGBP Learning from the IPCC Fourth Assessment Report*. WMO/TD 1418, GCOS-117, WCRP-127, and IGBP Report 58. Geneva: World Meteorological Organization.
- Wu H and Li Z-L (2009) Scale issues in remote sensing: A review on analysis, processing and modeling. *Sensors* 9: 1768–1793.
- Wu R and Kinter JL III (2009) Analysis of the relationship of U.S. droughts with SST and soil moisture distinguishing the time scales of droughts. *Journal of Climate* 22(17): 4520–4538.
- Yang Y and Davidson-Arnott R (2005) Rapid measurement of surface moisture content on a beach. *Journal of Coastal Research* 21(3): 447–452.
- Yeh TC, Wetherald RT, and Manabe S (1984) The effect of soil moisture on the short-term climate and hydrology change: A numerical experiment. *Monthly Weather Review* 112: 474–490.
- Zhang D and Anthes RA (1982) A high-resolution model of the planetary boundary-layer – sensitivity tests and comparisons with SESAME-79 data. *Journal of Applied Meteorology* 21: 1594–1609.
- Ziegler AD, Sutherland RA, and Giambelluca TW (2001) Acceleration of Horton overland flow and erosion by footpaths in an upland agricultural watershed in northern Thailand. *Geomorphology* 41: 249–262.