

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

7-31-2020

First record of *Planchonia stentae* (Brain, 1920) (Hemiptera: Coccomorpha: Asterolecaniidae) on *Asclepias curassavica* Linnaeus, 1753 (Gentianales: Asclepiadaceae) in Mexico, with observations on parasitic Encyrtidae

Hipolito Cortez-Madrigal

Arturo Ramírez-Cruz

Christof F. Stumpf

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0782

First record of *Planchonia stentae* (Brain, 1920)
(Hemiptera: Coccomorpha: Asterolecaniidae) on *Asclepias curassavica*
Linnaeus, 1753 (Gentianales: Asclepiadaceae) in Mexico, with
observations on parasitic Encyrtidae

Hipolito Cortez-Madrigal

Instituto Politécnico Nacional. Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional,
Unidad Michoacán. Michoacán. México

Arturo Ramírez-Cruz

Instituto Politécnico Nacional. Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional,
Unidad Michoacán. Michoacán. México

Christof F. Stumpf

Louisiana State University at Alexandria. Department of Biological Sciences. 8100 Hwy 71 South. Alexandria,
LA 71302-9121. USA

Date of issue: July 31, 2020

Hipolito Cortez-Madrigal, Arturo Ramírez-Cruz and Christof F. Stumpf
First record of *Planchonia stentae* (Brain, 1920) (Hemiptera: Coccothraupidae: Asterolecaniidae) on *Asclepias curassavica* Linnaeus, 1753 (Gentianales: Asclepiadaceae) in Mexico, with observations on parasitic Encyrtidae
Insecta Mundi 0782: 1–6

ZooBank Registered: urn:lsid:zoobank.org:pub:4C6D774F-9D53-4E0C-9729-E45FD5A7D28F

Published in 2020 by

Center for Systematic Entomology, Inc.

P.O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, insectamundi@gmail.com

Head Layout Editor: Robert G. Forsyth

Editorial Board: J. H. Frank, M. J. Paulsen

Founding Editors: Ross H. Arnett, Jr., Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agri-Food Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Robert G. Forsyth

First record of *Planchonia stentae* (Brain, 1920) (Hemiptera: Coccoomorpha: Asterolecaniidae) on *Asclepias curassavica* Linnaeus, 1753 (Gentianales: Asclepiadaceae) in Mexico, with observations on parasitic Encyrtidae

Hipolito Cortez-Madrigal

Instituto Politécnico Nacional. Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Unidad Michoacán. Michoacán. México. hcortezm@ipn.mx

Arturo Ramírez-Cruz

Instituto Politécnico Nacional. Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Unidad Michoacán. Michoacán. México. aramirez@ipn.mx

Christof F. Stumpf

Louisiana State University at Alexandria. Department of Biological Sciences. 8100 Hwy 71 South. Alexandria, LA 71302-9121. USA. cstumpf@lsua.edu

Abstract. In April 2018, specimens of an asterolecaniid were collected on infested wild plants of *Asclepias curassavica* Linnaeus, 1753 (Gentianales: Asclepiadaceae) in Jiquilpan, Michoacan, Mexico. The collected specimens were identified as *Planchonia stentae* (Brain, 1920) (Hemiptera: Asterolecaniidae). In this paper, we record for the first time the presence of *P. stentae* infesting *A. curassavica* in Mexico.

Key words. Parasitoid, pit scales, scarlet milkweed.

Introduction

Members of the family Asterolecaniidae (pit scales) are phytophagous insects that constitute one of the largest and more economically important scale insect taxa worldwide, with its members occurring in almost all zoogeographical regions of the world (Stumpf and Lambdin 1999, 2006). Females of many species of pit scales produce open galls, in which they suck plant sap (Stumpf and Lambdin 2006). Currently, this family consists of 25 genera and 247 species worldwide (García Morales et al. 2016).

One of the economically most important taxa of Asterolecaniidae includes the genus *Planchonia* Signoret, 1870 (Stumpf and Lambdin 2006), which contains ten species worldwide (García Morales et al. 2016). Specifically, the species *Planchonia stentae* (Brain, 1920) is significant due to its wide range of host plants, extensive distribution, and relative ability to be transported through movement of host plant material (Stumpf and Lambdin 2000, 2006).

Materials and Methods

Since 2012, a few scale insects have been observed on wild plants of *Asclepias curassavica* Linnaeus, 1753 (Asclepiadaceae) in the Cienega of Chapala region, but had not been collected. Adult female Asterolecaniidae infesting *A. curassavica* were first collected in April 2018 in Jiquilpan, Michoacan, Mexico (19°59'10.3"N, 102°42'37.6"W; 1560 meters above sea level) (Fig. 1A). Scale insect specimens were separated from the plants and preserved in 70% alcohol; subsequently, permanent slide-mounts of the adult females were prepared for species identification according to De Haro and Claps (1995). Samples of slide-mounted specimens were deposited in the Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional (CIIDIR), Unidad Michoacán, Instituto Politécnico Nacional, México, and alcohol-preserved specimens were deposited in the Entomological Collection of Louisiana State University at Alexandria, Alexandria, Louisiana, USA.

Results and Discussion

The female insect specimens were identified as *P. stentae* (Fig. 1B). We present the first official record for the presence of *P. stentae* in Mexico. This species is considered a major pest in Florida (Stumpf and Lambdin 2000), and also has the potential to interfere with Integrated Pest Management in Mexico. Although the presence of this species had already been recorded on *Echeveria* sp. (Crassulaceae) intercepted in Korea when imported from Mexico (Ji and Suh 2012; Suh et al. 2013), until identification its establishment in Mexico was not corroborated. Our findings also represent the first record of this species infesting *A. curassavica* and the family Asclepiadaceae in Mexico. So far, *P. stentae* has been recorded on a variety of host plants in South Africa (Brain 1920); Kenya, South Africa, USA (Russell 1941); Sao Tomé, South Africa, Zimbabwe (Giliomee 1966); USA (Gill 1993); Colombia (Stumpf and Lambdin 2000, 2006; Kondo 2001); Kenya, Puerto Rico, South Africa, USA (Stumpf and Lambdin 2000, 2006); Martinique (Germain and Grassart 2004; Matile-Ferrero and Étienne 2006); South Africa (Giliomee and Kozár 2008); Great Britain (Malumphy 2009); Mexico (Ji and Suh 2012; Suh et al. 2013) and Turks and Caicos Islands (Malumphy and Reid 2017). *Planchonia stentae* is the only pit scale species recorded from the host plant genus *Asclepias* (Stumpf and Lambdin 2006).

Scale insect taxonomy relies on the morphology of adult females. The genus *Planchonia* possesses fused, anvil-shaped anal plates, an arched plate, and an anal ring with six setae (Fig. 1C). Large 8-shaped pores (Fig. 1D) occur only on the dorsum and are found in marginal bands and in median swirls and bands. Rows of simple disk pores are present on dorsum and venter (Fig. 2) (Stumpf and Lambdin 2006). *Planchonia stentae* possesses three to four transverse multilocular pore bands ventrally around the vulva, with locular plates of multilocular pores having fringed edges (Fig. 1E), and ventral simple disk pores are only present in a submarginal row (Fig. 2) (Stumpf and Lambdin 2006). Morphologically, this species could be mistaken for *Planchonia arabis* Signoret, 1877, but *P. arabis* has at least five multilocular pore bands on the venter, the edges of locular plates are not fringed, and ventral simple disk pores are found in both submarginal and transverse rows (Stumpf and Lambdin 2006).

During the process of preparing specimens for slide-mounting, between two to four fully-developed adult parasitic wasps in the family Encyrtidae were found inside of some of the collected adult females of *P. stentae*. While *P. stentae* represents a new record of a phytophagous insect on *A. curassavica* in Mexico, the population density of this pit scale appears to be relatively low. We conclude that the polyembryonic encyrtid wasp found in this study could be an important population regulator of *P. stentae* populations in the Cienega de Chapala region. This finding is relevant, because there are no previous records of parasitoids attacking *P. stentae* (Stumpf and Lambdin 2006). The presence of parasitoids on *P. stentae* could be important for the management of agricultural pests, because *A. curassavica* hosts a large number of phytophagous insects, such as *Danaus plexippus* Linnaeus, 1758 (Lepidoptera: Nymphalidae) and *Aphis nerii* Boyer de Fonscolombe, 1841 (Hemiptera: Aphididae). Although *A. nerii* primarily feeds on the plant family Asclepiadaceae, this aphid can also colonize other plant families (Blackman and Eastop 1984) and can vector viral diseases, e.g., papaya ringspot virus (Mora-Aguilera et al. 1993). Nevertheless, the wide range of natural enemies that phytophagous insects associated with *A. curassavica* can support could help regulate pests in some agroecosystems as has been documented in citrus (Jacas and Urbaneja 2010; Godoy-Ceja and Cortez-Madrigal 2018). Therefore, the integration of *A. curassavica* as companion plant in agroecosystems has been suggested (Peña-Martínez et al. 2001; Cortez-Madrigal et al. 2016). Parasitoids controlling *P. stentae* on *A. curassavica* are valuable allies in this effort.

Future studies should be conducted in search of other host plants of *P. stentae* in Mexico, because the literature mentions about 40 host plants in Florida alone (Stumpf and Lambdin 2000), several of them present in Mexico. At the same time, parasitoid species involved in biological control of this pit scale should be identified. Our future goal is to estimate the pest potential of *P. stentae* in Mexico and to determine its biology.

Acknowledgments

We would like to thank Dr. S. N. Myartseva, Universidad Autónoma de Tamaulipas, México, for the identification of the family Encyrtidae, and Dr. C. Malumphy, Fera Science Ltd., and Dr. J. H. Giliomee, Stellenbosch University, for valuable literature references.

Special thanks go to the two reviewers, Greg Hodges, Florida Dept. of Agriculture and Consumer Services, Gainesville, FL, and Gary Miller, Systematic Entomology Laboratory, Beltsville, MD who were extremely helpful in improving this paper.

Literature Cited

- Blackman, R. L., and V. F. Eastop. 1984.** Aphids on the world's crops. An identification guide. John Wiley & Sons; London, UK. 413 p.
- Brain, C. K. 1920.** The Coccidae of South Africa-IV. Bulletin of Entomological Research 10(2): 95–128.
- Cortez-Madrigal, H., F. García-González, A. Guzmán-Larralde, J. A. Acuña-Soto, and G. Otero-Colina. 2016.** Conserving phytophagous arthropods to conserve natural enemies: *Asclepias curassavica* as the model. Southwestern Entomologist 41(3): 681–693.
- De Haro, M. E., and L. E. Claps. 1995.** Conociendo nuestra fauna III. Familia Dactylopiidae. (Insecta: Homoptera). Morfología, biología e importancia económica. Serie monográfica y didáctica N° 19. Instituto Superior de Entomología. INSUE. Facultad de Ciencias Naturales e Instituto Miguel Lillo. Universidad Nacional de Tucumán; Tucumán, Argentina. 20 p.
- García Morales, M., B. D. Denno, D. R. Miller, G. L. Miller, Y. Ben-Dov, and N. B. Hardy. 2016.** ScaleNet. Available at <http://scalenet.info>. (Last accessed April 2, 2020.)
- Germain, J. F., and L. Grassart. 2004.** First occurrence into Martinique (French West Indies) one pit scale and four mealybugs (Abstract only). p. 330. In: L. B. Erkilic and M. B. Kaydan (eds.). Proceeding of the X International Symposium on Scale Insect Studies. 19-23 April 2004. Plant Protection Research Institute; Adana, Turkey. 408 p.
- Giliomee, J. H. 1966.** A list of South African scale insects (Homoptera: Coccoidea) in the collection of the British Museum (Natural History) with information on their hosts plants and distribution. Annale Universiteit Van Stellenbosch. 41. Serie A 8: 412–428.
- Giliomee, J. H., and F. Kozár. 2008.** Review of the family Asterolecaniidae (Hemiptera: Coccoidea) in South Africa, with the description of a new species. African Entomology 16(2): 276–286.
- Gill, R. J. 1993.** The scale insects of California. Part 2. The minor families (Homoptera: Coccoidea). California Department of Food and Agriculture; Sacramento, California. 241 p.
- Godoy-Ceja, C. A., and H. Cortez-Madrigal. 2018.** Potencial de *Asclepias curassavica* L. (Apocynaceae) en el control biológico de plagas. Revista Mexicana de Ciencias Agrícolas 9(2): 303–315.
- Jacas, J. A., and A. Urbaneja. 2010.** Biological control in citrus in Spain: from classical to conservation biological control. p. 61–72. In: A. Ciancio and K. G. Mukerji (eds.). Integrated management of arthropod pests and insect borne diseases. Springer; London. 366 p.
- Ji, J., and S. J. Suh. 2012.** A list of scale insects (Hemiptera: Coccoidea) intercepted in quarantine on imported succulent plants in Korea 2006–2010. Insecta Mundi 0272: 1–5.
- Kondo, T. 2001.** Las cochinillas de Colombia (Hemiptera: Coccoidea). Biota Colombiana 2(1): 31–48.
- Malumphy, C. 2009.** A brief review of pit scales (Hemiptera: Asterolecaniidae) recorded in Britain. Cecidology 24: 46–49, 64.
- Malumphy, C., and S. Reid. 2017.** Improving bio-security in the United Kingdom overseas territories: Identification service for invasive alien invertebrate plant pests. Food and Environment Research Agency; UK. 23 p.
- Matile-Ferrero, D., and J. Étienne. 2006.** Cochenilles des antilles françaises et de quelques autres îles des Caraïbes (Hemiptera, Coccoidea). Revue Française d'Entomologie (N.S.) 28(4): 161–190.
- Mora-Aguilera, G., D. Nieto-Angel, D. Teliz, and C. L. Campbell. 1993.** Development of a prediction model for papaya ringspot in Veracruz, Mexico. Plant Disease 77(12): 1205–1211.
- Peña-Martínez, M. R., J. R. Lomelli, G. L. A. Trejo, and N. Villegas J. 2001.** Monitoreo de Afdos y Afdófagos de Importancia Agrícola. Escuela Nacional de Ciencias Biológicas. Instituto Politécnico Nacional; Mexico, D. F. 89 p.
- Russell, L. M. 1941.** A classification of the scale insect genus *Asterolecanium*. United States Department of Agriculture. Miscellaneous Publication No. 424. Washington, D.C. 322 p.
- Stumpf, C. F., and P. L. Lambdin. 1999.** A comparison of the pit scale genera of North and South America (Hemiptera: Coccoidea: Asterolecaniidae). Entomologica, Bari 33: 153–156.

- Stumpf, C. F., and P. L. Lambdin. 2000.** Distribution and known host records for *Planchonia stentae* (Hemiptera: Coccoidea: Asterolecaniidae). *Florida Entomologist* 83(3): 368–369.
- Stumpf, C. F., and P. L. Lambdin. 2006.** Pit scales (Sternorrhyncha: Coccoidea) of North and South America. Tennessee Agricultural Experiment Station. University of Tennessee Institute of Agriculture; Knoxville, Tennessee. 231 p.
- Suh, S. J., H. M. Yu, and K. J. Hong. 2013.** List of intercepted scale insects at Korean ports of entry and potential invasive species of scale insects to Korea (Hemiptera: Coccoidea). *Korean Journal of Applied Entomology* 52(2): 141–160.

Received May 16, 2020; accepted June 20, 2020.

Review editor Aline Barcellos.




Figure 1. Adult females of *Planchonia stentae*. **A)** Adult females (arrows) on the stem of *Asclepias curassavica*. **B)** Whole body. **C)** Anal ring (r) with setae (s), arched plate (a) and anal plate (asterisk). **D)** Lateral abdominal margin with 8-shaped pores in double or triple row (arrows). **E)** Multilocular pore bands (arrows), with locular plates of multilocular pores with fringed edges (insert); anal ring (r). Scales: B = 0.2 mm. C = 50 μ m. D = 20 μ m. E = 50 μ m. Insert = 6.0 μ m.




Figure 2. Illustration of mature adult female of *Planchonia stentae* with ventral characters on right side and dorsal characters on left side. Size: 1–2.5 mm. **a)** Body with mouthparts in upper third. **b)** Dorsal 8-shaped pores. **c)** Dorsal simple disk pores. **d)** Tubular ducts. **e)** Anal lobes with arched plate, anal plates, and anal ring (see Fig. 1C-1E for details of structures). **f)** Antennae. **g)** Spiracles. **h)** Ventral 8-shaped pores. **i)** Ventral simple disk pores. **j)** Quinquelocular pores. **k)** Multilocular pores. **l–m)** Multilocular pores with fringed edges. **n)** Setae. For more detailed taxonomic data, see Stumpf and Lambdin (2006). Illustration taken with permission from Stumpf and Lambdin (2006). © C. Stumpf.