

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

2013

Detection of Avian Influenza Viruses in Wild Waterbirds in the Rift Valley of Kenya Using Fecal Sampling

Victor O. Ofula

U.S. Army Medical Research Unit–Kenya

Alan B. Franklin

National Wildlife Research Center, alan.b.franklin@aphis.usda.gov

J. Jeffrey Root

USDA/APHIS/WS National Wildlife Research Center, jeff.root@aphis.usda.gov

Heather J. Sullivan

USDA/APHIS/WS National Wildlife Research Center, Heather.Sullivan@aphis.usda.gov

Patrick Gichuki

National Museums of Kenya

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc

 Part of the [Life Sciences Commons](#)

Ofula, Victor O.; Franklin, Alan B.; Root, J. Jeffrey; Sullivan, Heather J.; Gichuki, Patrick; Makio, Albina; Bulimo, Wallace; Abong'o, Bernard O.; Muchai, Muchane; and Schnabel, David, "Detection of Avian Influenza Viruses in Wild Waterbirds in the Rift Valley of Kenya Using Fecal Sampling" (2013). *USDA National Wildlife Research Center - Staff Publications*. 1566.
https://digitalcommons.unl.edu/icwdm_usdanwrc/1566

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Victor O. Ofula, Alan B. Franklin, J. Jeffrey Root, Heather J. Sullivan, Patrick Gichuki, Albina Makio, Wallace Bulimo, Bernard O. Abong'o, Muchane Muchai, and David Schnabel

Detection of Avian Influenza Viruses in Wild Waterbirds in the Rift Valley of Kenya Using Fecal Sampling

Victor O. Ofula,¹ Alan B. Franklin,² J. Jeffrey Root,² Heather J. Sullivan,² Patrick Gichuki,³ Albina Makio,¹ Wallace Bulimo,¹ Bernard O. Abong'o,⁴ Muchane Muchai,³ and David Schnabel¹

Abstract

Highly pathogenic avian influenza virus A/H5N1 has been reported in 11 African countries. Migratory waterbirds have the potential of introducing A/H5N1 into east Africa through the Rift Valley of Kenya. We present the results of a wild bird surveillance system for A/H5N1 and other avian influenza viruses based on avian fecal sampling in Kenya. We collected 2630 fecal samples in 2008. Viral RNA was extracted from pools of 3–5 fecal samples and analyzed for presence of avian influenza virus RNA by real-time RT-PCR. Twelve (2.3%) of the 516 sample pools were positive for avian influenza virus RNA, 2 of which were subtyped as H4N6 viruses. This is the first report of avian influenza virus in wild birds in Kenya. This study demonstrates the success of this approach in detecting avian influenza virus in wild birds and represents an efficient surveillance system for avian influenza virus in regions with limited resources.

Key Words: Africa—Aves—Avian influenza virus—Feces—Nondomestic animals—Waterfowl.

Introduction

INFLUENZA A VIRUSES ARE MEMBERS of the viral family *Orthomyxoviridae* and the genus *Influenzavirus A*. Influenza A viral envelopes contain two major glycoproteins—hemagglutinin (HA) and neuraminidase (NA) (Swayne and Halvorson 2003), which play key roles during cellular infection. Different HA/NA combinations allow viral subtype discrimination (Atkinson et al. 2009). Subtypes not usually isolated from humans, swine, or horses are referred to as avian influenza viruses (AIV). Currently, 16 HA and 9 NA subtypes of influenza A virus have been identified from wild birds (Fouchier et al. 2004, Olsen et al. 2006) and are considered to be AIV. Most influenza A viruses originate from wild waterfowl and shorebirds, which are the primary reservoirs for these viruses (Webster et al. 1992). However, these viruses can also spread to other wild bird species, wild mammals, domestic poultry, swine, and humans. Although AIV rarely causes morbidity and mortality in wild birds, two levels of AIV pathogenicity have been described in poultry—low pathogenic AIV (LPAIV) and highly pathogenic AIV (HPAIV) (Alexander 2000). LPAIV in wild birds can evolve into HPAIV once introduced into poultry (Horimoto et al. 1995, Banks et al. 2001, Duan et al. 2007). Thus, LPAIV circulating in wild birds pose an indirect threat to poultry and

humans, and it is critical to understand the ecology of LPAIV to prevent disease in these populations.

Following the large-scale outbreak of AIV strain A/H5N1 in migratory waterfowl at Qinghai Lake, China (Li et al. 2004), A/H5N1 virus subsequently spread through Central Asia, Europe, the Middle East, and into Africa (Wang et al. 2008). In Africa, A/H5N1 virus was first detected from 6 poultry outbreaks during January, 2006, in Nigeria (Joannis et al. 2006), with at least 2 of the outbreaks from viruses closely related to those isolated from wild waterfowl in Europe, suggesting possible transmission of the virus from migratory waterfowl (Ducatez et al. 2006). Since the initial outbreaks in Nigeria, there have been 1765 records of A/H5N1 in poultry to date in Africa, which included Egypt ($n=1423$), Nigeria ($n=296$), Sudan ($n=21$), Ghana ($n=6$), Benin ($n=5$), Ivory Coast ($n=5$), Burkina Faso ($n=4$), Niger ($n=2$), Cameroon ($n=1$), Djibouti ($n=1$), and South Africa ($n=1$) (WAHID 2005–2009a). In March, 2006, Egypt confirmed its first human case associated with this virus. As of June 22, 2011, there have been 155 human cases of A/H5N1, with 54 deaths recorded in Africa (WHO 2011). Over 96% of these cases and deaths occurred in Egypt. The spread of A/H5N1 virus across Africa raises serious concerns regarding the sustainability of the poultry sector and public health (Cattoli et al. 2009).

¹U.S. Army Medical Research Unit—Kenya, Nairobi, Kenya.

²National Wildlife Research Center, Fort Collins, Colorado.

³National Museums of Kenya, Nairobi, Kenya.

⁴Great Lakes University of Kisumu, Kisumu, Kenya.

Although considerable attention has focused on controlling A/H5N1 in Asia, Africa has largely been overlooked. For example, surveillance for A/H5N1 in Africa is sporadic relative to other continents, but the risk of human infection from A/H5N1 in Africa is estimated to be similar to that for Asia (WHO 2005). Several factors suggest that Africa may be an area of future concern. First, human population densities in Africa are increasing exponentially and are expected to increase from 26 people/km² to 60 people/km² in the next 50 years (Cohen 2003). Second, human contact with poultry is very high in Africa. Total poultry numbers in Africa are estimated at 1.1 billion (with 1.5 village fowl per person), and most poultry are free-ranging or in backyard systems with very low, if any, biosecurity (Guèye 1998, Sonaiya 2007).

Wild birds have been implicated in the movement of A/H5N1 virus within and across continents (Chen et al. 2006, Olsen et al. 2006). In Africa, A/H5N1 has been documented in wild birds during at least 8 outbreaks in Egypt, Ivory Coast, and Nigeria, usually in conjunction with poultry outbreaks (WAHID 2005–2009b). A/H5N1 has the potential to be widely introduced into Africa by wild birds migrating from Europe, where A/H5N1 is already present (Gaidet et al. 2007b), and circulated within Africa from countries already experiencing outbreaks. Because they have the potential to move A/H5N1 and other highly pathogenic strains of AIV, systematic AIV surveillance in wild birds may serve as an early warning for the introduction of AIV strains of concern in an area (Deliberto et al. 2009). Although surveillance for the presence of A/H5N1a has been conducted in wild birds in Africa (Gaidet et al. 2007a, Gaidet et al. 2007b, Nasirwa 2006), A/H5N1 has not been detected to date in wild bird populations. However, these surveys have provided valuable information on the presence of LPAIV in wild birds in Africa.

In Kenya, wild birds migrate from Eurasia and neighboring countries, such as Sudan and Ethiopia (Lewis and Pomeroy 1989, Scott and Rose 1996), which have experienced A/H5N1 outbreaks amongst their domestic poultry. In particular, the Rift Valley of Kenya serves as a major flyway for migrating palaeartic ducks, which stop over in this region during both their northern and southern migrations (Gichuki and Gichuki 1992). Consequently, Kenya remains at high risk for potential A/H5N1 introductions.

The objective of our study was to design and test a surveillance program that could detect AIV in Kenya using freshly deposited wild bird feces. Avian fecal sampling is an efficient way to monitor for the presence of AIV in areas used by wild birds, especially in countries, such as Kenya, where hunter-killed samples are not available because waterfowl hunting is prohibited. Environmental sampling is advantageous because capture and handling of wild birds is not required and sampling can be conducted in the field with minimal equipment and training. The basis for this approach is that wild birds shed infectious AIV (including A/H5N1) in their feces (Ito et al. 1995, Brown et al. 2006), which can then persist in water for several months (Stallknecht et al. 1990b, Brown et al. 2007), dependent on various environmental conditions.

Materials and Methods

We targeted sampling efforts by identifying waterbird species to be of primary concern for potential transmission of A/H5N1 into Kenya and areas where these species might

congregate. Waterbird species in Kenya were classified as either primary or secondary target species. Primary target species were palaeartic migratory waterfowl from Europe and Asia that had the potential to introduce A/H5N1 into Kenya from Europe or Asia where outbreaks of A/H5N1 occur. These included northern pintail (*Anas acuta*), garganey (*Anas querquedula*), northern shovelers (*Anas clypeata*), and Eurasian wigeon (*Anas penelope*) (Brown et al. 1982, Perennou et al. 1994). Secondary target species included afro-tropical waterfowl that breed and winter in Africa, such as storks and egrets (Ciconiformes), gulls (Charadriiformes), pelicans (Peliconiformes), and flamingoes (Phoenicopteriformes). The Rift Valley of Kenya was chosen as our study site because it is a major wintering area for palaeartic waterfowl migrating from Eurasia and is also a major flyway for waterbirds migrating to southern and western Africa (Pearson and Meadows 1992).

In developing a sampling design, we prioritized lakes in the Rift Valley by the types and numbers of primary and secondary target species occurring in those areas, with an emphasis on lakes where migratory waterfowl from Europe and Asia winter in Kenya (Fig. 1). Although outside the Rift Valley, the Dandora Sewage Ponds near Nairobi were also sampled because of their importance as a waterfowl congregation area in Kenya (Pearson and Meadows 1992, Nasirwa 2006). Each lake within the Rift Valley was ranked and weighted based on: (1) the maximum numbers of palaeartic waterfowl observed wintering at the site (Scott and Rose 1996, Owino et al. 2001, Owino et al. 2002, Wetlands International 2007), where sites were scored by the ranking of maximum numbers observed (6 being the highest in importance and 1 being the lowest); (2) maximum numbers of afro-tropical waterfowl observed at the site (Scott and Rose 1996, Owino et al. 2001, Owino et al. 2002, Wetlands International 2007), where sites were scored similarly to palaeartic waterfowl; (3) water type, where a score of 2 was assigned if the lake was fresh and 0 if the lake was saline (Lake Nakuru was assigned a score of 1 because the area contained elements of both saline and fresh water); and (4) human density, where a score of 1 was assigned for low density, 2 for medium density, and 3 for high density (e.g., Nairobi). Scores for palaeartic waterfowl were multiplied by 2 to weight them twice as heavily as afro-tropical waterfowl and the other criteria because of their importance in potentially introducing A/H5N1 into Kenya. For each site, scores were summed and then normalized to obtain weights, which represented proportions of the total samples (Table 1).

We chose a total sample size of 4500 samples to be collected, based on logistical constraints. We then calculated the optimal number of samples to be collected at each lake by multiplying the total number of samples to be collected by the weight for each location. In order of ranked importance, the areas sampled in this study were Dandora Sewage Ponds, Lake Naivasha, Lake Oloiden, Lake Bogoria, Lake Nakuru (including the adjacent Nakuru sewage ponds), Lake Elementeita, and Lake Baringo (Table 1).

Samples were collected in October, 2008, which is normally when migratory avian species move through Kenya (Owino et al. 2001, Owino et al. 2002). At each lake, flocks of target species were identified and then approached to collect freshly deposited fecal samples using Dacron swabs, which were stored in vials containing BA-1 virus transport medium with antibiotics, as previously described in Root et al. (2010).




FIG. 1. Map of Kenya showing the locations of water bodies in the Rift Valley where fecal samples from waterbirds were collected.

Samples were stored on ice immediately after collection and then frozen in liquid nitrogen until laboratory analysis. The species of birds that was most predominant at the site of collection were documented and GPS readings taken for each sampling point.

Real-time RT-PCR was performed on all fecal swabs to test for evidence of the presence of viral RNA. Up to 5 samples from similar species at the same location were pooled for laboratory analysis, and viral RNA was extracted from these pools and analyzed by real-time RT-PCR (Spackman et al. 2002) for the presence of AIV. Calibrated controls with known viral titers (10^2 EID₅₀/mL – 10^5 EID₅₀/mL) were also analyzed with real-time RT-PCR to construct 4-point standard

curves to estimate the viral concentrations in each sample. Samples found to be positive for the matrix gene were further characterized by H5- and H7-specific real-time RT-PCR assays (Spackman et al. 2002), the former of which is known to detect A/H5N1. Samples positive for AIV were analyzed further using virus isolation in chicken embryos (Szretter et al. 2006) and virus isolates were then sent to the National Veterinary Services Laboratory (Ames, IA) for subtyping using the HA-inhibition and NA-inhibition tests. The Fisher exact test (Zar 1984) was used to test whether avian fecal samples collected at freshwater lakes had a higher proportion of AIV-positive samples than samples collected at alkaline lakes and whether avian fecal samples collected from flocks containing

TABLE 1. NUMBERS OF SAMPLES ALLOCATED TO EACH LAKE BASED ON A WEIGHTED SAMPLING DESIGN AND SAMPLES ACTUALLY COLLECTED AT SIX SITES IN THE RIFT VALLEY, KENYA

Site	Criteria				Score	Weight	Number (%) of samples	
	Palaearctic waterfowl	Afro-tropical waterfowl	Water quality	Human density			Allocated (n = 4500)	Actual (n = 2630)
Dandora Ponds	12	6	2	3	23	0.288	1296 (28.8%)	526 (20.0%)
Lake Naivasha	10	5	2	2	19	0.238	1068 (23.8%)	840 (31.9%)
Lake Nakuru	8	3	1	2	14	0.175	787 (17.5%)	691 (26.3%)
Lake Elmenteita	6	2	0	1	9	0.113	506 (11.3%)	167 (6.3%)
Lake Bogoria	4	4	0	1	9	0.113	506 (11.3%)	365 (13.9%)
Lake Baringo	2	1	2	1	6	0.073	337 (7.3%)	41 (1.6%)

species with afrotropical distributions were different than those collected from flocks with palaearctic distributions.

Ethical and scientific approval was obtained from the Kenya Medical Research Institute Ethical Review Committee and the National Museums of Kenya. No wild birds were disturbed or handled during the conduct of this study.

Results

A total of 2630 samples were collected in this study and tested as 516 pools by real-time RT-PCR (Table 2). Although we were unable to meet our target of 4500 samples, our effort was roughly distributed according to our sampling design (Table 1). Overall, 2.3% (n=12) of the sample pools were positive for AIV RNA but none were positive by real-time RT-PCR for H5 or H7 subtypes. Samples collected from freshwater lakes had a significantly higher percent of AIV-positive pools (3.74%, 95% confidence interval [CI]=1.66–5.81%; Fisher exact p=0.002) than those collected at alkaline lakes (0.00%). Overall, Lake Oloiden had the highest percent of avian fecal samples collected that yielded AIV-positive pools (Table 2).

Based on the Fisher exact test, samples associated with flocks dominated by waterfowl (Anseriformes) had a significantly higher percentage of AIV-positive pools (4.5%,

95% CI=1.8–7.3%; Fisher exact p=0.006) than from other taxonomic orders (Table 3). Flocks dominated by coots (Gruiformes) and pelicans and cormorants (Pelecaniformes) had the next highest percentage of AIV-positive pools (1.7%, 95% CI=0.0–4.0; Table 3). AIV was not found in flocks dominated by storks and egrets (Ciconiiformes), flamingos (Phoenicopteriformes), or shorebirds (Charadriiformes) (Table 3). Of the 12 positive sample pools detected by real-time RT-PCR, 2 were subtyped as H4N6 viruses. Both of these sample pools were collected from a flock of cape teal at Lake Oloiden.

Discussion

The purpose of this study was to design an economical surveillance system for AIV that could be adopted by Kenyan scientists and potentially by other developing countries. The overall percentage of positive samples that we found was low, but not dissimilar to results from other surveillance efforts (Olsen et al. 2006, Gaidet et al. 2007a, Deliberto et al. 2009). Although inhibitors may reduce the detectability of AIV in feces using PCR, McLean et al. (2007) noted that overall prevalence from fecal sampling was similar to the prevalences found from sampling live birds in PCR-based surveillance efforts. In addition, this surveillance effort is the first to report the presence of AIV in wild birds in Africa, despite previous surveillance efforts (Nasirwa 2006, Gaidet et al. 2007a, b). Of interest, AIV was detected at Dandora Sewage Ponds, a site that had been surveyed in the past but with no AIV detected (Nasirwa 2006). Thus, this study indicates that our surveillance system was sensitive and able to detect AIV. In addition, this system could also be used to focus live bird capture and sampling efforts for the detection of species-specific infections with AIV.

Our study indicates circulation of AIV in select water bodies in Kenya. Overall, we found that waterfowl appeared to be the most common shedders of AIV, but that other avian orders in Kenya, such as Gruiformes and Pelecaniformes, should be considered for further scrutiny. Although fecal samples could not be definitively ascribed to a given species, flocks where samples were collected were almost always composed of a single species. Therefore, this collection strategy offers at least a moderate amount of precision in conducting targeted pathogen surveillance in wild birds in Kenya. In addition, feces of certain species, such as Egyptian geese, flamingoes, and marabou storks, were sufficiently distinct in form to identify them at the species level.

TABLE 2. PERCENTAGE OF POOLS FROM AVIAN FECAL SAMPLES TESTING POSITIVE FOR AVIAN INFLUENZA VIRUS WITH 95% CONFIDENCE INTERVALS, AT SEVEN STUDY SITES IN KENYA

Site	Water type	Number of sample pools		% AIV+ (95% CI)
		AIV-	AIV+	
Lake Baringo	Fresh	9	0	0.0 (—)
Lake Bogoria	Alkaline	80	0	0.0 (—)
Dandora Sewage Ponds	Fresh	85	2	2.3 (0.0, 5.5)
Lake Elementaita	Alkaline	37	0	0.0 (—)
Lake Naivasha	Fresh	115	2	1.7 (0.0, 4.1)
Lake Nakuru	Alkaline/fresh	132	1	0.8 (0.0, 2.2)
Lake Oloiden	Fresh	46	7	13.2 (4.1, 22.3)
Total		504	12	2.3 (1.0, 3.6)

AIV, Avian influenza virus; CI, confidence interval.

TABLE 3. PERCENTAGE OF POOLS FROM AVIAN FECAL SAMPLES TESTING POSITIVE FOR AVIAN INFLUENZA VIRUS, WITH 95% CONFIDENCE INTERVALS, IN DOMINANT BIRD SPECIES SAMPLED USING FECAL SAMPLES IN KENYA

Dominant species	Distribution†	No. of sample pools		% AIV+	95% CI
		AIV–	AIV+		
Pelicaniformes (<i>n</i> = 86 pools)					
Great white pelican	Palaearctic	57	0	0.0	—
Great cormorant	Palaearctic	28	1	3.5	0.00, 10.09
Subtotal		85	1	1.2	0.00, 3.43
Ciconiiformes (<i>n</i> = 61 pools)					
Goliath heron	Afrotropical	1	0	0.0	—
Great white egret	Palaearctic	1	0	0.0	—
Little egret	Palaearctic	3	0	0.0	—
Yellow-billed stork	Afrotropical	20	0	0.0	—
Marabou stork	Afrotropical	10	0	0.0	—
Sacred ibis	Afrotropical	25	0	0.0	—
African spoonbill	Afrotropical	1	0	0.0	—
Subtotal		61	0	0.0	—
Phoenicopteriformes (<i>n</i> = 42 pools)					
Greater flamingo	Palaearctic	17	0	0.0	—
Lesser flamingo	Palaearctic	25	0	0.0	—
Subtotal		42	0	0.0	—
Anseriformes (<i>n</i> = 221 pools)					
White-faced whistling duck	Afrotropical	32	2	5.9	0.00, 23.79
Egyptian goose	Afrotropical	103	3	2.8	0.00, 5.99
Cape teal	Afrotropical	43	3	6.5	0.00, 13.66
Yellow-billed duck	Afrotropical	25	1	3.9	0.00, 11.24
Red-billed duck	Afrotropical	1	0	0.0	—
Hottentot teal	Afrotropical	7	1	12.5	0.00, 35.42
Subtotal		211	10	4.5	1.78, 7.27
Gruiformes (<i>n</i> = 35 pools)					
Red-knobbed coot	Afrotropical	34	1	2.9	0.00, 8.38
Charadriiformes (<i>n</i> = 71 pools)					
Black-winged stilt	Palaearctic	20	0	0.0	—
Blacksmith plover	Afrotropical	9	0	0.0	—
Spur-winged plover	Palaearctic	4	0	0.0	—
Kittlitz's plover	Afrotropical	8	0	0.0	—
Common sandpiper	Palaearctic	1	0	0.0	—
Ruff	Palaearctic	3	0	0.0	—
Gray-headed gull	Afrotropical	13	0	0.0	—
White-cheeked tern	Palaearctic	2	0	0.0	—
Whiskered tern	Palaearctic	11	0	0.0	—
Subtotal		71	0	0.0	—
Total (<i>n</i> = 516 pools)		504	12	2.3	1.03, 3.63

AIV, Avian influenza virus; CI, confidence interval.

Nonetheless, we conducted our analyses at the ordinal, rather than species, taxonomic level because of the uncertainty of assignment of species to samples.

Overall, we detected a relatively low percentage (2.3%) of pooled samples to be positive for AIV. However, the percentage of positive pools obtained is not surprising. For example, Brown and Stallknecht (2008) concluded that prevalence of AIV in North America decreases as waterfowl populations migrate, with prevalences as low 1–2% in wintering areas. There was a higher chance of detecting AIV from samples collected around freshwater bodies than alkaline water bodies. This finding was expected on the basis of previous work on AIV survival in water of different chemistry (Stallknecht et al. 1990a; Brown et al. 2007). These data can be used by the Kenyan disease surveillance community to target resources for animal and human surveil-

lance near freshwater lakes in the future. However, our sampling effort was low in some areas (*e.g.*, Lake Baringo) and lacked the power to detect AIV in those areas. Therefore, we suggest that future sampling be conducted at a larger scale to better approximate necessary sample sizes for future surveillance efforts.

To our knowledge, the AIV subtype H4N6 has not been reported in Africa, even though it is fairly common in Europe (Suss et al. 1994, Alexander 2007). In Europe, H4N6 has been found in garganey and mallards, both of which migrate through Kenya (Lewis and Pomeroy 1989). This provides evidence for the potential spread of AIV from Europe to Kenya, with implications for the possible introduction of pathogenic A/H5N1 virus into Kenya. This potential is being currently explored through genetic sequencing of the H4N6 viruses we found.

Acknowledgments

We would like to acknowledge Samuel Owaka for his assistance in sample collections; Finley Osuna, Meshack Wadegu, Ben Kiawa, and Darcy Orahoo for assistance in laboratory testing; and Jeremy Ellis for his assistance with map development. We would also like to thank Dr. Rosemary Sang for providing laboratory space at the Kenya Medical Research Institute, Centre for Virus Research, Nairobi, Kenya. Funding for this work was provided by the U.S. Department of Defense, Armed Forces Health Surveillance Center, Division of GEIS Operations.

Material has been reviewed by the Walter Reed Army Institute of Research. There is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author, and are not to be construed as official, or as reflecting true views of the Department of the Army or the Department of Defense. Research was conducted in compliance with the Animal Welfare Act and other federal statutes and regulations relating to animals and experiments involving animals and adheres to principles stated in the *Guide for the Care and Use of Laboratory Animals*, NRC Publication, 2011 edition.

Author Disclosure Statement

No competing financial interests exist.

References

- Alexander DJ. A review of avian influenza in different bird species. *Vet Microbiol* 2000; 74:3–13.
- Alexander DJ. Summary of avian influenza activity in Europe, Asia, Africa, and Australasia, 2002–2006. *Avian Dis* 2007; 51:161–166.
- Atkinson W, Hamborsky J, McIntyre L, et al. *Epidemiology and prevention of vaccine-preventable diseases*. Washington DC, 2009.
- Banks J, Speidel ES, Moore E, et al. Changes in the haemagglutinin and the neuraminidase genes prior to the emergence of highly pathogenic H7N1 avian influenza viruses in Italy. *Arch Virol* 2001; 146:963–973.
- Brown JD, Stallknecht DE, Beck JR, et al. Susceptibility of North American ducks and gulls to H5N1 highly pathogenic avian influenza viruses. *Emerg Infect Dis* 2006; 12:1663–1670.
- Brown JD, Swayne DE, Cooper RJ, et al. Persistence of H5 and H7 avian influenza viruses in water. *Avian Dis* 2007; 51:285–289.
- Brown JD, Stallknecht DE. Wild bird surveillance for the avian influenza virus. *Methods Mol Biol* 2008; 436:85–97.
- Brown LH, Urban EK, Newman K. *The Birds of Africa*. New York: Academic Press, 1982.
- Cattoli G, Monne I, Fusaro A, et al. Highly pathogenic avian influenza virus subtype H5N1 in Africa: A comprehensive phylogenetic analysis and molecular characterization of isolates. *PLoS One* 2009; 4.
- Chen H, Li Y, Li Z, et al. Properties and dissemination of H5N1 viruses isolated during an influenza outbreak in migratory waterfowl in Western China. *J Virol* 2006; 80:5976–5983.
- Cohen JE. Human population: The next half century. *Science* 2003; 302:1172–1175.
- Deliberto TJ, Swafford SR, Nolte DL, et al. Surveillance for highly pathogenic avian influenza in wild birds in the USA. *Integr Zool* 2009; 4:426–439.
- Duan L, Campitelli L, Fan XH, et al. Characterization of low-pathogenic H5 subtype influenza viruses from Eurasia: Implications for the origin of highly pathogenic H5N1 viruses. *J Virol* 2007; 81:7529–7539.
- Ducatez MF, Olinger CM, Owoade AA, et al. Avian Flu: Multiple introductions of H5N1 in Nigeria. *Nature* 2006; 442:37.
- Fouchier RAM, Schneeberger PM, Rozendaal FW, et al. Avian influenza A virus (H7N7) associated with human conjunctivitis and a fatal case of acute respiratory distress syndrome. *Proc Natl Acad Sci USA* 2004; 101:1356–1361.
- Gaidet N, Dodman T, Caron A, et al. Avian influenza viruses in water birds, Africa. *Emerg Infect Dis* 2007a; 13:626–629.
- Gaidet N, Dodman T, Caron A, et al. Influenza surveillance in wild birds in Eastern Europe, the Middle East, and Africa: Preliminary results from an ongoing FAO-led survey. *J Wildlife Dis* 2007b; 43:S22–S28.
- Gichuki CM, Gichuki NN. Wetland birds of Kenya. In: Crafter SA, Njuguna SG, Howard GW, eds. *Wetlands of Kenya*. Nairobi, Kenya: National Museums of Kenya, 1992:37–46.
- Guèye EHF. Village egg and fowl meat production in Africa. *World's Poultry Sci J* 1998; 54:82–86.
- Horimoto T, Rivera E, Pearson J, et al. Origin and molecular changes associated with emergence of a highly pathogenic H5N2 influenza virus in Mexico. *Virology* 1995; 213:223–230.
- Ito T, Okazaki K, Kawaoka Y, et al. Perpetuation of influenza A viruses in Alaskan waterfowl reservoirs. *Arch Virol* 1995; 140:1163–1172.
- Joannis T, Lombin LH, De Benedictis P, et al. Confirmation of H5N1 avian influenza in Africa. *Vet Rec* 2006; 158:309–310.
- Lewis A, Pomeroy D. *A Bird Atlas of Kenya*. Brookfield, Vermont: AA Balkema Publishing, 1989.
- Li KS, Guan Y, Wang J, et al. Genesis of a highly pathogenic and potentially pandemic H5N1 influenza virus in eastern Asia. *Nature* 2004; 430:209–213.
- McLean RG, Hall JS, Franklin AB, et al. Avian influenza in wild birds: Environmental sampling strategy for the rapid detection of avian influenza viruses. In: D. L. Nolte, W. M. Arjo, and D. H. Stalman, eds. *Proceedings of the 12th Wildlife Damage Management Conference* 2007:87–93.
- Nasirwa O. Field sampling mission for avian influenza surveillance—Kenya report. National Museums of Kenya, Department of Ornithology, 2006.
- Olsen B, Munster VJ, Wallensten A, et al. Global patterns of influenza A virus in wild birds. *Science* 2006; 312:384–388.
- Owino AO, Oyugi JO, Nasirwa OO, et al. Patterns of variation in waterbird numbers on four Rift Valley lakes in Kenya, 1991–1999. *Hydrobiologia* 2001; 458:45–53.
- Owino AO, Bennun LA, Nasirwa O, et al. Trends in waterbird numbers in the southern Rift Valley of Kenya, 1991–2000. *Waterbirds* 2002; 25:191–201.
- Pearson, DJ, Meadows, BS. Numbers, distribution and seasonality of Palaeartic duck in Kenya. *Scopus* 1992; 15:109–119.
- Perennou C, Mundkur T, Scott DA, et al. The Asian waterfowl census 1987–1991: Distribution and status of Asian waterfowl. UK: International Waterfowl and Wetlands Research Bureau, 1994.
- Root JJ, Bentler KT, Nemeth NM, et al. Experimental infection of raccoons (*Procyon lotor*) with West Nile virus. *Am J Trop Med Hygiene* 2010; 83:803–807.
- Scott DA, Rose PM. *Atlas of Anatidae Populations in Africa and Western Eurasia*. Wetland International, Wageningen, The Netherlands, 1996, 41.
- Sonaiya EB. Family poultry, food security and the impact of HPAI. *World's Poultry Sci J* 2007; 63:132–138.

- Spackman E, Senne DA, Myers TJ, et al. Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. *J Clin Microbiol* 2002; 40:3256–3260.
- Stallknecht DE, Kearney MT, Shane SM, et al. Effects of pH, temperature, and salinity on persistence of avian influenza viruses in water. *Avian Dis* 1990a; 34:412–418.
- Stallknecht DE, Shane SM, Kearney MT, et al. Persistence of avian influenza viruses in water. *Avian Dis* 1990b; 34: 406–411.
- Suss J, Schafer J, Sinnecker H, et al. Influenza virus subtypes in aquatic birds of eastern Germany. *Arch Virol* 1994; 135: 101–114.
- Swayne DE, Halvorson DA. Influenza. In: Saif YM, Barned HJ, Glisson JR, Fadly AM, McDougald LR, Swayne DE, eds. *Diseases in Poultry*. Ames, Iowa: Iowa State University Press, 2003:135–160.
- Szretter KJ, Balish AL, Katz JM. Influenza: Propagation, quantification, and storage. *Curr Protoc Microbiol* 2006; Chapter 15.
- WAHID. Detailed country(ies) disease incidence: Highly path avian influenza. World Animal Health Information Database, 2005–2009a.
- WAHID. Disease outbreak maps: Highly path avian influenza. World Animal Health Information Database, 2005–2009b.
- Wang G, Zhan D, Li L, et al. H5N1 avian influenza re-emergence of Lake Qinghai: Phylogenetic and antigenic analyses of the newly isolated viruses and roles of migratory birds in virus circulation. *J Gen Virol* 2008; 89:697–702.
- Webster RG, Bean WJ, Gorman OT, et al. Evolution and ecology of influenza A viruses. *Microbiol Rev* 1992; 56:152–179.
- Wetlands International. Ramsar sites information service. Wetlands International, 2007.
- WHO. Avian influenza and the pandemic threat in Africa: Risk assessment for Africa. Epidemic and pandemic alert response. World Health Organization, 2005.
- WHO. World Health Organization cumulative number of confirmed human cases of avian influenza A/(H5N1) reported to WHO. World Health Organization, 2011.
- Zar JH. *Biostatistical Analysis*. Englewood Cliffs, New Jersey: Prentice Hall, 1984.

Address correspondence to:

Dr. David Schnabel

Division of Preventive Medicine

Walter Reed Army Institute of Research

503 Robert Grant Rd

Silver Spring, MD 20910

E-mail: david.c.schnabel@us.army.mil