


1973

G73-57 Hot Weather Livestock Stress

Allen C. Wellman

University of Nebraska - Lincoln

Follow this and additional works at: <http://digitalcommons.unl.edu/extensionhist>

 Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

Wellman, Allen C., "G73-57 Hot Weather Livestock Stress" (1973). *Historical Materials from University of Nebraska-Lincoln Extension*. 1658.

<http://digitalcommons.unl.edu/extensionhist/1658>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



NebGuide

PUBLISHED BY COOPERATIVE EXTENSION SERVICE
INSTITUTE OF AGRICULTURE AND NATURAL RESOURCES
UNIVERSITY OF NEBRASKA - LINCOLN



G 73-57

Hot Weather Livestock Stress

Allen C. Wellman
Extension Economist, Marketing

During periods of high temperatures and humidity, livestock losses can occur from "Hot Weather Stress."

Hot weather stress is particularly hazardous to closely confined livestock (those in feedlots, sorting and holding pens, trucks and rail cars).

High relative humidity when the temperature is at 80 degrees or higher adds to the likelihood of profit-stealing losses if necessary precautions are not taken.

Nebraska livestock producers can make their livestock handling and marketing plans flexible

Table 1. Livestock Weather Hazard Guide.

Dry Blub temp.	Relative Humidity Intervals (%)																			
	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
75									70	70	71	71	72	72	73	73	74	74	75	75
76							70	70	70	71	72	72	72	73	74	74	74	75	76	76
77						70	70	71	71	72	72	73	73	74	74	75	75	76	76	77
78				70	70	71	71	72	72	73	73	74	74	75	75	76	76	77	78	78
79			70	70	71	72	72	73	73	74	74	75	75	76	77	77	78	78	79	80
80		70	70	71	72	72	73	73	74	74	75	75	76	77	77	78	78	79	80	81
81		70	71	71	72	73	73	74	74	75	75	76	77	77	78	78	79	80	80	81
82		70	71	71	72	73	73	74	74	75	75	76	77	77	78	79	79	80	81	82
83	70	71	71	72	73	73	74	74	75	75	76	77	78	78	79	DANGER	81	82	82	83
84	70	71	72	72	73	74	74	75	75	76	77	78	78	79	80	80	81	82	83	84
85	71	72	72	73	74	74	75	75	76	77	78	78	79	80	81	81	82	83	84	85
86	71	72	73	74	74	75	76	77	78	78	79	80	81	81	82	83	84	84	85	86
87	72	73	73	74	75	76	77	77	78	79	80	81	81	82	83	84	85	85	86	87
88	72	73	74	75	76	76	77	78	79	80	81	81	82	83	84	EMERGENCY	87	87	88	89
89	73	74	74	75	76	77	78	79	80	80	81	82	83	84	85	86	86	87	88	89
90	73	74	75	76	77	78	79	79	80	81	82	83	84	85	86	87	87	88	89	90
91	74	75	76	76	77	78	79	80	81	82	83	84	85	86	86	87	88	89	90	91
92	74	75	76	77	78	79	80	81	82	83	84	84	85	86	87	88	89	90		
93	75	76	77	78	79	80	80	81	82	83	84	85	87	87	88	89	90			
94	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90				
95	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90					
96	76	77	78	79	80	81	82	84	84	86	87	88	89	90	91					
97	77	78	79	80	81	82	83	84	85	86	87	88	90	91						
98	77	78	79	80	82	83	84	85	86	87	88	89	90							
99	78	79	80	81	82	83	84	86	87	88	88	90								
100	78	79	80	82	83	84	85	86	87	88	90	91								
105	80	82	83	84	86	87	89	90	91											

Temperatures above 100 are always "Danger" and if the relative humidity is above 25 per cent, the situation is "Emergency".
Source: Livestock Conservation, Inc.

enough or take necessary precautions to reduce or eliminate livestock hot weather stress by following the Livestock Weather Hazard Guide (*Table 1*).

Alert livestockmen will adjust ventilation and bedding to the prevailing temperatures. If livestock must be transported, vehicles should be bedded with sand, sawdust, shavings or combination of these in the summer. Avoid the use of straw, particularly oats straw, in vehicles with solid sides or tight boxes during hot weather. Use "wet" bedding in "shirt sleeve" weather. Sprinkling animals in confined areas may be advisable when the temperature is above 80 degrees outside.

How To Use the Livestock Weather Hazard Guide

Obtain the current or expected temperature and relative humidity by listening to a weather forecast or read your own thermometer and hygrometer. Bear in mind what the temperature and humidity is likely to be by the time the animals reach their destination.

Locate the temperature (actual or forecast) in the column at the LEFT of *Table 1.*)

Follow the line to the RIGHT until you come to the "relative humidity" which is equal to or less than that reported or forecast. *This number tells you what weather stress category (Alert, Danger or Emergency) you are in and how safe your confined livestock may be.*

This stress is closely related to the "discomfort index" for humans as developed by the Weather Bureau, originally termed the "temperature-humidity" index (THI).

Stress Categories

ALERT (THI of 75-78)

A forecast of temperature and humidity conditions in this range at time of handling, loading or before animals reach their destination calls for an "alert."

Additional precautions may be needed to avoid excessive losses or prepare for higher THI. DANGER (THI of 79-83)

Temperature and humidity readings in this range are not only dangerous to confined livestock but there is a need to adopt additional measures to avoid severe losses.

EMERGENCY (THI of 84 and higher)

A severe situation has developed, consider changing livestock handling and shipping plans, or if plans cannot be changed, these four suggestions at a minimum should be followed:

1. All handling stress should be kept at a minimum.
2. Keep animals in position for free circulation of air.
3. Provide shade if at all possible.
4. Make water readily available for drinking. If water is to be used to cool the animals, avoid "shock" from cold water in too huge quantities.

A continuous sprinkling or coarse mist will lower the temperature to a safe level with a minimum of danger to the animals.

Loading rested hogs onto wet bedding will minimize the heat stress problem during transit.

The best solution is to plan your livestock handling and shipping activities for the periods when the THI reading is below 75. Moving livestock when the THI is above 75 should be considered risky at best.

File Under: ANIMALS, GENERAL
C-1, Management