

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

2015

Exposure to Rabies in Small Indian Mongooses (*Herpestes auropunctatus*) from Two Regions in Puerto Rico

Are R. Berentsen

USDA National Wildlife Research Center, Are.R.Berentsen@aphis.usda.gov

Shylo R. Johnson

USDA/APHIS/WS National Wildlife Research Center

Amy T. Gilbert

USDA/APHIS/WS National Wildlife Research Center, Amy.T.Gilbert@aphis.usda.gov

Kurt C. VerCauteren

USDA-APHIS-Wildlife Services, kurt.c.vercauteren@aphis.usda.gov

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Life Sciences Commons](#)

Berentsen, Are R.; Johnson, Shylo R.; Gilbert, Amy T.; and VerCauteren, Kurt C., "Exposure to Rabies in Small Indian Mongooses (*Herpestes auropunctatus*) from Two Regions in Puerto Rico" (2015). *USDA National Wildlife Research Center - Staff Publications*. 1734. https://digitalcommons.unl.edu/icwdm_usdanwrc/1734

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Exposure to Rabies in Small Indian Mongooses (*Herpestes auropunctatus*) from Two Regions in Puerto Rico

Are R. Berentsen,^{1,2} Shylo R. Johnson,¹ Amy T. Gilbert,¹ and Kurt C. VerCauteren¹ ¹US Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services, National Wildlife Research Center, 4101 LaPorte Ave., Fort Collins, Colorado 80521, USA; ²Corresponding author (email: Are.R.Berentsen@aphis.usda.gov)

ABSTRACT: The small Indian mongoose (*Herpestes auropunctatus*) was introduced to several Caribbean Islands to control rat (*Rattus* spp.) damage to sugarcane plantations. Mongooses failed at suppressing rat populations and are now considered pests throughout most of their introduced range. Importantly, mongooses are rabies reservoirs on several Caribbean Islands. In Puerto Rico, mongooses have been implicated in up to 70% of reported animal rabies cases. There is no rabies vaccination program for wildlife in Puerto Rico, and data on rabies in mongooses are limited. We conducted a serosurvey of mongooses in two different ecologic environments in Puerto Rico: El Yunque National Forest and Cabo Rojo National Wildlife Refuge. We collected 119 serum samples from 112 mongooses, 44 (39.3%) of which were positive for rabies virus–neutralizing antibodies. We also collected oral swabs from 147 mongooses, including 88 from which we also collected serum. No oral swabs were positive for rabies virus RNA. Our data support previous research suggesting rabies virus is circulating within the mongoose population on Puerto Rico.

Key words: Caribbean, *Herpestes auropunctatus*, mongoose, Puerto Rico, rabies, wildlife disease.

The small Indian mongoose (*Herpestes auropunctatus*) was introduced to the Caribbean Islands in the 1870s, primarily to control rat (*Rattus* spp.) damage to sugarcane (Hoagland et al. 1989). Mongooses are now considered pests throughout most of their introduced range and are the primary rabies reservoir on several Caribbean Islands (Everard and Everard 1992). In Puerto Rico, mongooses account for up to 75% of reported rabies cases (Krebs et al. 1998; Dyer et al. 2014). In Puerto Rico, the first laboratory-confirmed rabies-infected mongooses were reported

in 1950, although references to rabies date back prior to mongoose introduction (Tierkel et al. 1952). Between 1986 and 1990, 71.5% (236/330) of suspect mongooses tested in Puerto Rico were positive for rabies virus, as were 2.6% (4/152) of randomly trapped mongooses (Everard and Everard 1992). In comparison, Everard et al. (1981) reported rabies infection in 1.3% of >11,000 mongooses in Grenada between 1968 and 1977. Velez (1998) found 19.3% of mongooses sampled in portions of El Yunque National Forest (YNF), Puerto Rico, positive for rabies virus–neutralizing antibodies (RVNA) when tested by enzyme-linked immunosorbent assay (ELISA), although none were rabies virus positive. Few data are available on rabies exposure in mongooses on Puerto Rico, as most studies have occurred on other Caribbean islands, primarily Grenada (Everard et al. 1981; Zieger et al. 2014). We evaluate sera for RVNA and saliva samples for rabies virus RNA to investigate mongoose exposure to rabies in two ecologically different environments on Puerto Rico.

We conducted our study at YNF and Cabo Rojo (CR) National Wildlife Refuge (Fig. 1). The YNF is a subtropical rainforest approximately 40 km southeast of San Juan. In YNF, samples were collected from a 1.0-km² region of Palo Colorado and Sierra Palma forest types in October 2011 and March 2012. The CR is >7 km² of subtropical dry forest in southwestern Puerto Rico. We collected samples from CR in two ~1-km² regions, one each on the eastern and western sides of the




FIGURE 1. El Yunque National Forest and Cabo Rojo National Wildlife Refuge, Puerto Rico.

refuge, during September 2011, April–May 2012, and August 2014. Both sample sites are dominated by scrub-class vegetation (Weaver and Schwagerl 2008).

We used cage traps (Tomahawk Trap Company, Hazelhurst, Wisconsin, USA) baited with canned tuna to live capture mongooses (Quinn and Whisson 2005). Mongooses were hazed out of the trap into a cone-shaped canvas bag, physically restrained, and immobilized with an intramuscular injection of Telazol® (tiletamine-zolazepam, Fort Dodge Laboratories, Fort Dodge, Iowa, USA) at a dose rate of 5 mg/kg (Kreeger and Armemo 2007). We collected 0.5–1.0 cc of whole blood by venipuncture of the cranial vena cava, as described for ferrets (Briscoe and Syring 2004), and injected a passive integrated transponder tag (Avid Identification Systems, Inc., Norco, California, USA) subcutaneously

between the shoulder blades of each captured mongoose. We used a sterile cotton swab to collect saliva samples and stored swabs in a bovine albumin 1 viral transport medium to reduce potential bacterial or fungal contamination (Shriner et al. 2012). Blood samples were centrifuged, and sera were transferred by pipette into cryovials. Serum samples and oral swabs were frozen at -10 C for up to 14 d and transferred to -80 C until analysis.

Sera were analyzed for RVNA end-point titers by using the rapid fluorescent focus inhibition test (Smith et al. 1996) by the rabies laboratory at Kansas State University (Manhattan, Kansas, USA). Mongooses with sera $>0.1\text{ IU/mL}$ were considered positive for RVNA. We compared prevalences by using the Logistic procedure in SAS (SAS Institute Inc., Cary, North

Carolina, USA). We accepted statistical significance at $\alpha=0.05$.

Oral swabs were processed by Atlanta Health Associates (Cumming, Georgia, USA). Total RNA was extracted by using TRIzol reagent (Invitrogen™, Ambion®, Carlsbad, California, USA). Heminested reverse-transcription PCR was performed for amplification of a partial sequence of the rabies virus nucleoprotein gene by using primers 1066:304 for the primary and 1087:304 for the heminested reactions. Positive (RNA from mouse brain infected with the mongoose-variant rabies virus) and negative (water) control samples were used with each PCR run. Reaction products were visualized by gel electrophoresis (Freuling et al. 2015).

We collected 41 sera from 39 mongooses at YNF. Fourteen sera were RVNA positive from 13 mongooses (nine males and four females), including one recaptured female whose titer decreased from 2.8 to 1.2 IU/mL between sampling periods (138 d). The geometric mean titer of positive mongooses at YNF was 0.52 IU/mL (range: 0.1–2.8 IU/mL), and antibody prevalence was 33% (13 unique positives and 39 unique captures; 95% confidence interval [CI]: 19–48%). At CR, 78 sera were collected from 73 mongooses. Thirty-two sera were RVNA positive from 31 individuals (20 males and 11 females). Five animals were recaptured during subsequent seasons: two females and one male, initially captured in 2011, were recaptured in 2012. Among recaptures, one male and one female seroconverted between captures (from <0.1 IU/mL to 0.4 and 1.4 IU/mL for the male and female, respectively), while the second female remained antibody negative. One male captured in 2012 (recaptured in 2014) seroconverted between captures (<0.1 IU/mL in 2012 and 10.7 IU/mL in 2014). A positive female captured in 2011 was recaptured in 2014, and the RVNA titer had decreased from 0.6 IU/mL to 0.5 IU/mL (1,073 d). Geometric mean titer of antibody-positive mongooses at CR was

1.62 IU/mL (range: 0.1–50.0 IU/mL), and antibody-prevalence was 42% (31 positive and 73 unique captures; 95% CI: 31–54%). Antibody prevalence (including recaptured animals) did not differ between sites ($\chi^2=0.53$; $P=0.46$) or seasons (wet vs. dry; $\chi^2=0.02$; $P=0.88$), or among years ($\chi^2=1.28$; $P=0.53$).

Oral swabs from 146 unique mongooses (72 from CR and 74 from YNF) were tested for rabies virus RNA. This included swabs from 88 individual mongooses whose sera were screened, and 33 of which were RVNA positive. All swabs were negative for viral RNA.

The RVNA detected in mongoose sera suggests prior peripheral virus exposure and adaptive immune response induction rather than an active nervous system infection at the time of sampling. A peripheral neutralizing antibody response may not be routinely induced or if it is induced, tends to be observed during late stages of clinical infection (Lafon 2013), which were not observed among mongooses in this study. Furthermore, viral shedding in saliva by infected rabid animals is intermittent and, though possible to detect several days in advance of clinical signs, is typically detected during clinical infection (Hanlon et al. 2007). Zieger et al. (2014) found that of 171 mongooses sampled on Grenada, only two (1.7%) were positive for rabies virus, whereas 33 (19.3%) had RVNA titers at the 0.1 IU/mL threshold. Rabies virus RNA was not found in saliva from either infected mongoose in the Zieger et al. (2014) study, and neither mongoose showed clinical signs of infection.

Overall, 39% of mongooses sampled in this study were RVNA positive. This is slightly higher than previous estimates from Puerto Rico (19.3%; Velez 1998) and Grenada (30.0%, Everard et al. 1981; 19.3%, Zieger et al. 2014), although caution must be exercised during comparison as Velez (1998) used an ELISA assay for antibody detection and Everard et al. (1981) used a lower cutoff than our study

to report positive RVNA results. The number of rabies-positive mongooses in our study is unknown because mongooses were not lethally sampled, but prevalence estimates from similar environments of Grenada are 1.7% (Zieger et al. 2014) and 1.3% (Everard et al. 1981). Rabies transmission routes among mongooses are poorly understood, but bite contacts during intraspecific aggression are likely the primary route.

Phylogenetic studies suggest the rabies virus circulating in mongooses of Puerto Rico is related to the North Central skunk variant and cosmopolitan dog lineages of rabies virus (Nadin-Davis et al. 2008). Virus strains from Puerto Rico (Nadin-Davis et al. 2008), Grenada (Zieger et al. 2014), and Cuba (Nadin-Davis et al. 2006) do not cluster in a monophyletic group, suggesting independent introductions to different Caribbean islands.

Rabies is maintained by mongooses as a reservoir species rather than a spillover host. Although domestic dogs (*Canis familiaris*) can account for up to 26% of rabies cases in Puerto Rico, infection is typically attributed to the mongoose variant (Dyer et al. 2014). It is not possible to distinguish whether multiple exposures occurred among recaptured animals in this study, but our findings support reports by Everard et al (1981), indicating potential for long-term maintenance of RVNA in mongoose. The seroconversion events detected among recaptured animals provides strong evidence of active rabies virus circulation among mongooses in Puerto Rico. Future research should include virus isolation from saliva of actively infected mongooses. With no wildlife vaccination program in place, mongoose rabies in Puerto Rico continues to pose a threat to domestic animal and human health.

We thank P. Quiñones, N. Moers, C. Ellis, E. Blizzard, and S. Eaton for field assistance. We also thank F. Cano, O. Diaz, and R. Lopez-Ortiz for logistic assistance. Capture and handling of mon-

gooses was approved by the US Department of Agriculture National Wildlife Research Center's animal care and use committee under protocol QA-1856.

LITERATURE CITED

- Briscoe JA, Syring R. 2004. Techniques for emergency airway and vascular access in special species. *Semin Avian Exot Pet Med* 13:118–131.
- Dyer JL, Yager P, Orciani L, Greenberg L, Wallace R, Hanlon CA, Blanton JD. 2014. Rabies surveillance in the United States during 2013. *J Am Vet Med Assoc* 245:1111–1123.
- Everard COR, Baer GM, Alls ME, Moore SA. 1981. Rabies serum neutralizing antibody in mongooses from Grenada. *Trans R Soc Trop Med Hyg* 5:654–666.
- Everard COR, Everard JD. 1992. Mongoose rabies in the Caribbean. *Ann N Y Acad Sci* 653:356–366.
- Freuling CM, McElhinney LM, Fooks AR, Müller TF. 2015. Gel-based reverse transcription-polymerase chain reaction. In: *Current laboratory techniques in rabies diagnosis, research and prevention*, Vol. 2, Rupprecht CE, Nagarajan T, editors. Elsevier, San Diego, California, pp. 119–128.
- Hanlon CA, Niezgodka M, Rupprecht CE. 2007. Rabies in terrestrial animals. In: *Rabies*, 2nd Ed., Jackson AC, Wunner WH, editors. Academic Press, London, UK, pp. 201–258.
- Hoagland DB, Horst GR, Kilpatrick CW. 1989. Biogeography and population biology of the mongoose in the West Indies. In: *Biogeography of the West Indies, past, present and future*, Woods CA, editor. Sandhill Crane Press, Gainesville, Florida, pp. 611–634.
- Krebs JW, Smith JS, Rupprecht CE, Childs JE. 1998. Rabies surveillance in the United States during 1997. *J Am Vet Med Assoc* 213:1713–1728.
- Kreeger TJ, Arnemo JM. 2007. *Handbook of wildlife chemical immobilization*, 3rd Ed. Sunquest Printing, Vancouver, British Columbia, Canada, 432 pp.
- Lafon M. 2013. Immunology. In: *Rabies*, 3rd Ed., Jackson A, editor. Academic Press, San Diego, California, pp. 387–408.
- Nadin-Davis SA, Torres G, de Los Angeles Ribas M, Guzman M, Cruz de La Paz R, Morales M, Wandeler AI. 2006. A molecular epidemiological study of rabies in Cuba. *Epidemiol Infect* 134:1313–1324.
- Nadin-Davis SA, Velez J, Malaga C, Wandeler AI. 2008. A molecular epidemiological study of rabies in Puerto Rico. *Virus Res* 131:8–15.
- Quinn JH, Whisson DA. 2005. The effects of anthropogenic food on the spatial behavior of small Indian mongooses (*Herpestes javanicus*) in a subtropical rainforest. *J Zool* 267:339–350.

- Shriner SA, VanDalen KC, Mooers N, Ellis JW, Sullivan HJ, Root JJ, Pelzel AM, Franklin AB. 2012. Low-pathogenic avian influenza virus in wild house mice. *PLoS One* 7:e39206.
- Smith JS, Yager PA, Baer GM. 1996. A rapid fluorescent focus inhibition test (RFFIT) for determining rabies virus-neutralizing antibody. In: *Laboratory techniques in rabies*, 4th Ed., Meslin FX, Kaplan MM, Koprowski H, editors. World Health Organization, Geneva, Switzerland, pp. 181–190.
- Tierkel ES, Arbona G, Rivera A, De Juan A. 1952. Mongoose rabies in Puerto Rico. *Public Health Rep* 67:274.
- Velez JV. 1998. *Presencia de anticuerpos antirábicos y diagnóstico de la rabia en mangostas (Herpestes auropunctatus) capturadas en El Bosque Nacional del Caribe (El Yunque)*. MS Thesis, University of Puerto Rico, San Juan, Puerto Rico, 125 pp.
- Weaver PL, Schwagerl JJ. 2008. Secondary forest succession and tree planting at the Laguna Cartagena and Cabo Rojo wildlife refuges in southwestern Puerto Rico. *Ambio* 37:598–603.
- Zieger U, Marston DA, Sharma R, Chikweto A, Tiwari K, Sayyid M, Louison B, Goharriz H, Voller K, Breed AC, et al. 2014. The phylogeography of rabies in Grenada, West Indies, and implications for control. *PLoS Negl Trop Dis* 8:e3251.

Submitted for publication 22 January 2015.

Accepted 25 March 2015.