

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Historical Materials from University of  
Nebraska-Lincoln Extension

Extension

---

October 2013

## EC113 Seed Treatment for Stinking Smut (Bunt) in Wheat

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>

---

"EC113 Seed Treatment for Stinking Smut (Bunt) in Wheat" (2013). *Historical Materials from University of Nebraska-Lincoln Extension*. 2098.

<https://digitalcommons.unl.edu/extensionhist/2098>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

EC 1136  
**RECEIVED**

DEC 2 1910  
COLLEGE OF AGRICULTURE  
LIBRARY

NEBRASKA

AGRICULTURAL EXTENSION WORK

Circular

Neb. Agr. College & U. S. Dept. of Agr. Cooperating

1136

W. H. Brokaw, Director, Lincoln

SEED TREATMENT FOR STINKING SMUT (Bunt) IN WHEAT

Soaking method: Run the seed thru a good fanning mill before treatment. This will remove many of the shriveled and light weight kernels, spore balls, Many of the free smut spores, and dirt. Make up a solution consisting of 1 pint Formalin to 35 gallons of water. Put the seed in loosely filled burlap sacks, dip into the solution and soak for five minutes. Drain on slats over the barrels or tanks to prevent wasting the solution. Place the seed on a clean floor or canvas, and cover with sacks or canvas for two hours. Dry by spreading in a thin layer and stirring occasionally with a rake or shovel. If the seed is sown when swollen, open the drill to offset this. Disinfect the drill with a one to 10 solution of Formalin. It is recommended that Formalin in sealed bottles be purchased.

(over)



R02378 93842

NEBRASKA

AGRICULTURAL EXTENSION WORK

Circular

Nebr. Agr. College &amp; U. S. Dept. of Agr. Cooperating

113

W. H. Brokaw, Director, Lincoln

## SEED TREATMENT FOR STINKING SMUT (Bunt) IN WHEAT

Sprinkling method: This method is not as effective as the soaking method and is not to be recommended when the soaking method can be used. Where the wheat is only slightly affected and it is run thru a fanning mill with a good air blast to remove the spore balls this method may be successfully used.

Sprinkle on the seed a solution of one pint of formalin to 35 gallons of water, using about one gallon to the bushel, shoveling over at the same time to make sure of uniform wetting. Cover in a pile with bags or canvas and handle as in the soaking method. The effectiveness of this treatment depends on the thoro wetting of each seed. If the seed is swollen, open the drill to offset this. Disinfect the drill with a one to 10 solution of Formalin. It is recommended that Formalin in sealed bottles be purchased.

(over)

E.C. # 113

copy 1