

2018

Cottontail rabbits shed clade 2.3.4.4 H5 highly pathogenic avian influenza A viruses

J. Jeffrey Root

National Wildlife Research Center, jeff.root@aphis.usda.gov

Angela M. Bosco-Lauth

USDA National Wildlife Research Center

Nicole L. Marlenee

USDA National Wildlife Research Center

Richard A. Bowen

USDA National Wildlife Research Center

Follow this and additional works at: http://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Life Sciences Commons](#)

Root, J. Jeffrey; Bosco-Lauth, Angela M.; Marlenee, Nicole L.; and Bowen, Richard A., "Cottontail rabbits shed clade 2.3.4.4 H5 highly pathogenic avian influenza A viruses" (2018). *USDA National Wildlife Research Center - Staff Publications*. 2142.
http://digitalcommons.unl.edu/icwdm_usdanwrc/2142

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Cottontail rabbits shed clade 2.3.4.4 H5 highly pathogenic avian influenza A viruses

J. Jeffrey Root¹ · Angela M. Bosco-Lauth² · Nicole L. Marlenee² · Richard A. Bowen²

Received: 8 March 2018 / Accepted: 31 May 2018

© This is a U.S. government work and its text is not subject to copyright protection in the United States; however, its text may be subject to foreign copyright protection 2018

Abstract

During 2014–2015, clade 2.3.4.4 H5Nx highly pathogenic (HP) avian influenza A viruses (IAV) were first detected in North America and subsequently caused one of the largest agricultural emergencies in U.S. history. Recent evidence has suggested that cottontail rabbits can shed multiple IAV subtypes. We experimentally infected cottontail rabbits with three HP H5Nx IAVs. All rabbits tested shed virus on at least one day by at least one route. Cottontail rabbits appear to be an exception to the limited capacity for replication that has been previously reported for certain other mammalian species inoculated with clade 2.3.4.4 HP H5Nx avian influenza A viruses.

During late 2014, highly pathogenic (HP) H5N8 avian influenza A virus (IAV) and reassortant H5N2 viruses were identified in Washington, USA from wild and/or captive birds fed wild waterfowl [6]. The H5N2 virus spread rapidly in the U.S. during the winter and spring of 2015, reaching Iowa by mid-April [17]. The effects of these viruses on the U.S. poultry industry were tremendous, and while multiple states were affected, in Minnesota and Iowa alone, approximately 40 million birds either died from HP IAV infections or were culled [17]. In Iowa, the economic impact of this outbreak to the state has been estimated to be greater than 1 billion U.S. dollars [3].

The potential roles of peridomestic wildlife in the epidemiology of avian IAVs have received increasing attention over recent years [5, 9, 16]. The 2015 HP avian IAV outbreaks in the U.S., as well as other emergent, high consequence strains (both low pathogenic and HP that have high zoonotic or agricultural health implications) elsewhere (e.g., Asian H7N9 and H5N1), have motivated researchers

to pursue several studies associated with questions involving the replication competence of some of these synanthropic animals [2, 4, 14]. However, to date, few studies have addressed this issue in clade 2.3.4.4 HP H5 viruses.

During recent years, some lagomorph species (e.g., rabbits, hares, and pika) have been shown to be either experimentally susceptible (e.g., cottontail rabbits) or naturally exposed (e.g., plateau pika [*Ochotona curzoniae*]) to various IAVs [12, 19, 21]. Further, a recent study indicated that cottontail rabbits have the capacity to transmit an IAV back to birds through shared resources [13]. Motivated by these IAV susceptibilities in wild lagomorphs, the objective of this study was to characterize the infection dynamics in a common mammalian farm-side resident, the cottontail rabbit (*Sylvilagus* sp.), experimentally inoculated with three HP H5Nx clade 2.3.4.4 IAVs that have been recently detected in North America.

Three viruses were used in the experimental infections. These included A/Northern pintail/Washington/40964/2014 (H5N2) (hereinafter = Northern pintail), A/turkey/Minnesota/9845-4/2015 (H5N2) (hereinafter = turkey), and A/gyrfalcon/Washington/41088-6/2014 (H5N8) (hereinafter = gyrfalcon). All viruses were kindly provided by the National Veterinary Services Laboratory, Ames, IA. Viruses were propagated in embryonated chicken eggs and subsequently titrated with plaque assays (see methods below).

Twelve cottontail rabbits (*Sylvilagus* sp.) were wild-caught in Larimer County, CO with box traps. The twelve cottontail rabbits were dusted for ectoparasites and bled to

Handling Editor: Ayato Takada.

J. Jeffrey Root and Angela M. Bosco-Lauth contributed equally to this work.

✉ J. Jeffrey Root
Jeff.root@aphis.usda.gov

¹ United States Department of Agriculture, National Wildlife Research Center, Fort Collins, CO, USA

² Colorado State University, Fort Collins, CO, USA

assess if they had pre-existing IAV antibodies (see below). In addition, a subcutaneous transponder (IPPT-300 transponders, Bio Medic Data Systems) that provided an individual identification number and temperature readings was implanted into each cottontail. Following brief quarantine periods, animals were transferred to an enhanced animal biosafety level-three (ABSL-3) facility at Colorado State University. All cottontails were housed in rabbit cages placed within HEPA-filtered isolator cages outfitted with a nest box. Cottontail rabbits were maintained with rabbit chow, alfalfa, and fruit. Food and water were replenished daily. Animal care and use protocols were approved by the National Wildlife Research Center and Colorado State University.

The twelve cottontail rabbits, all assessed to be negative to IAV antibodies, were divided into three experimental groups of four rabbits for each of the three viruses listed above and were inoculated with approximately $10^{5.3}$ plaque forming units (PFU) (turkey virus) or $10^{5.9}$ PFU (Northern pintail and gyrfalcon viruses) in 200 μ l BA-1 [16] diluent vehicles by the nasal route. Each nostril received approximately 100 μ l. Following inoculations, one-half of the cottontail rabbits were sampled on odd days ($n = 2$ for each virus) and the other one-half were sampled on even days ($n = 2$ for each virus) over 1–8 days post-infection (DPI). This sampling time period was chosen based on previous studies indicating that cottontail rabbits typically cease shedding or shed other IAVs at low levels by the end of this time-frame [12]. Daily sampling consisted of an oral swab (stored in 1 mL of BA-1 viral transport media) and a nasal flush (1 mL of

BA-1) for sampled animals, as well as general health observations of all animals. In addition, body temperature data was recorded each day an animal was processed, when possible. All swab and nasal flush samples were stored at -80° C prior to testing. Animals were euthanized on 18 DPI. To accomplish nasal inoculations and animal processing, cottontails were anesthetized with 10mg/kg ketamine and 1 mg/kg xylazine in combination intramuscularly.

Oral swab and nasal flush samples were tested by plaque assay as described in a previous study [1]. The limit of detection for both sample types was 10 PFU/mL. Pre-experiment sero-negativity was determined with agar gel immunodiffusion (AGID) tests [10], a procedure that has been successfully employed during previous studies with cottontail rabbit sera [12, 15]. Post-experiment serology was conducted with standard hemagglutination-inhibition (HI) assays using equine erythrocytes and an H5 virus with cottontail rabbit sera collected on 18 DPI. Serological responses were delineated as HI titers of ≥ 10 .

Cottontail rabbits were infected by the three tested viruses, as all cottontail rabbits in each virus group ($n = 4$) shed virus on at least one DPI (Tables 1 and 2) but did not exhibit any overt signs of disease. The highest virus titers were noted from nasal samples from rabbits infected with the H5N2 turkey virus, as two rabbits shed at levels of $\geq 10^{5.1}$ PFU/mL (Table 1). Oral shedding of this virus was also relatively high, as one rabbit shed a titer of $10^{4.3}$ PFU/mL, while three of four rabbits shed at levels of $> 10^{3.0}$ PFU/mL (Table 2). Of interest, the rabbit that shed the highest titer in this treatment group was the only animal that shed

Table 1 Nasal shedding and serological responses of cottontail rabbits (*Sylvilagus* sp.) experimentally infected with clade 2.3.4.4 highly pathogenic H5N2 and H5N8 avian influenza A viruses

Rabbit number	Virus	Days post infection								HI status ^f
		1	2	3	4	5	6	7	8	
1	Turkey ^a	5.1 ^d		4.5		1.6		<1		+
2	Turkey		5.3		2.4		4.6		2.2	+
3	Turkey	3.9		4.2		ND ^e		ND		ND
4	Turkey		3.4		1.3		<1		<1	+
5	Northern Pintail ^b	3.8		2.3		<1		<1		+
6	Northern Pintail		2.1		<1		ND ^e		ND	ND
7	Northern Pintail	3.9		2.8		<1		<1		+
8	Northern Pintail		3.6		<1		<1		<1	+
9	Gyrfalcon ^c	3.5		3.0		2.3		<1		+
10	Gyrfalcon		2.5		2.2		<1		<1	+
11	Gyrfalcon	2.5		1.0		<1		<1		+
12	Gyrfalcon		2.9		1.3		<1		<1	+

^aA/turkey/Minnesota/9845-4/2015 (H5N2)

^bA/Northern pintail/Washington/40964/2014 (H5N2)

^cA/gyrfalcon/Washington/41088-6/2014 (H5N8)

^dValues represent \log_{10} PFU/mL

^eND = not done. Animal died from complications with anesthesia during a previous sampling period

^fHI conducted with sera collected on 18 DPI. A “+” indicates a positive serological response (see methods)

Table 2 Oral shedding of cottontail rabbits (*Sylvilagus* sp.) experimentally infected with clade 2.3.4.4 highly pathogenic H5N2 and H5N8 avian influenza A viruses

Rabbit number	Virus	Days post infection							
		1	2	3	4	5	6	7	8
1	Turkey ^a	3.1 ^d		3.3		<1		<1	
2	Turkey		4.3		1.8		3.4		1.8
3	Turkey	2.3		3.4		ND ^e		ND	
4	Turkey		2.1		<1		<1		<1
5	Northern Pintail ^b	3.0		1.3		<1		<1	
6	Northern Pintail		2.1		<1		ND ^e		ND
7	Northern Pintail	2.4		2.3		<1		<1	
8	Northern Pintail		2.5		<1		<1		<1
9	Gyrfalcon ^c	2.0		1.8		2.1		<1	
10	Gyrfalcon		<1		1.0		<1		<1
11	Gyrfalcon	1.7		<1		<1		<1	
12	Gyrfalcon		C ^f		C		C		C

^aA/turkey/Minnesota/9845-4/2015 (H5N2)

^bA/Northern pintail/Washington/40964/2014 (H5N2)

^cA/gyrfalcon/Washington/41088-6/2014 (H5N8)

^dValues represent log₁₀ PFU/mL

^eND = not done. Animal died from complications with anesthesia during a previous sampling period

^fBacterial contamination noted in assays from oral swab samples, likely due to colonization of the oral cavity of this particular rabbit

each day it was sampled (i.e., 2, 4, 6, and 8 DPI) by both the oral and nasal routes. Overall, the turkey virus produced the highest shedding levels for cottontail rabbits on most DPI as measured by daily maximums for both nasal ($10^{5.3}$ PFU/mL) and oral ($10^{4.3}$ PFU/mL) samples (Figure 1). One rabbit shed virus through at least 8 DPI (the last day it was sampled), which is consistent with a previous study [12].

Viral shedding of the H5N2 Northern pintail virus tended to be of a lower titer and of a shorter duration when compared to the turkey virus. All rabbits shed this virus by the nasal route, and three of four rabbits shed at levels of $\geq 10^{3.6}$ PFU/mL (Table 1). In addition, all animals shed virus by the oral route on at least one DPI. The highest titer shed by the oral route was $10^{3.0}$ PFU/mL (Table 2). Two of four cottontails associated with this virus only shed virus at detectable levels the first day they were sampled (2 DPI). The H5N2 Northern pintail virus was shed up to 3 DPI by the oral and nasal routes (Tables 1 and 2).

Overall, the H5N8 gyrfalcon virus produced the lowest levels of viral shedding in cottontail rabbits in terms of peak viral titers. The highest titer from a nasal flush sample was $10^{3.5}$ PFU/mL, while the highest titer from an oral swab sample was $10^{2.1}$ PFU/mL, of the three animals evaluated for oral shedding for this virus (Tables 1 and 2). In one of four individuals, bacterial contamination was noted in assays from oral swab samples, likely due to colonization of the oral cavity of this particular rabbit. Although peak levels of shedding of this virus were lower than the Northern pintail virus, the duration of shedding was longer for the gyrfalcon

virus (up to 5 DPI by one rabbit by both shedding routes; Tables 1 and 2).

Overall, 10 of 10 cottontail rabbits seroconverted for the three viruses. Serology tests indicated 3/3 animals seroconverted following inoculations with the turkey virus, 3/3 animals seroconverted following inoculations with the Northern pintail virus, and 4/4 animals seroconverted following inoculations with the gyrfalcon virus (Table 1; see table footnotes for denominator explanations).

Recent literature suggests that cottontail rabbits are susceptible to multiple LP IAV strains, and can be susceptible to low doses of virus [12, 14, 15]. However, to our knowledge, this is the first evaluation of HP viruses in cottontail rabbits. Consistent with other studies evaluating cottontail rabbits, the current study also indicates that this species can shed relatively high viral titers by multiple routes. Of interest, the virus that produced the highest levels of viral shedding in cottontails during the current study was isolated from poultry, thereby suggesting that this species may pose more of a threat for viral movement between poultry barns once the virus is established at a farm, as compared to the rabbits acquiring a viral infection from a wild bird on the farm premises. Nonetheless, the higher levels of oral and nasal shedding observed in cottontail rabbits inoculated with a poultry virus (Figure 1) were largely the result of high titers of this virus from two of four rabbits. Thus, individual heterogeneity in shedding responses to the three viruses likely accounts for at least some the observed differences in the viral titers that were shed.




Fig. 1 Average nasal (top) and oral (bottom) shedding of three different highly pathogenic avian influenza A viruses in experimentally infected cottontail rabbits. Samples with values of < 10 PFU/mL or those that were not taken (see Table 1) are evaluated as zero

Influenza viral isolates from wild lagomorphs have been reported from the Qinghai Lake region in China, where intensive wildlife influenza investigations have been undertaken. For example, H5N1, H9N2, and H7N2 IAVs have been isolated from plateau pika in Qinghai Province [18, 19, 21]. Further, antibodies reactive with H5 and H9 subtypes have also been reported in this species from this region [20, 21]. This suggests that a comparable AI hotspot in North America, one where wild lagomorphs and waterfowl are in close proximity to each other, could facilitate natural IAV infections in cottontails. At this point in time, no large-scale IAV surveillance studies have been conducted in cottontail rabbits; however, this could represent a useful investigation if an appropriate location, harboring large numbers of cottontail rabbits and waterfowl in an area with known IAV activity, is identified.

Ferrets that were experimentally inoculated with the same Northern pintail and gyrfalcon viruses as used in

this study showed minimal nasal shedding, thereby leading these authors to conclude that this species has limited capacity for replication of these viruses [7]. However, others have suggested efficient replication of these viruses in the respiratory tracts of this species [11]. No viral RNA was detected from nasal swabs from swine infected with the two viruses mentioned above when sampled on 1, 3, or 5 DPI; however, broncho-alveolar lavage fluids did show evidence of viral RNA on 3 and 5 DPI [8]. Consistent with the viral titers in the lungs of mice infected with two of the viruses used in the current study [7], cottontail rabbits tended to shed higher levels of the Northern pintail virus as compared to the gyrfalcon virus during early DPI.

Similar to previous studies involving relatively high inoculation doses [12], the viral shedding dynamics in the current study were fairly brief, with most animals ceasing viral shedding by 6–7 DPI. Nonetheless, one animal shed virus through 8 DPI (the last day samples were taken) and shed relatively high titers out to 6 DPI ($10^{4.6}$ PFU/mL by the nasal route and $10^{3.4}$ PFU/mL by the oral route). For obvious reasons, cottontail rabbits that shed higher titers and shed for longer periods of time likely pose a greater epidemiological risk in most situations.

As has been reported previously for multiple LP IAV strains [12, 14], the current study indicates that cottontail rabbits can shed relatively high levels of some HP IAVs along with a lack of any obvious signs of disease. As a symptomless carrier of IAVs, if infected, cottontail rabbits are unlikely to provide any type of obvious warning of the presence of a virus on poultry facilities without the use of active surveillance. Thus, reducing wildlife attractants and preventing wildlife access to facility premises, such as poultry houses, are key to mitigating the potential risk posed by this and certain other wildlife species [17].

Acknowledgments We thank the National Wildlife Research Center (NWRC) animal care staff for excellent animal care, J. Ellis and M. Grady for assistance with animal capture/processing, J. Ellis for assistance with AGID assays, S. Shriner for assistance with figures, and M. Torchetti for facilitating virus transfer. The opinions and conclusions of this article are those of the authors and do not necessarily represent those of the U.S. Department of Agriculture (USDA) or Colorado State University. The mention of commercial products herein is for identification purposes only and does not constitute endorsement or censure.

Funding This research was supported by the U.S. Department of Agriculture.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

References

- Achenbach JE, Bowen RA (2011) Transmission of avian influenza A viruses among species in an artificial barnyard. *PLoS ONE* 6:e17643
- Brown JD, Stallknecht DE, Berghaus RD, Swayne DE (2009) Infectious and lethal doses of H5N1 highly pathogenic avian influenza virus for house sparrows (*Passer domesticus*) and rock pigeons (*Columbia livia*). *J Vet Diag Invest* 21:437–445
- DecisionInnovationSolutions (2015) Economic impact of highly pathogenic avian influenza (HPAI) on poultry in Iowa. Document prepared for Iowa Farm Bureau. pp 1–16
- Forrest HL, Kim JK, Webster RG (2010) Virus shedding and potential for interspecies waterborne transmission of highly pathogenic H5N1 influenza virus in sparrows and chickens. *J Virol* 84:3718–3720
- Hall JS, Bentler KT, Landolt G, Elmore SA, Minnis RB, Campbell TA, Barras SC, Root JJ, Pilon J, Pablonia K, Driscoll C, Slate D, Sullivan H, McLean RG (2008) Influenza infection in wild raccoons. *Emerg Infect Dis* 14:1842–1848
- Ip HS, Torchetti MK, Crespo R, Kohrs P, DeBruyn P, Mansfield KG, Baszler T, Badcoe L, Bodenstern B, Shearn-Bochsler V, Killian ML, Pedersen JC, Hines N, Gidlewski T, DeLiberto T, Sleeman JM (2015) Novel Eurasian highly pathogenic influenza A H5 viruses in wild birds, Washington, USA, 2014. *Emerg Infect Dis* 21:886–890
- Kaplan BS, Russier M, Jeevan T, Marathe B, Govorkova EA, Russell CJ, Kim-Torchetti M, Choi YK, Brown I, Saito T, Stallknecht DE, Krauss S, Webby RJ (2016) Novel highly pathogenic avian A(H5N2) and A(H5N8) influenza viruses of clade 2.3.4.4 from North America have limited capacity for replication and transmission in mammals. *MSphere* 1:00003–00016
- Kaplan BS, Torchetti MK, Lager KM, Webby RJ, Vincent AL (2017) Absence of clinical disease and contact transmission of HPAI H5NX clade 2.3.4.4 from North America in experimentally infected pigs. *Influenza Respir Viruses* 11:464–470
- Nemeth NM, Thomas NO, Orathood DS, Anderson TD, Oesterle PT (2010) Shedding and serologic responses following primary and secondary inoculation of house sparrows (*Passer domesticus*) and European starlings (*Sturnus vulgaris*) with low-pathogenicity avian influenza virus. *Avian Pathol* 39:411–418
- NVSL (2003) Avian influenza agar gel immunodiffusion test to detect serum antibodies to type A influenza viruses. NVSL document AVPRO0100.05
- Pulit-Penalzo JA, Sun X, Creager HM, Zeng H, Belser JA, Maines TR, Tumpey TM (2015) Pathogenesis and transmission of novel highly pathogenic avian influenza H5N2 and H5N8 viruses in ferrets and mice. *J Virol* 89:10286–10293
- Root JJ, Shriner SA, Bentler KT, Gidlewski T, Mooers NL, Spraker TR, VanDalen KK, Sullivan HJ, Franklin AB (2014) Shedding of a low pathogenic avian influenza virus in a common synanthropic mammal—the cottontail rabbit. *PLoS ONE* 9:e103513
- Root JJ, Shriner SA, Ellis JW, VanDalen KK, Sullivan HJ, Franklin AB (2015) When fur and feather occur together: interclass transmission of avian influenza A virus from mammals to birds through common resources. *Sci Rep* 5:12354
- Root JJ, Bosco-Lauth AM, Bielefeldt-Ohmann H, Bowen RA (2016) Experimental infection of peridomestic mammals with emergent H7N9 (A/Anhui/1/2013) influenza A virus: implications for biosecurity and wet markets. *Virology* 487:242–248
- Root JJ, Shriner SA, Ellis JW, VanDalen KK, Sullivan HJ (2017) Low viral doses are sufficient to infect cottontail rabbits with avian influenza A virus. *Arch Virol* 162:3381–3388
- Shriner SA, VanDalen KK, Mooers NL, Ellis JW, Sullivan HJ, Root JJ, Franklin AB (2012) Low-pathogenic avian influenza viruses in wild house mice. *PLoS ONE* 7:e39206
- Shriner SA, Root JJ, Lutman MW, Kloft JM, VanDalen KK, Sullivan HJ, White TS, Milleson MP, Chandler SC, Wolf PC, Turnage CT, McCluskey BJ, Vincent AL, Torchetti MK, Gidlewski T, Deliberto TJ (2016) Surveillance for highly pathogenic H5 avian influenza A virus in synanthropic wildlife associated with poultry facilities during an acute outbreak. *Sci Rep* 6:36237
- Su S, Xing G, Wang J, Li Z, Gu J, Yan L, Lei J, Ji S, Hu B, Gray GC, Yan Y, Zhou J (2016) Characterization of H7N2 avian influenza virus in wild birds and pikas in Qinghai-Tibet Plateau area. *Sci Rep* 6:30974
- Yan Y, Gu JY, Yuan ZC, Chen XY, Li ZK, Lei J, Hu BL, Yan LP, Xing G, Liao M, Zhou JY (2017) Genetic characterization of H9N2 avian influenza virus in plateau pikas in the Qinghai Lake region of China. *Arch Virol* 162:1025–1029
- Yu Z, Cheng K, Sun W, Xin Y, Cai J, Ma R, Zhao Q, Li L, Huang J, Sang X, Li X, Zhang K, Wang T, Qin C, Qian J, Gao Y, Xia X (2014) Lowly pathogenic avian influenza (H9N2) infection in plateau pika (*Ochotona curzoniae*), Qinghai Lake, China. *Vet Microbiol* 173:132–135
- Zhou J, Sun W, Wang J, Guo J, Yin W, Wu N, Li L, Yan Y, Liao M, Huang Y, Luo K, Jiang X, Chen H (2009) Characterization of the H5N1 highly pathogenic avian influenza virus derived from wild pikas in China. *J Virol* 83:8957–8964