

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and
Plant Health Inspection Service

2018

Nutritional depletion of total mixed rations by red-winged blackbirds and projected impacts on dairy cow performance

James C. Carlson

USDA APHIS Wildlife Services, james.c.carlson@aphis.usda.gov

Randal S. Stahl

USDA APHIS Wildlife Services

John J. Wagner

Colorado State University

Terry E. Engle

Colorado State University

Shelagh T. DeLiberto

USDA APHIS Wildlife Services, shelagh.t.deliberto@aphis.usda.gov

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Life Sciences Commons](#)

Carlson, James C.; Stahl, Randal S.; Wagner, John J.; Engle, Terry E.; DeLiberto, Shelagh T.; Reid, Dustin A.; and Werner, Scott J., "Nutritional depletion of total mixed rations by red-winged blackbirds and projected impacts on dairy cow performance" (2018). *USDA National Wildlife Research Center - Staff Publications*. 2172.

https://digitalcommons.unl.edu/icwdm_usdanwrc/2172

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

James C. Carlson, Randal S. Stahl, John J. Wagner, Terry E. Engle, Shelagh T. DeLiberto, Dustin A. Reid, and Scott J. Werner

Nutritional depletion of total mixed rations by red-winged blackbirds and projected impacts on dairy cow performance

James C. Carlson¹, Randal S. Stahl¹, John J. Wagner², Terry E. Engle², Shelagh T. DeLiberto¹, Dustin A. Reid¹ and Scott J. Werner^{1*}

¹U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services, National Wildlife Research Center, 4101 LaPorte Avenue, Fort Collins, Colorado, USA

²Department of Animal Science, Colorado State University, College of Agricultural Sciences, Fort Collins, Colorado, USA

Received 2 November 2017; accepted for publication 15 June 2018

This Research Communication describes an investigation of the nutritional depletion of total mixed rations (TMR) by pest birds. We hypothesized that species-specific bird depredation of TMR can alter the nutritional composition of the ration and that these changes can negatively impact the performance of dairy cows. Blackbirds selected the high energy fraction of the TMR (i.e., flaked corn) and reduced starch, crude fat and total digestible nutrients during controlled feeding experiments. For Holsteins producing 37.1 kg of milk/d, dairy production modeling illustrated that total required net energy intake (NE_i) was 35.8 Mcal/d. For the reference TMR unexposed to blackbirds and the blackbird-consumed TMR, NE_i supplied was 41.2 and 37.8 Mcal/d, and the resulting energy balance was 5.4 and 2.0 Mcal/d, respectively. Thus, Holsteins fed the reference and blackbird-consumed TMR were estimated to gain one body condition score in 96 and 254 d, and experience daily weight change due to reserves of 1.1 and 0.4 kg/d, respectively. We discuss these results in context of an integrated pest management program for mitigating the depredation caused by pest birds at commercial dairies.

Keywords: *Agelaius phoeniceus*, dairy production, bird damage management, nutrition.

Multiple species of cohabitant birds, including red-winged blackbirds (*Agelaius phoeniceus*) and European starlings (*Sturnus vulgaris*), congregate in large flocks and exploit the abundant and nutritious food sources found at dairies and feedlots (Besser et al. 1968; Dolbeer et al. 1978). Estimates of bird damage in commercial dairies in Wisconsin, New York and Pennsylvania suggest that red-winged blackbirds were one of the most common birds species associated with depredation (Shwiff et al. 2012). Red-winged blackbirds have been previously observed to most frequently consume corn, averaging 30% of their diet (White et al. 1985). In contrast, European starlings offered individual components of total mixed rations (TMR) preferentially selected Propel[®] energy nugget (Nestle Purina, St. Louis, Missouri, USA), averaging 49% of their daily diet (Carlson et al. 2018).

The purpose of this study was to investigate the nutritional depletion of TMR by red-winged blackbirds and their associated impacts to dairy production. We previously estimated

the nutritional depletion of TMR by European starlings (Carlson et al. 2018). Our estimates from the NRC (2001) production model suggested that cows fed TMR exposed to starlings would lose one body condition score (BCS) in 91 d and cows fed the reference TMR (unexposed to starling depredation) would lose one BCS in 161 d (Carlson et al. 2018). We hypothesized that species-specific bird depredation of TMR can alter the nutritional composition of the ration and that these changes can negatively impact the performance of dairy cows. We therefore replicated our controlled feeding experiment and dairy production modeling with red-winged blackbirds offered individual TMR components and a high-energy TMR to investigate the nutritional depletion of dairy rations by red-winged blackbirds and the associated impacts on dairy cow performance.

Materials and methods

We conducted a controlled feeding experiment with red-winged blackbirds offered individual TMR components. On each of four test days, we offered blackbirds within

*For correspondence; e-mail: Scott.J.Werner@aphis.usda.gov

group cages ($n = 10$ cages; five birds per cage) 100 g of eight individual TMR components including steamed-flaked corn, Propel[®] energy nugget, canola meal, cracked corn, ground corn, soybean meal, cotton seed and lactating mineral. All components were sourced from a commercial dairy located in Larimer County, Colorado 80538. Blackbird consumption of individual TMR components is reported as percent consumed throughout the 4-d test. The live-capture, animal care and use procedures were approved by the Institutional Animal Care and Use Committee of the National Wildlife Research Center (QA-2369, J.C. Carlson-Study Director).

For the purpose of estimating the nutritional offsets caused by red-winged blackbirds, we subsequently offered a high energy TMR (30–160 d in milk) to experimentally-naïve blackbirds within group cages ($n = 10$ cages; 5 birds per cage, 1 kg TMR per cage) for four consecutive days. Parameterization of the dairy production model (NRC, 2001) was based upon animal condition and feed formulation data provided by a commercial dairy in northern Colorado, nutrition data provided by Cumberland Valley Analytical Services (CVAS, Hagerstown, Maryland USA) and our component preference data.

All nutritional data were analyzed using ANOVA within mixed linear models (Proc Mixed, SAS 9.2, SAS Institute Inc., Cary, North Carolina USA). Fixed effects included treatment status (i.e., reference and blackbird-consumed rations) and cage was included as a random effect. Denominator degrees of freedom were calculated using the Satterthwaite approximation. We controlled for false discoveries using the Benjamini-Hochberg procedure (Benjamini & Hochberg, 1995). For all analyses, the false discovery rate was set at $\alpha = 0.05$. Univariable analyses were ranked by P -value from smallest (1) to largest (m). Cutoff values for the rejection of null hypotheses were calculated as $(\text{rank}/m) \times \alpha$.

Results and discussion

Red-winged blackbirds preferred TMR components with high starch and high fat content. Among all foods consumed, various corn products (flaked, cracked and ground corn) accounted for 78% of all foods consumed by red-winged blackbirds during the preference experiment. Blackbirds primarily consumed flaked corn (42%) and the energy nugget (13%), and cracked corn and ground corn comprised 4 and 1% of blackbird consumption, respectively. Each of the other TMR components comprised <0.5% of blackbird consumption. Thus, blackbirds are likely attracted to concentrated animal feeding operations because of specific nutrients associated with expensive feed components (e.g., processed corn, fat supplements).

The preference and consumption of various TMR components by pest birds may significantly alter the cost to producers and the decision-making for bird damage management. The preference of red-winged blackbirds for steam-flaked corn (42% consumption) was greater than that

observed for Propel[®] energy nugget, canola meal, cracked corn, ground corn, soybean meal, cotton seed and lactating mineral. In contrast, the preference of European starlings for Propel[®] energy nugget (49% consumption) was greater than that observed for steam-flaked corn, corn gluten, dry distillers grains, canola meal, corn silage and mineral supplement (Carlson et al. 2018). Thus, species-specific data for offending bird species are necessary for (1) the generation of accurate estimates of bird damage and (2) the prescription and implementation of cost-effective damage management strategies.

The nutritional data illustrated that red-winged blackbird consumption significantly altered the nutritional characteristics of TMR (Table 1). Relative to the reference rations unexposed to blackbirds, blackbird-consumed rations had less net energy for lactation ($P < 0.0001$), maintenance ($P < 0.0001$) and gain ($P < 0.0001$). Blackbird-consumed rations also had lower concentrations of starch ($P < 0.0001$), crude fat ($P < 0.0001$) and total digestible nutrients ($P < 0.0001$), and greater concentrations of acid detergent fiber ($P < 0.0017$), neutral detergent fiber ($P < 0.0005$), potassium ($P < 0.0001$) and calcium ($P < 0.0017$) compared to the reference rations. Relative to reference rations, we observed a 9.5% reduction in corn silage, a 9.6% reduction in energy nugget, a 23.6% reduction in ground corn and a 24.7% reduction in steam-flaked corn in TMR exposed to red-winged blackbirds. These results enabled us to identify metabolizable energy sources that need to be excluded from depredating blackbirds.

Dairy production modeling (NRC, 2001) illustrated that cow performance can be negatively impacted by blackbird consumption of TMR (Table 2). For Holsteins producing 37.1 kg of milk/d, total required net energy intake (NE_I) was 35.8 Mcal/d. For the reference TMR, NE_I supplied was 41.2 Mcal/d and for the blackbird-consumed TMR NE_I supplied was 37.8 Mcal/d. The resulting energy balance for reference and blackbird-consumed rations was 5.4 and 2.0 Mcal/d, respectively. Thus, blackbird depredation of cattle feed can reduce dairy production through the nutritional depletion of TMR.

We also observed species-specific impacts of pest birds on dairy cow performance. Body condition scores integrate the influences of many nutritional aspects of TMR to distinguish differences in the fat reserves, or the nutritional needs among individual cows. The high-energy reference TMR used for this blackbird experiment would cause the gain of one BCS in 96 d (vs. 255 d for the blackbird-exposed TMR; Table 2) for cows producing 37.1 kg of milk per day ($NE_I = 35.8$ Mcal/d). In contrast, the late-lactation reference TMR used for our starling feeding experiments would cause the loss of one BCS in 161 d (vs. 91 d for the starling-exposed TMR) for cows producing 31.75 kg of milk per day ($NE_I = 31.5$ Mcal/d; Carlson et al. 2018). These TMR-specific differences notwithstanding, the absolute difference in BCS was 159 d in this blackbird study and 70 d in the starling study (Carlson et al. 2018). Because fat supplements are approximately \$1.10 per kg (USD) and corn products are approximately \$0.14 per kg, we conclude that starlings can have a greater monetary impact to dairy

Table 1. Nutritional analysis of reference and blackbird-consumed total mixed rations

Variable	Metric	Reference mean (X)	Blackbird mean (X)	P-value	Rank [†]	Cutoff value [‡]
Net Energy Lactation	Mcal/kg	1·839	1·700	<0·0001	4·5	0·0102
Net Energy Maintenance	Mcal/kg	1·825	1·739	<0·0001	4·5	0·0102
Net Energy Gain	Mcal/kg	1·197	1·120	<0·0001	4·5	0·0102
Starch	%DM	31·128	28·221	<0·0001	4·5	0·0102
Crude Fat	%DM	5·482	4·625	<0·0001	4·5	0·0102
Potassium	%DM	1·159	1·252	<0·0001	4·5	0·0102
Total Digestible Nutrients	%DM	76·184	73·459	<0·0001	4·5	0·0102
Non Fiber Carbohydrates	%DM	42·634	40·25	<0·0001	4·5	0·0102
Neutral Detergent Fiber	%DM	24·825	26·725	0·0005	9·0	0·0205
Acid Detergent Fiber	%DM	16·134	17·793	0·0017	10·5	0·0239
Calcium	%DM	1·179	1·283	0·0017	10·5	0·0239
Ash	%DM	7·724	8·195	0·002	12	0·0273
Crude Protein	%DM	19·347	20·203	0·0028	13	0·0295
Phosphorus	%DM	0·448	0·468	0·0031	14	0·0318
Copper	PPM	30·282	31·938	0·0073	15	0·0341
Magnesium	%DM	0·376	0·396	0·0317	16	0·0364
Manganese	PPM	136·125	142·373	0·0356	17	0·0386
Zinc	PPM	253·186	279·656	0·0691	18	0·0409
Dry Matter	%DM	96·409	96·543	0·1994	19·5	0·0443
Moisture	%DM	3·591	3·456	0·1994	19·5	0·0443
Sodium	%DM	0·444	0·451	0·4531	21	0·0477
Iron	PPM	361·218	361·531	0·9863	22	0·05

[†]Rank order of P-values from analyses of nutritional cattle feed samples

[‡]Benjamini Hochberg cutoff values for rejection of null hypotheses

Table 2. National Research Council, dairy production model output for reference and blackbird-consumed total mixed rations

NRC model output [†]	Reference TMR	Blackbird-consumed TMR
NE _i required (Mcal/d)	35·8	35·8
NE _i supplied (Mcal/d)	41·2	37·8
NE _i balance (Mcal/d)	5·4	2·0
Days to gain one BCS	96	255
Weight change (kg/d)	1·1	0·4
RDP required (g/d)	2541	2312
RDP supplied (g/d)	2584	2524
RDP balance (g/d)	42	212
RUP required (g/d)	1300	1395
RUP supplied (g/d)	1853	1689
RUP balance (g/d)	552	294
MP – bacterial (g/d)	1383	1258
MP – RUP (g/d)	1447	1305
MP – endogenous (g/d)	120	109

[†]RDP and RUP are rumen degradable protein and rumen undegradable protein, respectively. MP is microbial protein

production than red-winged blackbirds. We have previously observed the colonization of thousands of blackbirds subsequent to the lethal control of starlings at particular dairies. Thus, the impacts of pest birds and the cost-effective management associated with these impacts are dependent on the species composition and the density of the pest-bird community at a particular dairy operation.

Because pest birds may not nutritionally deplete all TMR consumed by cows each day (Carlson et al. 2018), our

results may overestimate the nutritional depletion caused by blackbirds at commercial dairies. The density of blackbirds in our controlled feeding experiments is comparable to more than 100 blackbirds per cow under field conditions. Most commercial dairies will likely not experience 100 pest birds per cow nor the associated extent of nutritional depletion as estimated by our study. Our approach and our results, however, demonstrate that (1) we can estimate the species-specific nutritional depletion caused by pest birds and (2) dairy production models can be used to predict production losses to dairies impacted by bird depredation.

Similar to European starlings (Glahn et al. 1983), red-winged blackbirds are apparently influenced by the form and size of livestock feed. For example, blackbirds strongly preferred flaked corn over ground corn in our feeding experiments. Previous research has shown that European starlings could not consume $\geq 1\cdot27$ cm-diameter (Deppenbusch et al. 2011) extruded feed pellets. Altering the size of flaked corn and fat nugget, or combining highly desired components into approximately 1·27 mm-diameter pellets or finely milled particles may be a non-lethal component of integrated pest management strategies for the mitigation of the species-specific impacts of bird depredation (Twedt & Glahn, 1982; Glahn et al. 1983; Deppenbusch et al. 2011; Carlson et al. 2018).

References

Benjamini Y & Hochberg Y 1995 Controlling for false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society* 57 289–300

- Besser JF, DeGrazio JW & Guarino JL** 1968 Costs of wintering starlings and red-winged blackbirds at feedlots. *Journal of Wildlife Management* **32** 179–180
- Carlson JC, Stahl RS, DeLiberto ST, Wagner JJ, Engle TE, Engeman RM, Olson CS, Ellis JW & Werner SJ** 2018 Nutritional depletion of total mixed rations by European starlings: projected effects on dairy cow performance and potential intervention strategies to mitigate damage. *Journal of Dairy Science* **101** 1777–1784
- Depenbusch BE, Drouillard JS & Lee CD** 2011 Feed depredation by European starlings in a Kansas feedlot. *Human-Wildlife Interactions* **5** 58–65
- Dolbeer RA, Wornecki PA, Strickley JR & White SB** 1978 Agricultural impact of a winter population of blackbirds and starlings. *Wilson Bulletin* **90** 31–44
- Glahn JF, Twedt DJ & Otis DL** 1983 Estimating feed loss from starling use of livestock feed troughs. *Wildlife Society Bulletin* **11** 366–372
- National Research Council (NRC)** 2001 *Nutrient Requirement of Dairy Cattle*, Seventh edition. National Research Council, National Academy Press, Washington, DC
- Shwiff SA, Carlson JC, Glass H, Suckow J, Lowney MS, Moxcey KM, Larson B & Linz GM** 2012 Producer survey of bird livestock interactions in commercial dairies. *Journal of Dairy Science* **95** 6820–6829
- Twedt DJ & Glahn JF** 1982 Reducing starling depredation at livestock feeding operations through changes in management practices. *Proceedings of the Vertebrate Pest Conference* **10** 159–163
- White SB, Dolbeer RA & Bookhout TA** 1985 Ecology, bioenergetics, and agricultural impacts of a winter-roosting population of blackbirds and starlings. *Wildlife Monographs* **93** 1–42