

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

4-10-2015

Prevalence of internal parasites in beef cows in the United States: Results of the National Animal Health Monitoring System's (NAHMS) beef study, 2007–2008

Bert E. Stromberg

Louis C. Gasbarre

Lora R. Ballweber

David A. Dargatz

Judith M. Rodriguez

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Bert E. Stromberg, Louis C. Gasbarre, Lora R. Ballweber, David A. Dargatz, Judith M. Rodriguez, Christine A. Koprak, and Dante Zarlenga

Prevalence of internal parasites in beef cows in the United States: Results of the National Animal Health Monitoring System's (NAHMS) beef study, 2007–2008

Bert E. Stromberg, Louis C. Gasbarre, Lora R. Ballweber, David A. Dargatz, Judith M. Rodriguez, Christine A. Koprak, Dante S. Zarlenga

Abstract

During the United States Department of Agriculture (USDA) National Animal Health Monitoring System's (NAHMS) 2007–2008 beef study, 567 producers from 24 US States were offered the opportunity to collect fecal samples from weaned beef calves and have them evaluated for the presence of parasite eggs (Phase 1). Participating producers were provided with instructions and materials for sample collection. Up to 20 fresh fecal samples were collected from each of the 99 participating operations. Fresh fecal samples were submitted to one of 3 randomly assigned laboratories for evaluation. Upon arrival at the laboratories, all samples were processed for the enumeration of strongyle, *Nematodirus*, and *Trichuris* eggs using the modified Wisconsin technique. The presence or absence of coccidian oocysts and tapeworm eggs was also noted. In submissions where the strongyle eggs per gram exceeded 30, aliquots from 2 to 6 animals were pooled for DNA extraction. Extracted DNA was subjected to genus level polymerase chain reaction (PCR) identification for the presence of *Ostertagia*, *Cooperia*, *Haemonchus*, *Oesophagostomum*, and *Trichostrongylus*. In this study, 85.6% of the samples had strongyle type, *Nematodirus*, and *Trichuris* eggs. Among the samples evaluated, 91% had *Cooperia*, 79% *Ostertagia*, 53% *Haemonchus*, 38% *Oesophagostomum*, 18% *Nematodirus*, 7% *Trichuris*, and 3% *Trichostrongylus*. The prevalence of coccidia and tapeworm eggs was 59.9% and 13.7%, respectively.

Résumé

Pendant l'étude de 2007–2008 chez les bovins effectuée par le Système national de surveillance des maladies animales (NAHMS) du Département de l'agriculture des États-Unis (USDA), 567 producteurs provenant de 24 états américains se sont vus offrir l'opportunité de prélever des échantillons de fèces de veaux sevrés et de les faire analyser pour la présence d'œufs de parasite (Phase 1). On a fourni aux producteurs participants les instructions et le matériel pour le prélèvement d'échantillon. Jusqu'à 20 échantillons de fèces fraîches furent prélevés de chacune des 99 opérations participantes. Les échantillons de fèces fraîches furent soumis de manière aléatoire pour évaluation à l'un des trois laboratoires participants. Suite à l'arrivée au laboratoire, tous les échantillons étaient traités pour énumération des strongles, de *Nematodirus*, et d'œufs de *Trichuris* en utilisant la technique de Wisconsin modifiée. La présence ou l'absence d'œufkystes de coccidie et d'œufs de vers plats furent également notées. Dans les échantillons soumis et dont le nombre d'œufs de strongles par gramme dépassait 30, des aliquots de 2 à 6 animaux étaient regroupés pour extraction de l'ADN. L'ADN extrait était soumis à une réaction d'amplification en chaîne par la polymérase (PCR) pour une identification au genre de la présence d'*Ostertagia*, de *Cooperia*, d'*Haemonchus*, d'*Oesophagostomum*, et de *Trichostrongylus*. Dans la présente étude, 85,6 % des échantillons avaient des strongles, du *Nematodirus*, et des œufs de *Trichuris*. Parmi les échantillons évalués, 91 % avaient du *Cooperia*, 79 % de l'*Ostertagia*, 53 % de l'*Haemonchus*, 38 % de l'*Oesophagostomum*, 18 % du *Nematodirus*, 7 % du *Trichuris*, et 3 % du *Trichostrongylus*. Les prévalences de coccidies et d'œufs de vers plats étaient respectivement de 59,9 % et 13,7 %.

(Traduit par Docteur Serge Messier)

College of Veterinary Medicine, University of Minnesota, St. Paul, Minnesota (Stromberg); United States Department of Agriculture (USDA), Agricultural Research Service (ARS), Bovine Functional Genomics Lab, Beltsville, Maryland (Gasbarre); Colorado State University, Veterinary Diagnostic Laboratory, Fort Collins, Colorado (Ballweber); USDA, Animal and Plant Health Inspection Service (APHIS), Veterinary Service (VS) Centers for Epidemiology and Animal Health, Fort Collins, Colorado (Dargatz, Rodriguez, Koprak); USDA, ARS, Animal Parasitic Diseases Lab, Beltsville, Maryland, USA (Zarlenga).

Address all correspondence to Dr. Zarlenga; telephone: 301-504-8754; fax: 301-504-8979; e-mail: dante.zarlenga@ars.usda.gov

Dr. Gasbarre's current address is Gasbarre Consulting, Buffalo, Wyoming 82834, USA.

Received March 6, 2015. Accepted April 10, 2015.

U.S. government works are not subject to copyright.

Introduction

Internal parasites, primarily helminths, are common pathogens of the cattle industry and substantially impact the economics of cattle production worldwide. Helminths can reduce the reproductive performance of the herd, reduce weaning weights, and in general, negatively impact animal health. These negative impacts are due to the destructive effects directly on host tissues; indirect responses, such as suppression of the host immune responses; or the loss of appetite (1), which can impact all aspects of animal well-being.

Data from the United States Department of Agriculture (USDA) National Health Monitoring System's (NAHMS) 2007–2008 beef study show that cow/calf producers consider parasites to be a major problem in production (Ballweber et al, unpublished data). Previous studies in stocker cattle (2) showed a significant difference in weight gain in the parasite free (drug-treated) groups relative to non-treated groups, ranging from 0.132 to 0.272 kg average daily gain (ADG) for the various treatment groups. In a more confined study involving a beef cow/calf operation over a 2-year period, Stromberg, et al (3) demonstrated a negative effect of gastrointestinal (GI) nematodes on productivity where the calves of treated cows had a mean weaning weight of 18.5 kg over the non-medicated group. There was also a 12% improvement in reproductive performance for the nematode-free cows. As such, the presence of helminth parasites generally equates to a negative impact on production; however, data are lacking that define the magnitude of this economic impact.

In a 1993 cattle parasite survey that sampled over 5500 herds, Myers and Keith (4) found that calves, yearlings, cows, and bulls had > 90% prevalence of helminths in the western, northern, and southern regions of the United States (US). Similar findings have been noted in beef cattle outside of the US. A survey using tracer calves and necropsies in the cerrado region of Brazil (5), culled cows in Ireland (6), and a serological survey of replacement stock in The Netherlands (7) all showed a substantial prevalence of parasites.

Inferences of parasite importance and/or prevalence on animal production are often predicated on studies done on a single farm/ranch. In some cases the parasites are identified and/or quantified from animals using fecal egg counts and coproculture; however, tracer calves grazed with the herd have also been used where adult parasites are recovered and morphologically identified. These narrowly focused studies though important, provide prevalence data that are confined to a particular state or region of the state. The primary objective of this study was to evaluate parasite prevalence in weaned beef calves throughout the major cow-calf producing regions of the US. These data are presented herein.

Materials and methods

Study locations

A stratified random sample of over 4000 operations was chosen by the USDA's National Agricultural Statistics Service (NASS) for initial contact. The 567 beef cow/calf operations that participated in a national study of animal health and management conducted by the USDA NAHMS were provided an opportunity to collect fecal samples from weaned calves for evaluation of parasite burdens.

This population of operations has been described previously (8). Briefly, the eligible production facilities consisted of a stratified random sample of operations in 24 States with the largest beef cow populations that had 1 or more beef cows on October 1, 2007, and agreed to respond to a questionnaire administered during an interview. All participants remained anonymous within the NAHMS reporting system.

Sample collection

Producers interested in participating in the parasite survey were provided with the necessary materials and instructions to collect and ship the samples. Sample collection from weaned calves occurred at the discretion of the producer from March 1 through December 2, 2008. All animals were 6 to 18 mo of age, had grazed on pasture for at least 4 wk prior to the collection, and had not been treated with an anthelmintic in the previous 45 d. Fecal samples were collected from up to 20 calves (9) either directly from the rectum or from fresh fecal pats. Producers were asked to collect "golf ball" sized samples into an unused examination glove, transfer these into individual plastic bags and refrigerate overnight. The chilled samples were shipped with ice packs to one of 3 randomly assigned laboratories participating in the study: Colorado State University, the USDA, ARS, Beltsville, or the University of Minnesota.

Laboratory counting procedures

Upon arrival at the laboratory, samples were logged in and refrigerated. The eggs present in all fecal samples were quantified using the Wisconsin Double Centrifugal Floatation or the Modified Wisconsin technique (10,11). Each laboratory conducted the test according to routine procedures used in that laboratory. Parasite eggs were counted and identified as strongyle type, *Nematodirus*, or *Trichuris*. Samples were also evaluated as positive or negative for coccidia oocysts and *Moniezia* eggs.

Data collection

Information on routine deworming practices for the operation was also available from a previously administered questionnaire. At the time of sample collection, producers were asked to complete a short questionnaire about the group of animals being sampled. The requested information included the number of animals in the group sampled, any prior treatment with anthelmintics, and the anthelmintic used. Egg count data were recorded and transmitted to USDA, NAHMS, for dissemination to the owners and for statistical analysis.

Genus identification by polymerase chain reaction (PCR)

Samples having strongyle egg counts > 30 were collected and pooled for genus identification. Following zinc sulfate flotation (12), eggs were washed and frozen at -80°C in 0.5 mL PCR tubes in a minimal volume of water; each sample had no less than 100 eggs. The tubes were transferred uncapped and frozen to a thermal cycler (Perkin Elmer DNA Thermal Cycler 480; Waltham, Massachusetts, USA) preheated to 95°C , incubated a minimum of 15 min, and thereafter until the sample volumes were reduced to 20 μL . The eggs were subsequently treated as described in the Tissue and Hair Extraction kit (Promega Biotec; Madison, Wisconsin, USA). Released DNA was

Table I. Number of responding operations by geographic region

Region	Phase 1: Operations completing 1st questionnaire	Phase IIA: Operations completing 2nd questionnaire	Phase IIB: Operations completing 3rd questionnaire and eligible for sample collection	Phase III: Operations submitting useable sample for evaluation
West	370	138	105	26
Central	612	196	175	45
South Central ^a	483	233	190	28
East ^a	694			
Total	2159	567	470	99

^a Regions were combined for Phases II and III of survey.

Table II. Number of responding operations by herd size

Herd size	Phase 1: Operations completing 1st questionnaire	Phase IIA: Operations completing 2nd questionnaire	Phase IIB: Operations completing 3rd questionnaire and eligible for sample collection	Phase III: Operations submitting useable sample for evaluation
1–49	819	163	127	18
50–99	386	96	81	17
100–199	381	125	104	26
200+	573	183	158	38
Total	2159	567	470	99

for each parasite was calculated by region, by quarter, and overall. Regional and quarterly operation-level percentages were compared using Chi-squared tests (SAS Proc Freq; SAS Institute).

Results

Overall 291 producers indicated interest in participating in the parasite prevalence component of the study and ordered sample collection materials. However, only 102 (35%) operations actually submitted samples. Three producers submitted samples that were discarded because they did not follow collection guidelines. These included operations where the animals were too young, where collection took place too soon (< 45 d) after the last anthelmintic treatment, or where the samples were held and not shipped within 24 to 48 h of collection. Thus 99 operations were used in the analysis and accounted for 1772 fecal samples.

Overall 85.6% (95% CI = 80.9, 89.6) of the samples collected had strongyle type eggs, 18.0% (14.2, 22.8) had *Nematodirus* spp., and 7.1% (5.3, 9.8) had *Trichostrongylus* spp. eggs (Figure 2). When these data were sub-divided into the months of year in which they were collected (March–May, June–August, and September–December), significant differences were observed between collection periods for *Nematodirus* ($P = 0.031$) and *Trichostrongylus* ($P = 0.010$). In addition, 59.9% (53.1, 67.0) of the samples contained coccidian oocysts and 13.7% (10.8, 18.4) had tapeworm eggs (Figure 2).

All operations exhibited cattle shedding strongyle eggs in each collection period and for all regions. Mean strongyle shedding was

32.5 epg (25.3, 43.4) over all samples. Regional differences were observed with 30.3 (22.7, 44.7), 58.1 (36.2, 85.3), and 11.3 (7.5, 18.8) epg for Central, South, and West, respectively ($P < 0.0001$). The overall average for *Nematodirus* spp. was 0.8 epg (0.5, 1.2) and there was a significant difference between regions, with the South mean epg less than that of the Central and West regions ($P = 0.0003$). The mean epg for *Trichostrongylus* spp. was 0.15 (0.10, 0.23), with no differences between regions. The genera present identified by PCR were *Cooperia* spp. (91%), *Ostertagia* (79%), *Haemonchus* (53%), *Oesophagostomum* (38%), and *Trichostrongylus* (3%), and by egg morphology, *Nematodirus* spp (18%), and *Trichostrongylus* (7.1%).

Discussion

Anthelmintic use rather than pasture management has become the dominant strategy for controlling GI nematodes of cattle (1). The introduction of thiabendazole followed by other benzimidazoles changed the mindset in the way producers managed parasite infections. This reliance on pharmaceuticals was extended with the introduction and extensive use of the avermectins/milbemycins (1). One would expect that the extensive use of anthelmintics over the past 40 y would have reduced parasite prevalence across the US. However, there have been few large-scale studies evaluating the impact of anthelmintic therapy on the populations of parasites throughout the US.

Historically, most studies have reported parasite prevalence from one or a few farms in a specific state or region. In this study,




Figure 2. Percent of positive samples relative to time period of survey: 1 = Mar-May; 2 = Jun-Aug; 3 = Sep-Dec; All = Mar-Dec.

cow/calf operations in 24 States, representing 87.8% (28.6 million) of the US beef inventory and 79.6% (603 000 operations) of the cow/calf inventory were offered the opportunity to assess parasite burdens and types. Among the 2159 operations initially contacted by NASS, 567 agreed to have their contact information forwarded to APHIS and of these 470 became eligible for the parasite sampling activity. Within this group, 291 ordered kits for parasite sampling and 99 operations from 21 States collectively shipped a total of 1772 samples for analysis. The remaining States were either not a source of cow/calf operations in the US or had no operations wishing to pursue sample analysis. Clearly, the issue of representative sampling and bias comes into play within such a study where approximately 5% of those operations initially contacted, eventually agreed to ship and comply with proper sampling procedures for analysis. Among common reasons for lack of participation were the burden of sample collection/submission/follow through, confidentiality despite assurance of anonymity, and insufficient interest in parasites. As such, this survey represents a convenience sample and really cannot be used as an estimate for the original national population. That being said, the survey still encompassed operations from 21 of the 24 States initially contacted. Further, the operations that agreed to participate spanned the full range of herd size (Table II) with a predilection toward operations greater than 100 head of cattle.

The concept of representative sampling is a relative term. It is clear from the participation levels, that to draw conclusions regarding within state or local dissemination of parasitism would be of limited value because participation at the state level was low; however, the goal of the survey was more holistic in nature and designed to garner information regionally and nationally. To this end, the results showed that even amidst the wide use of anthelmintics and newly developed treatments, infection trends and species dissemination had not changed since Porter (14) had first reported data in 1942.

Among the 1772 samples that were analyzed, 85.6% had one or more strongyle eggs, and 18.0% and 7.1% had *Nematodirus* spp. or *Trichuris* spp. eggs, respectively. This is not appreciably different from earlier observations (15) where 75.3% of calves and yearlings were positive for strongyle eggs in Iowa. The Iowa study also reported that 26.8% of the animals were passing *Nematodirus* spp. eggs. In 1956, a South Carolina study found that 40% of over 2000

fecal samples were positive for strongyle eggs and 0.7% and 1.2% for *Nematodirus* and *Trichuris*, respectively (16). In Illinois, Levine and Aves (17) found that all cattle surveyed were shedding strongyle eggs and that counts ranged between 43 to over 600 egg. A Georgia study found 83.9% of all beef cattle were positive for parasites by fecal examination (18). A more recent survey showed high prevalence (> 80%) of strongyle eggs in US beef and dairy operations (4).

The current study found that the average sample contained 32.5 strongyle egg across all regions surveyed. However, regional variation was observed, where the average egg were 30.3, 58.1, and 11.3 for the Central, South, and West regions, respectively. Interestingly, the lowest egg shedding was observed in the West and the highest in the South. This observation is consistent with the warm and moist, parasite friendly environment of the South.

Coccidia oocysts were found in 59.9% of all the fecal samples in this study with little variation between regions. The numbers presented here differ from those published from 2 Wisconsin studies (10,19) in which a prevalence of 84% compared to 58.0% was observed in the Central region defined by the current study. Samples in both studies were from animals of similar age. In a third study conducted in South Carolina (16), the prevalence was 21.5% compared with 63.1% in the region defined as South in the current study. The South Carolina samples were from primarily adult animals suggesting that acquired immunity may have played a factor in the lower prevalence values.

The prevalence of tapeworm eggs (*Moniezia*) was relatively low at 13.7% with few differences observed among regions. In the 1962 Wisconsin study (10), only 5.1% of cattle were found infected with tapeworms compared with 11.9% found in the Central region of the US in this study. These data suggest little has changed over the past 50 y.

Numerous surveys have identified the presence of internal parasite species based on necropsy findings. Porter (14) found an overall prevalence of 91% for *Cooperia punctata* in cattle in the southeastern US (Alabama, Florida, Georgia, Louisiana, Mississippi). Other nematode species identified were *Haemonchus contortus* (83%), *Ostertagia ostertagi* (74%), *Bunostomum* sp. (62%), *Oesophagostomum radiatum* (59%), *Trichostrongylus axei* (47%), *Cooperia pectinata* (32%), and *Strongyloides papillosus* (21%). These data are similar to our identifications based on PCR; *Cooperia* spp. (91.2%), *Ostertagia* sp. (79.4%), *Haemonchus* spp. (52.9%), and *Oesophagostomum* (38%). Likewise, the prevalence of *Nematodirus helvetianus* (15%) was similar to the 18.3% prevalence of *Nematodirus* sp. in the current study. The *Trichostrongylus* (3%) observation was considerably lower than the 47% prevalence observed by Porter (14).

Surprisingly there appears to have been little change in parasite prevalence (regardless of diagnostic technique) and numbers of eggs shed in the past 40 to 60 y, even with the extensive use of benzimidazoles and macrocyclic lactones, and less reliance on pasture management. It is interesting to postulate that in the early years of parasite control (40s, 50s, and 60s), parasitism was at a steady state as demonstrated by Porter (14). With few large-scale studies to detail prevalences in the intervening years, parasite numbers may have waned. A smaller more limited report supports that hypothesis

where a tracer calf study in Minnesota found *Ostertagia ostertagi* (37.5%), *Trichostrongylus* spp. (36.6%), *Cooperia* spp. (20.1%), and *Haemonchus* sp. (4.9%) (3). Now, as resistance continues to emerge and production systems have become more high intensity, we are once again approaching that “steady state” in dissemination of GI nematodes observed back in the 40s, 50s, and 60s. Presented another way, would a similar study, done several years after the introduction of the macrocyclic lactones and before the appearance of resistance have displayed a different level of parasitism? Perhaps a joint role for animal management and anthelmintic therapy must be considered rather than simply relying on the use of anthelmintics to effectively control parasitic nematodes.

Acknowledgments

This study was carried out as part of the USDA APHIS National Animal Health Monitoring System and, as such, the collection of data and samples was funded by the USDA. Supplemental funding was provided by the USDA, Agricultural Research Service to support the evaluation of the fecal samples.

References

1. Stromberg BE, Gasbarre LC. Gastrointestinal nematode control programs with an emphasis on cattle. *Vet Clin North Am Food Anim Pract* 2006;22:543–565.
2. Ballweber LR, Smith L, Stuedemann JA, Yazwinski T, Skogerboe TL. The effectiveness of a single treatment with doramectin or ivermectin in the control of gastrointestinal nematodes in grazing yearling stocker cattle. *Vet Parasitol* 1997;72:53–68.
3. Stromberg BE, Vatthauer RJ, Schlotthauer JC, et al. Production responses following strategic parasite control in a beef cow/calf herd. *Vet Parasitol* 1997;68:315–322.
4. Myers GH, Keith EA. Nationwide cattle survey: Zeroing in on parasites. *Large Anim Vet* 1993;48:30–32.
5. Bianchin I, Honer MR. Helminth parasites of beef cattle in the cerrado region of Brazil. *Trop Anim Health Prod* 1987;19:39–45.
6. Murphy TM, Fahy KN, McAuliffe A, Forbes AB, Clegg TA, O'Brien DJ. A study of helminth parasites in culled cows from Ireland. *Prev Vet Med* 2006;76:1–10.
7. Ploeger HW, Borgsteede FH, Sol J, et al. Cross-sectional serological survey on gastrointestinal and lung nematode infections in first and second-year replacement stock in the Netherlands: relation with management practices and use of anthelmintics. *Vet Parasitol* 2000;90:285–304.
8. USDA, 2010, Beef 2007–08, Part IV: Reference of Beef Cow-calf Management Practices in the United States, 2007–08. [Website on the Internet]. USDA:APHIS:VS, CEAH. Fort Collins, Colorado, USA #523.0210. Available from: http://www.aphis.usda.gov/animal_health/nahms/beefcowcalf/downloads/beef0708/Beef0708_dr_PartIV.pdf Last accessed June 26, 2015.
9. Gasbarre LC, Leighton EA, Bryant D. Reliability of a single fecal egg per gram determination as a measure of individual and herd values for trichostrongyle nematodes of cattle. *Am J Vet Res* 1996;57:168–171.
10. Cox DD, Todd AC. Survey of gastrointestinal parasitism in Wisconsin dairy cattle. *J Am Vet Med Assoc* 1962;141:706–709.
11. Bliss DH, Kvasnicka WG. The fecal examination: A missing link in food animal practice. *Compendium on continuing education for the practicing veterinarian* 1997;19(Suppl 4):S104–S109.
12. Herlich H. Attempts to produce protection against *Ostertagia ostertagi* in cattle. *Am J Vet Res* 1976;37:61–64.
13. Zarlenga DS, Chute MB, Gasbarre LC, Boyd PC. A multiplex PCR assay for differentiating economically important gastrointestinal nematodes of cattle. *Vet Parasitol* 2001;97:199–209.
14. Porter DA. Incidence of gastrointestinal nematodes of cattle in the Southeastern United States. *Am J Vet Res* 1942;3:304–308.
15. Zimmermann WJ, Hubbard ED. Gastrointestinal parasitism in Iowa cattle. *J Am Vet Med Assoc* 1961;139:555–559.
16. Hitchcock DJ. A survey of gastrointestinal parasites of cattle of South Carolina. *J Am Vet Med Assoc* 1956;129:34–35.
17. Levine ND, Aves IJ. The incidence of gastrointestinal nematodes in Illinois cattle. *J Am Vet Med Assoc* 1956;129:331–332.
18. Ciordia H. Occurrence of gastrointestinal parasites in Georgia cattle. *Am J Vet Res* 1975;36:457–461.
19. Hasche MR, Todd AC. Prevalence of bovine coccidia in Wisconsin. *J Am Vet Med Assoc* 1959;134:449–451.