

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

2019

Reducing Prairie Dog Populations and Damage by Castration of Dominant Males

Gary W. Witmer

USDA/APHIS National Wildlife Research Center

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc

 Part of the [Natural Resources and Conservation Commons](#), [Natural Resources Management and Policy Commons](#), [Other Environmental Sciences Commons](#), [Other Veterinary Medicine Commons](#), [Population Biology Commons](#), [Terrestrial and Aquatic Ecology Commons](#), [Veterinary Infectious Diseases Commons](#), [Veterinary Microbiology and Immunobiology Commons](#), [Veterinary Preventive Medicine, Epidemiology, and Public Health Commons](#), and the [Zoology Commons](#)

Witmer, Gary W., "Reducing Prairie Dog Populations and Damage by Castration of Dominant Males" (2019). *USDA National Wildlife Research Center - Staff Publications*. 2270.

https://digitalcommons.unl.edu/icwdm_usdanwrc/2270

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Reducing Prairie Dog Populations and Damage by Castration of Dominant Males

Gary W. Witmer, USDA/APHIS National Wildlife Research Center, Fort Collins, CO

ABSTRACT: Black-tailed prairie dogs (*Cynomys ludovicianus*) occur widely across the prairie states of North America. They compete with livestock for forage, transmit plague, and damage lawns, landscaping, and property. Interest in non-lethal methods, such as immunocontraception, is growing; however, reductions in the population due to contraception may be offset by increases in survival because adults and yearlings are not subject to the energetic demands of reproduction, and lower densities may increase the amount of resources available to growing offspring. Surgical sterilization provides a means for modeling these effects. Thus, we castrated males prior to the 1998 breeding season to simulate the potential effects of some contraceptives on body mass and survival. During the summer following treatment, the proportion of male and female adults/yearlings and juveniles captured did not differ between treatment and control colonies; however, the proportion of adults and yearlings captured decreased with later trapping periods. Hence we cannot recommend castration of dominant males to reduce colony expansion and damage by prairie dogs. Other methods of fertility control (GonaCon and diazacon) have shown more promise in prairie dogs.

KEY WORDS: *Cynomys ludovicianus*, fertility control, prairie dog, wildlife damage

Proceedings of the 18th Wildlife Damage Management Conference.
(J.B. Armstrong, G.R. Gallagher, Eds.). 2019. Pp. 28-31

INTRODUCTION

Prairie dogs (*Cynomys ludovicianus*) are a social rodent species of the grass prairies of the United States. They pose many challenges to resource managers in highly disturbed settings, such as urban/suburban areas, where conflicting interests persist regarding the presence of prairie dogs (Witmer et al. 2003, Zinn and Andelt 1999). The history, biology, ecology, and status of prairie dogs has been reviewed by Clippinger (1989), Fagerstone and Ramey (1996), Hoogland (2003), Mulhern and Knowles (1996), and U.S. Fish and Wildlife Service (2000). There is a need to better

monitor colonies and the changes that they undergo as well as a need to plan for future events. Municipalities have designed management plans to reduce conflicts by using public input, zoned management areas, and a variety of management techniques and tools. Individual populations must often be managed very differently.

Fertility control offers another potential solution to control expanding prairie dog colonies. The topic of wildlife fertility control was reviewed, including chemicals, delivery systems, advantages, disadvantages, regulatory issues, and challenges (Fagerstone et al. 2010). Previous

field studies (Nash et al. 2007) indicate that the steroid diazacholesterol can effectively limit prairie dog reproduction if delivered in adequate amounts to the animals over a sufficiently long period of time before the breeding season. Yoder and Miller (2011) showed that GonaCon can also inhibit reproduction in prairie dogs; however, this material must be injected. Physical castration is another possible means to control prairie dog populations. Prairie dogs live in colonies and those are comprised of extended family groups called coterie. Each coterie is comprised of one or more adult males (including a dominant male), several adult females, and their juveniles.

Our objective was to determine if physical castration of the adult males in coterie would reduce the reproduction of those coterie. We hypothesized that the castrated adult males would still prevent other males from impregnating adult females in their coterie. This study was conducted by an M.S. graduate student of ours at Colorado State University who never published the results of the study (Schwartz 2002). Hence, in this paper, I summarize the study; for more detail, see Schwartz 2002.

STUDY AREA AND METHODS

The field study was conducted on the Pine Ridge Natural Area of Fort Collins, Colorado. The 30-ha area is partially surrounded by residential housing. Prairie dogs were live-trapped, ear-tagged, and dyed with a marker so that each had a unique combination of letters and numbers. Observation blinds were built so that the prairie dogs could be observed without disturbing them. Through extensive observations, the individual coterie could be identified along with the adult (sexually active) males in each coterie. The reproductive status of captured adult females could also be determined. Fourteen coterie were used in the study. These were paired so

that each pair was of similar size in terms of the number of prairie dogs. In each pair, one was randomly selected to serve as the control while the other was the treatment coterie.

Once the adult males were identified, they were live-trapped and brought to an animal room of the USDA National Wildlife Research Center in Fort Collins for processing. The males from the treatment coterie were anesthetized and physically castrated by a veterinarian; the control coterie males were subjected to a sham procedure. Each male was returned to its original coterie shortly after recovery from the anesthetic.

We observed the dominant male in each coterie to make sure he continued to behave as the dominant male. The following spring, we observed the coterie to determine the reproductive output of each and that the original dominant male was still present. We also live-trapped the females to determine their reproductive status. We used logistic regression, ANOVA, and paired t-tests to analyze the data.

RESULTS AND DISCUSSION

There was no significant difference between the number of males per coterie in the treatment (mean = 2.9) versus the control (mean = 2.0) groups. The number of males with pigmented scrotum (i.e., sexually active) was not significantly different between treatment (14 of 20) and control (12 of 14) groups. All coterie contained at least one male with a pigmented scrotum.

There was no significant difference in the portion of females that lactated prior to treating the adult males in the treatment (26%) and control (39%) coterie. All females at the start of the study had sealed vulvas, meaning they had not yet been sexually active. There was no significant difference in the number of females in treatment (mean = 4.0) and control (5.0) groups.

After the breeding season, there was no significant difference between the number of lactating females in the treatment (19 of 24)

and control (27 of 28) groups. Litters first emerged in late April. The date that juveniles first emerged did not differ significantly between treatment and control groups. There were fewer juveniles born in the treatment groups (68; 32 male and 35 females) than in the control groups (139: 71 males and 67 females). However, it should be noted that 58 of the control juveniles were born on one large control coterie containing 12 females. There was no significant difference in the number of juveniles per coterie in the treatment (mean = 9.7) and the control (19.9) groups. Nor was there a significant difference in the number of pups born per female in the treatment (mean = 2.5) and control (4.0) groups.

It is clear that castrating the adult males in coterie did not slow reproduction in those coterie. It was noted that immigrant males (and in some cases females) moved between coterie. This may have led to copulation and pregnancies of adult females in some coterie. It has also been noted that female prairie dogs will often mate with more than one male (Hoogland 1995). This helps ensure pregnancy, but may also reduce the chances of infanticide of young by non-parent males.

This study indicates that, because of extra-coterie copulations, reproductive inhibitors that target adult males will not reducing population growth in black-tailed prairie dogs unless almost all adult males are treated. To be successful at reducing colony size and expansion, dual-gender contraceptive agents such as GonaCon and diazacon will likely be needed.

ACKNOWLEDGMENTS

This study was funded by the USDA National Wildlife Research Center (NWRC). This study was conducted under the NWRC IACUC-approved study protocol QA-543. I thank Aaron Schwartz and William Andelt of Colorado State University for their substantial efforts on this study. Mention of a company or

commercial product does not represent endorsement by the federal government.

LITERATURE CITED

- Clippinger, N. 1989. Habitat suitability index models: black-tailed prairie dog. Biological Report 82(10.156). U.S. Fish and Wildlife Service, Washington, D.C. 21 pp.
- Fagerstone, K., L. Miller, G. Killian, and C. Yoder. 2010. Review of issues concerning the use of reproductive inhibitors, with particular emphasis on resolving human-wildlife conflicts in North America. Integrative Zoology 1:15-30.
- Fagerstone, K., and C. Ramey. 1996. Rodents and lagomorphs. Pp. 83-132 In: Rangeland Wildlife. P. Krausman, editor. The Society for Range Management, Denver, Colorado.
- Hoogland, J. 1995. The black-tailed prairie dog: social life of a burrowing mammal. University of Chicago Press, Chicago, Illinois.
- Hoogland, J. 2003. Black-tailed prairie dog. Pp 232-247 In: Wild Mammals of North America. The Johns Hopkins University Press, Baltimore, Maryland.
- Mulhern, D., and C. Knowles. 1996. Black-tailed prairie dog status and future conservation planning. Pp. 19-29 In: Conserving biodiversity on native rangelands (D. Uresk, G. Schenbeck, and J. O'Rourke, eds.). USDA Forest Service General Technical Report RM-GTR-298. Fort Collins, Colorado. 38 pp.

- Nash, P., C. Furclow, K. Bynum, C. Yoder, L. Miller, and J. Johnston. 2007. 20,25-diazacholesterol as an oral contraceptive for black-tailed prairie dog population management. *Human-Wildlife Conflicts* 1:60-67.
- Schwartz, A. M. 2002. Surgical sterilization of the black-tailed prairie dog (*Cynomys ludovicianus*): a model for the effects of reproductive inhibition and hormonal reduction in the male. Thesis, Colorado State University, Fort Collins, Colorado.
- U.S. Fish and Wildlife Service. 2000. Endangered and threatened wildlife and plants: 12-month finding for a petition to list the black-tailed prairie dog as threatened. *Fed. Register* 65(24):5476-5488.
- Witmer, G., M. Brennan, D. Dees, et al. 2003. Black-tailed prairie dog management in urban-suburban settings: opportunities and challenges. *Trans. N. Amer. Wildl. and Natur. Resource Conf.* 68:209-221.
- Yoder, C., and L. Miller. 2011. Effect of GonaCon vaccine on black-tailed prairie dogs: immune response and health effects. *Vaccine* 29, 233-239.
- Zinn, H. C., and W. F. Andelt. 1999. Attitudes of Fort Collins, Colorado residents toward prairie dogs. *Wildlife Society Bulletin* 27:1098–1106.