

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2013

Antigenicity of Recombinant Maltose Binding Protein- Mycobacterium avium subsp. paratuberculosis Fusion Proteins with and without Factor Xa Cleaving

Ratna B. Gurung

University of Sydney, Camden & Ministry of Agriculture and Forests, Thimphu,
ratna.gurung@sydney.edu.au

Douglas J. Begg

University of Sydney

Auriol C. Purdie

University of Sydney

John Bannantine

USDA-ARS National Animal Disease Center, john.bannantine@usda.gov

Richard J. Whittington

University of Sydney

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Gurung, Ratna B.; Begg, Douglas J.; Purdie, Auriol C.; Bannantine, John; and Whittington, Richard J., "Antigenicity of Recombinant Maltose Binding Protein-Mycobacterium avium subsp. paratuberculosis Fusion Proteins with and without Factor Xa Cleaving" (2013). *Publications from USDA-ARS / UNL Faculty*. 2380.

<https://digitalcommons.unl.edu/usdaarsfacpub/2380>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Antigenicity of Recombinant Maltose Binding Protein-*Mycobacterium avium* subsp. *paratuberculosis* Fusion Proteins with and without Factor Xa Cleaving

Ratna B. Gurung,^{a,b} Douglas J. Begg,^a Auriol C. Purdie,^a John P. Bannantine,^c Richard J. Whittington^a

Faculty of Veterinary Science, University of Sydney, Camden, NSW, Australia^a; Department of Livestock, Ministry of Agriculture and Forests, Thimphu, Bhutan^b; Bacterial Diseases of Livestock Research Unit, USDA-ARS National Animal Disease Center, Ames, Iowa, USA^c

***Mycobacterium avium* subsp. *paratuberculosis* causes Johne's disease (JD) in ruminants. Proteomic studies have shown that *M. avium* subsp. *paratuberculosis* expresses certain proteins when exposed to *in vitro* physiological stress conditions similar to the conditions experienced within a host during natural infection. Such proteins are hypothesized to be expressed *in vivo*, are recognized by the host immune system, and may be of potential use in the diagnosis of JD. In this study, 50 recombinant maltose binding protein (MBP)-*M. avium* subsp. *paratuberculosis* fusion proteins were evaluated using serum samples from sheep infected with *M. avium* subsp. *paratuberculosis*, and 29 (58%) were found to be antigenic. Among 50 fusion proteins, 10 were evaluated in MBP fusion and factor Xa-cleaved forms. A total of 31 proteins (62%) were found to be antigenic in either MBP fusion or factor Xa-cleaved forms. Antigenicity after cleavage and removal of the MBP tag was marginally enhanced.**

Johne's disease (JD) in ruminants is a chronic infection of the intestines caused by *Mycobacterium avium* subsp. *paratuberculosis*. Economic losses arise due to culling, reduced production of milk and wool, and mortalities (1, 2). The disease is characterized by a long incubation period, and subclinical infection creates a potential source of infection for uninfected animals. The most common method of detection of JD is by measuring immune responses in the infected host. The two widely used assays to measure cell-mediated and antibody-mediated immune responses are the gamma interferon (IFN- γ) assay and the enzyme-linked immunosorbent assay (ELISA), respectively. These depend on *M. avium* subsp. *paratuberculosis*-specific antigens for stimulating IFN- γ from memory T cells or detecting specific antibodies in blood samples. Currently used antigens in these assays are French-pressed proteins or purified protein derivatives (PPDs) derived from whole cells of *M. avium* subsp. *paratuberculosis*, containing a large number of antigens. The diagnostic specificity of commercial antibody ELISAs is generally high, but sensitivity is poor (3, 4). The accuracy of an ELISA also can be adversely affected by potential cross-reactions due to exposure of the host to environmental mycobacteria. The specificity of an assay can be enhanced by absorbing the serum against *Mycobacterium phlei* proteins (5, 6). However, it is difficult to enhance ELISA sensitivity due to the poor Th2 responses during the lengthy latent period of the disease and the large population of infected animals with latent subclinical infections in an exposed flock or herd. Therefore, to enhance assay performance, new *M. avium* subsp. *paratuberculosis*-specific antigens that are expressed during latency need to be identified.

Evidence of dormancy in *M. avium* subsp. *paratuberculosis* when the organism was present in the soil/pasture environment was reported (7). *In vitro* studies simulating the stress conditions of natural infection reported dormancy-associated proteins in *M. avium* subsp. *paratuberculosis* (8–10). These findings led to a hypothesis that *M. avium* subsp. *paratuberculosis* expresses stress/dormancy-related proteins during infection of the host. The use of *M. avium* subsp. *paratuberculosis* proteins that are expressed *in*

in vivo following pathogen entry into the host as diagnostic antigens may be of value in the detection of an early stage of *M. avium* subsp. *paratuberculosis* infection. Indeed, some of the *M. avium* subsp. *paratuberculosis* proteins known to be differentially regulated under stress conditions were found to be antigenic in serum collected from sheep infected with *M. avium* subsp. *paratuberculosis*, and these findings support the hypothesis that stress proteins expressed *in vitro* are also expressed *in vivo* (11–13).

A large number of recombinant *M. avium* subsp. *paratuberculosis* antigens have been investigated for their diagnostic potential in cell- and antibody-mediated assays (11, 13–22). Some of these *M. avium* subsp. *paratuberculosis* proteins were from groups of proteins that were differentially regulated under physiological stress conditions. Although many proteins were found to be antigenic, no obvious candidate has yet been identified as having suitable diagnostic sensitivity and specificity.

A major limitation for characterization of recombinant *M. avium* subsp. *paratuberculosis* proteins is their expression as inclusion bodies or insoluble proteins, especially when prepared using histidine (His) as an affinity purification tag (13). Production of antigens from insoluble proteins involves processes that may be detrimental to biological activity. Expression of maltose binding protein (MBP) fusion proteins facilitates maintenance of the solubility, structure, and functions of recombinant proteins through downstream processing (23, 24). Several recombinant MBP-*M. avium* subsp. *paratuberculosis* fusion proteins (MBP fusion proteins) were found to be antigenic in sheep, cattle, and mice infected with *M. avium* subsp. *paratuberculosis* (15, 16, 25). How-

Received 16 September 2013 Accepted 26 September 2013

Published ahead of print 16 October 2013

Address correspondence to Ratna B. Gurung, ratna.gurung@sydney.edu.au.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/CVI.00596-13

U.S. government works are not subject to copyright.

TABLE 1 Evaluation and comparison of recombinant *M. avium* subsp. *paratuberculosis* proteins (MBP fusion and His-tagged proteins)

Protein type and function	Protein(s)
MBP fusion proteins	
Phosphate metabolism	MAP0435c
Cold shock protein	MAP0810
Putative virulence factor	MAP1272c
Universal stress proteins	MAP1339
Cell wall synthesis	MAP2058c
Signal recognition	MAP3968
ATP and purine biosynthesis	MAP2450c, MAP3393c
Cell division	MAP0068, MAP1889c
Heat shock protein chaperone	MAP3268, MAP3701c
Response regulators	MAP0834c, MAP3200
Protein synthesis	MAP1027c, MAP4125
Proteolysis	MAP1834c, MAP2280c, MAP2281c
Amino acid metabolism	MAP1293, MAP1297, MAP1846c, MAP2864c
Antioxidant enzymes	MAP1588c, MAP1589c, MAP1653, MAP4340
Hypothetical protein	MAP0593c, MAP0184c, MAP1586, MAP3555, MAP3864
Fatty acid metabolism	MAP0508, MAP0516c, MAP1017c, MAP2698c, MAP2872c, MAP3190, MAP3577, MAP3651c
Cellular processes	MAP0187c, MAP0540, MAP1560, MAP1885c, MAP2411, MAP2487c, MAP2705c, MAP3007, MAP3538, MAP3567
His-tagged proteins	
Putative virulence factor	MAP1272c
Fatty acid metabolism	MAP2698c
Cellular processes	MAP2487c, MAP3567

ever, MBP alone, with a molecular mass of about 42.5 kDa, is known to have a small amount of seroreactivity; because of this, it must be used as a control in ELISAs (23). Furthermore, it is not known if the MBP protein masks the immune recognition of a protein of interest. Therefore, cleavage of the MBP tag from the recombinant *M. avium* subsp. *paratuberculosis* proteins may be beneficial for their use.

Factor Xa is a protease that specifically cleaves after the arginine residue in its preferred site Ile-(Glu or Asp)-Gly-Arg sequence (26) and can be used to separate the MBP affinity purification tag from the protein of interest following expression and purification (27, 28). This protease was used in the current study for removal of the MBP purification tag. The aims of this study were to evaluate the antigenicity of *M. avium* subsp. *paratuberculosis* recombinant proteins hypothesized to be upregulated under stress conditions and to investigate their potential use in early diagnosis. These proteins were examined with and without the MBP tag to determine if the proteins cleaved of MBP had better antigenicity, and some were compared with the corresponding His-tagged recombinant protein.

MATERIALS AND METHODS

Antigens. The *M. avium* subsp. *paratuberculosis* proteins in this study (Table 1) were selected based on their expression in response to *in vitro* physiological stress conditions (8–10). Fifty *M. avium* subsp. *paratuber-*

culosis recombinant proteins used in this study were produced as MBP fusion proteins at the Bacterial Diseases of Livestock Research Unit, USDA-ARS Agricultural Research Service National Animal Disease Center (NADC) (Ames, IA), and one (MAP1272c) was produced as a His-tagged recombinant *M. avium* subsp. *paratuberculosis* protein (29). The MBP fusion proteins were produced as described previously (30). Briefly, the full-length coding sequence of the *M. avium* subsp. *paratuberculosis* protein was amplified using gene-specific primers and was cloned into the pMAL-c2 translational fusion expression vector. The vector and amplified products were digested with XbaI and HindIII, and the ligated products were transformed into *Escherichia coli* DH5 α cells. The overexpressed proteins were extracted and purified by affinity chromatography with amylose resin columns (New England BioLabs). Three His-tagged recombinant *M. avium* subsp. *paratuberculosis* proteins (MAP2698c, MAP2487c, and MAP3567) were produced at the Faculty of Veterinary Science, University of Sydney (Sydney, Australia), as previously described (12). Briefly, gene-specific primers were designed to include *attB1* and *attB2* sites at the 5' end of each sequence. The complete open reading frames of each gene were amplified by PCR using Gateway technology (Invitrogen, Australia). Amplified and purified PCR products were cloned into the donor vector pDONR221 (Invitrogen, Australia) and transformed into One Shot TOP10 chemically competent *E. coli* cells (Invitrogen, Australia) to produce an entry clone. Purified entry clones were subcloned into the destination vector pET160-DEST with an N-terminal 6 \times His and Lumio tag (Champion pET160 Gateway expression kit with Lumio technology; Invitrogen, Australia) and transformed into One Shot TOP10 chemically competent *E. coli* cells to produce an expression clone. One Shot BL21 Star (DE3) cells (Invitrogen, Australia) were transformed with the purified expression clone. The transformed culture was induced with 1 mM isopropyl- β -1-thiogalactopyranoside to express the recombinant proteins. Recombinant proteins were extracted and purified by affinity liquid chromatography (AKTApurifier system; GE Healthcare). The four His-tagged recombinant *M. avium* subsp. *paratuberculosis* proteins (1 from the NADC and 3 from the University of Sydney) were compared with the corresponding MBP fusion proteins.

Proteolytic cleavage of MBP fusion proteins. A pilot study was performed with MBP fusion proteins (MAP0435c, MAP1846c, and MAP1017c) to determine the optimal time for enzymatic cleavage of the *M. avium* subsp. *paratuberculosis* protein from the MBP tag. Factor Xa (Amersham Biosciences) was reconstituted to a final concentration of 1 unit/ μ l in nuclease-free water at 4°C. MBP fusion proteins were diluted to 1 mg/ml in phosphate-buffered saline (PBS). Factor Xa (1 μ l) was added to the MBP fusion protein (100 μ l) in a 1.5-ml screw-cap tube and mixed briefly with a vortex mixer. Reaction buffer (100 μ l) was added to the mixture, which was vortex mixed briefly and incubated for 16 to 40 h at room temperature (RT) (22 to 24°C). Fifteen-microliter aliquots were collected after 16, 18, 20, 25, 30, 35, and 40 h of incubation for SDS-PAGE analysis. The optimal cleavage time was determined for each protein based on the appearance of two bands that corresponded to the expected molecular masses of MBP and the relevant *M. avium* subsp. *paratuberculosis* protein.

Due to the limited volume (500 μ l) of MBP fusion proteins, only 22 proteins with volumes and concentrations adequate for cleavage and antigenicity evaluation were available. A reaction mixture containing 1 ml of MBP fusion protein (1 mg), 10 μ l (10 units) of factor Xa, and 1 ml of reaction buffer in a 5-ml screw-cap tube was prepared, vortex mixed briefly, and incubated for up to 40 h at RT. Aliquots of 15 μ l were collected at various incubation times (16, 18, 20, 25, 30, and 40 h) and examined by SDS-PAGE.

SDS-PAGE analysis. SDS-PAGE analyses of proteins were performed using 12% precast polyacrylamide gels (Mini-PROTEAN TGX precast gel, product no. 456-1043; Bio-Rad). Briefly, a 15- μ l aliquot of each protein sample (MBP-LacZ, MBP-*M. avium* subsp. *paratuberculosis* fusion protein, factor Xa-cleaved *M. avium* subsp. *paratuberculosis* protein, pu-




FIG 1 SDS-PAGE analysis of pilot scale factor Xa cleavage of MBP fusion proteins. (A) MBP-MAP0435c; (B) MBP-MAP1017c; (C) MBP-MAP1846. Lanes 1, molecular mass markers (PageRuler Plus prestained protein ladder); lanes 2, MBP-LacZ (54 kDa); lanes 3, MBP-*M. avium* subsp. *paratuberculosis* fusion protein before cleavage; lanes 4 to 10, MBP-*M. avium* subsp. *paratuberculosis* fusion proteins cleaved at different time points, i.e., 16 h (lanes 4), 18 h (lanes 5), 20 h (lanes 6), 25 h (lanes 7), 30 h (lanes 8), 35 h (lanes 9), and 40 h (lanes 10).

rified MBP, or purified *M. avium* subsp. *paratuberculosis* protein) was placed in a 1.5-ml screw-cap tube. The protein samples were mixed with 3 μ l of reducing sample buffer and heated for 5 min in a boiling water bath. Protein samples (18 μ l) were loaded onto a precast gel, and electrophoresis was performed at a constant voltage of 180 V (SmartPower 4000 power pack) for 65 min or until the visible line of bromophenol blue reached the bottom of the gel, using a Mini-PROTEAN 3 cell system (Bio-Rad). Protein bands were stained with Coomassie brilliant blue (0.1% Coomassie brilliant blue G-250, 3% *ortho*-phosphoric acid, 10% ammonium sulfate, and 20% methanol). The protein bands were visualized with a Geldoc system (Bio-Rad).

Affinity chromatography purification of cleaved proteins. Purification of cleaved MBP fusion proteins was performed with an AKTApurifier fast performance liquid chromatography (FPLC) system (GE Healthcare) using a MBPTrap HP dextran-Sepharose high performance column (5 ml; GE Healthcare), following the manufacturer's manual. Briefly, the column was equilibrated with 5 column volumes (CVs) of binding buffer at a flow rate of 2 ml/min. A cleaved protein sample solution (1 ml) was applied using a 1-ml sample loop, followed by washing with 5 CVs of binding buffer to remove unbound protein (*M. avium* subsp. *paratuberculosis* protein), at a flow rate of 5 ml/min. The flowthrough protein peaks

were collected in 2-ml fractions. The bound MBP was eluted with 5 CVs of elution buffer at a flow rate of 5 ml/min, and protein peaks were collected in 2-ml fractions. The column was reequilibrated with 5 CVs of binding buffer at a flow rate of 5 ml/min. The cleaved and purified *M. avium* subsp. *paratuberculosis* proteins were dialyzed against PBS overnight at 4°C using a 12-kDa-cutoff membrane tube. Proteins were concentrated by centrifuging the dialyzed protein solution at $2,000 \times g$ for 10 min at 4°C (AllegraX-12R; Beckman Coulter), using Amicon Ultra-15 centrifugal filter units (nominal molecular mass limit of 10 kDa). The protein yield in the retentate was estimated using a spectrophotometer (NanoDrop 1000; Thermo Scientific) set at 280 nm.

Serum samples. A total of 46 sheep serum samples were obtained from the serum archive maintained at the Faculty of Veterinary Science, University of Sydney. Twenty-three serum samples were from unexposed/uninfected sheep from Western Australia (31) that were certified to be free of JD based on the negative test results for their flocks. Another 23 serum samples were obtained from exposed/infected sheep that either tested positive by tissue/fecal culture or had histopathological lesions consistent with ovine JD. Serum samples obtained from infected animals also tested positive with the Institut Pourquier ELISA (32), with a sample-to-positive ratio of >70%, and were categorized as strong reactors ($n = 8$), medium

reactors ($n = 8$), or low reactors ($n = 7$). The infected sheep were fecal culture positive (shedder; $n = 15$) or negative (nonshedder; $n = 8$). Histopathological lesions were categorized according to a previously described method (33), as follows: no or low-grade lesion (grade 0 to 2; $n = 4$), paucibacillary lesion (grade 3a plus grade 3c; $n = 2 + 8$), or multibacillary lesion (grade 3b; $n = 9$).

Positive- and negative-control sheep sera were obtained from sheep from the University of Sydney that were in an independent experimental infection trial (34). The negative-control serum was obtained from a sheep not exposed to *M. avium* subsp. *paratuberculosis* that had tested negative by fecal culture, biopsied tissue culture, histopathological analysis, and ELISA. Positive-control serum was obtained from a Gudair vaccine-immunized sheep, and the serum was determined to be positive by the Institut Pourquier ELISA.

ELISA methodology. Antibody ELISA was performed to evaluate the seroreactivity of MBP fusion proteins, cleaved *M. avium* subsp. *paratuberculosis* proteins, and His-tagged recombinant *M. avium* subsp. *paratuberculosis* proteins, using serum samples from *M. avium* subsp. *paratuberculosis*-infected and uninfected sheep. Briefly, 50 μ l/well of antigen at the specified final concentrations of MBP fusion antigens (see Tables 3 and 4), cleaved *M. avium* subsp. *paratuberculosis* antigens (5 μ g/ml) (see Table 4), or His-tagged *M. avium* subsp. *paratuberculosis* antigens (see Table 5) in carbonate buffer (0.1 M carbonate buffer [pH 9.6]) were coated on 96-well flat-bottom microplates (Nunc MaxiSorp; Nunc) and incubated overnight at 4°C. The plates were machine washed 5 times with purified reverse-osmosis (RO) water with Tween 20. ELISA plates coated with MBP fusion proteins and MBP-LacZ were blocked with 4% skim milk, and plates coated with cleaved *M. avium* subsp. *paratuberculosis* proteins and His-tagged *M. avium* subsp. *paratuberculosis* proteins were blocked with 1% (vol/vol) fetal calf serum (FCS) for 30 min each.

The serum samples were diluted (1:100) in a diluent (0.1% [vol/vol] FCS in PBS with Tween 20 [PBST]) containing 1.3 mg/ml heat-killed *M. phlei* protein (Elizabeth Macarthur Agricultural Institute, New South Wales, Australia) and were absorbed overnight at 4°C, with constant end-to-end shaking. The absorbed serum was centrifuged at 2,500 \times g for 10 min at RT to separate the supernatant from the particulate *M. phlei*. The absorbed serum supernatant (50 μ l) was added to the required wells and incubated for 1 h at RT. The plate was machine washed 5 times with RO water with Tween 20 prior to the addition of horseradish peroxidase-labeled mouse anti-sheep IgG monoclonal antibody conjugate (50 μ l, 1:40,000, clone GT-34; Sigma, New South Wales, Australia) in diluent (0.1% [vol/vol] FCS in PBST) and then incubated for 1 h at RT. The plate was machine washed as described above, and 100 μ l of 3,3',5',5'-tetramethylbenzidine (TMB) substrate was added. The plate was incubated for 20 min in the dark, after which the chromogenic reaction was stopped with the addition of stop solution (50 μ l of 2 M sulfuric acid). The optical density at 450 nm (OD_{450}) was measured using a plate reader (Multiskan Ascent; Thermo Scientific, Victoria, Australia). Serum samples were also tested with MBP-LacZ (5 μ g/ml) to examine seroreactivity to MBP. The ELISA results are presented as mean OD_{450} values.

Data analysis. Statistical analysis was performed using GenStat 12.1 (VSN International Ltd., United Kingdom) and GraphPad Prism 4.0 (GraphPad Software Inc., La Jolla, CA) software. The seroreactivities of MBP fusion antigens, factor Xa-cleaved *M. avium* subsp. *paratuberculosis* antigens, and His-tagged *M. avium* subsp. *paratuberculosis* antigens between groups were compared by analysis of variance (ANOVA) with a Bonferroni correction for multiple comparisons, as described previously (35). The area under the receiver operating characteristic curve (AUC_{ROC}) was calculated for the ability of an assay using each antigen to discriminate between uninfected and *M. avium* subsp. *paratuberculosis*-infected sheep. An assay with an AUC_{ROC} value of 1.0 is considered to be perfect, and one with a value of 0.5 is considered worthless (36).

TABLE 2 Expected molecular masses of MBP fusion and cleaved proteins

Cluster	Protein	CT ^a (h)	Molecular mass (kDa) of:	
			Fusion protein ^b	<i>M. avium</i> subsp. <i>paratuberculosis</i> protein
Amino acid metabolism	MAP1297	40	67.8	25.3
	MAP1846c	40	73	30.5
Antioxidant enzymes	MAP1588c	20	61.3	18.8
	MAP1589c	16	64.1	21.6
	MAP1653	16	58.9	16.4
ATP biosynthesis	MAP3393c	20	60	17.5
Cell division	MAP0068	40	60	17.5
Cellular processes	MAP1560	16	57.7	15.2
	MAP1885c	16	60.9	18.4
	MAP2411	16	58	15.5
	MAP2705c	20	56.4	13.9
Fatty acid metabolism	MAP3538	16	58.4	15.9
	MAP1017c	16	70.3	27.8
	MAP2698c	20	73.9	31.4
Heat shock protein	MAP3190	16	75.8	33.3
	MAP3268	16	58.9	16.4
Hypothetical protein	MAP3701c	40	58.7	16.2
	MAP0593c	40	57.3	14.8
Phosphate metabolism	MAP3555	16	61.3	18.8
	MAP0435c	16	61.1	18.6
Proteolysis	MAP2280c	20	65.7	23.2
Universal stress proteins	MAP1339	20	57.9	15.4

^a CT, cleavage time.

^b Fusion protein mass is the total mass of MBP (42.5 kDa) and the *M. avium* subsp. *paratuberculosis* protein.

RESULTS

Cleavage of MBP fusion proteins on a pilot scale. Three MBP fusion proteins were cleaved by factor Xa protease in a pilot study and analyzed by SDS-PAGE (Fig. 1A to C). Factor Xa was able to cleave MBP-MAP0435c and MBP-MAP1017c after 16 h of incubation at RT. Factor Xa was able to cleave MBP-MAP1846 completely only after 30 h of incubation. Cleavage of all three fusion proteins revealed the expected band sizes for MBP (42.5 kDa) and *M. avium* subsp. *paratuberculosis* proteins. However, multiple bands were observed for the proteins MAP0435c and MAP1017c. The pilot study showed that different proteins required different cleavage times.

Cleavage of MBP fusion proteins on a large scale. Twenty-two MBP fusion proteins were cleaved by factor Xa at different incubation times. Eleven proteins were completely cleaved with 16 h of incubation, six proteins required 20 h, and five proteins required 40 h for complete cleavage (Table 2). The SDS-PAGE analyses of cleaved products are shown in Fig. 2A to F. For each MBP fusion protein, the MBP tag was separated from the *M. avium* subsp. *paratuberculosis* protein of expected molecular mass. Four pro-




FIG 2 SDS-PAGE of factor Xa-cleaved *M. avium* subsp. *paratuberculosis* proteins. Lanes 1, molecular mass markers (PageRuler Plus prestained protein ladder); lanes 2, MBP-LacZ; lanes 3, 5, 7, and 9, MBP-*M. avium* subsp. *paratuberculosis* fusion proteins; lanes 4, 6, 8, and 10, factor Xa-cleaved MBP and *M. avium* subsp. *paratuberculosis* proteins.

teins, i.e., MBP-MAP0593c, MBP-MAP3701c, MBP-MAP1297, and MBP-MAP0068c, showed weak bands of undigested fusion proteins even with 40 h of incubation. Fusion proteins such as MBP-MAP0435c, MBP-MAP3393c, MBP-MAP3268, MBP-MAP1653, MBP-MAP1846, MBP-MAP2280c, MBP-MAP1885c, and MBP-MAP0068c were found to be cleaved, but there were multiple bands of *M. avium* subsp. *paratuberculosis* proteins.

SDS-PAGE analysis of cleaved and purified proteins. Due to the limited volumes of MBP fusion proteins received from the USDA Agricultural Research Service (500 μ l each), only 22 proteins were used for factor Xa cleavage experiments. Of 22 cleaved proteins, only 10 proteins had adequate volumes for affinity liquid chromatography purification.

The flowthrough (cleaved *M. avium* subsp. *paratuberculosis* proteins) and eluate (cleaved MBP tag) fractions were verified by SDS-PAGE analysis (Fig. 3A to E). The expected molecular masses are shown in Table 2 and Fig. 3A to E. All of the cleaved *M. avium* subsp. *paratuberculosis* proteins were obtained with >95% purity, based on the band appearance on SDS-PAGE analysis. The yields of purified and concentrated *M. avium* subsp. *paratuberculosis* proteins were approximately 20 μ g/ml (MAP2698c and MAP3555), 30 μ g/ml (MAP1846, MAP1885c, and MAP2280c), 40 μ g/ml

(MAP1017c and MAP0593c), 60 μ g/ml (MAP3538), or 70 μ g/ml (MAP1588c and MAP3190). Separated bands for cleaved *M. avium* subsp. *paratuberculosis* proteins MAP1846, MAP1885c, and MAP3555 were faint, due to the low protein concentrations.

Antigenicity of MBP fusion proteins. Fifty MBP fusion proteins were evaluated, and 29 (58%) were found to be detected by antibodies in sera obtained from *M. avium* subsp. *paratuberculosis*-infected sheep (Tables 3 and 4). The remaining 42% (21/50) of the proteins were not able to differentiate infected from uninfected animals. The seroreactivity of the control MBP-LacZ was not significantly different in the infected and uninfected groups ($P > 0.05$). The greatest ability to differentiate the infected group from the uninfected group was observed for the MAP0516c protein, which is encoded by *echA20* and is involved in fatty acid metabolism ($AUC_{ROC} = 0.758$, $P = 0.001$), MAP2872c, which is encoded by *fabG5* and is a 3-ketoacylreductase ($AUC_{ROC} = 0.765$, $P = 0.001$), and MAP1834c, which is encoded by *prcA* and is a proteome subunit protein ($AUC_{ROC} = 0.765$, $P = 0.002$).

Among the MBP fusion proteins evaluated, 19/50 (38%) were able to produce significantly higher OD values in serum samples obtained from sheep with paucibacillary infections, compared with sera from unexposed sheep ($P < 0.05$). Simi-




FIG 3 SDS-PAGE of factor Xa-cleaved and purified MBP fusion proteins. Lanes 1, molecular mass markers (PageRuler Plus prestained protein ladder) (M); lanes 2, MBP-LacZ; lanes 3 to 10, proteins shown as MBP fusion protein and factor Xa-cleaved protein, eluate, and flowthrough fraction.

larly, 26/50 proteins (52%) were able to produce significantly higher OD values in serum samples that were low reactors in the Institut Pourquier ELISA, in comparison with sera from unexposed sheep ($P < 0.05$) (Tables 3 and 4). However, none of the proteins was able to detect infections that were associated with no or low-grade lesions.

Antigenicity of cleaved and MBP fusion proteins. Successfully cleaved and purified *M. avium* subsp. *paratuberculosis* proteins were evaluated and ELISA results were compared with those for fused forms (Table 4). The OD values of infected sheep sera were significantly higher than those of the uninfected sheep sera for all factor Xa-cleaved proteins except MAP1885c. The MBP fusion proteins MAP0593c and MAP2698c were not able to differentiate between infected and uninfected sheep, but when they were evaluated as cleaved proteins purified from the MBP fusion proteins, the OD values of the infected group were significantly higher ($P < 0.05$) than those of the uninfected group. Protein MAP1885c was not able to differentiate between the infected and uninfected groups in either the fused or cleaved forms. Based on AUC_{ROC} values, factor Xa cleavage did not enhance the immunoreactivity of four proteins (MAP1846, MAP1588c, MAP1885c, and MAP3555). The *M. avium* subsp. *paratuberculosis*-specific immunoreactivity of two proteins (MAP2698c and MAP0593c) was enhanced while that of three proteins (MAP3538, MAP1017c, and MAP2280c) was marginally enhanced by cleavage of MBP. In

general, the ability of cleaved proteins to differentiate the infected and uninfected groups was enhanced in only 50% of the proteins (5/10 proteins).

Antigenicity evaluation of His-tagged recombinant *M. avium* subsp. *paratuberculosis* proteins. All four His-tagged recombinant *M. avium* subsp. *paratuberculosis* proteins were able to differentiate between the infected and uninfected sheep sera ($P < 0.05$) (Table 5). The order of most antigenic to least antigenic proteins was MAP1272c > MAP3567 > MAP2487c > MAP2698c, with AUC_{ROC} values of 0.90, 0.75, 0.70, and 0.69, respectively.

The seroreactivity of two different protein expression systems (His tag and MBP fusion) were compared for MAP2698c, MAP1272c, MAP2487c, and MAP3567. All four proteins were able to differentiate between the infected and uninfected groups of sheep in both expression systems. The OD values obtained from an infected group of sheep using His-tagged recombinant proteins were significantly higher ($P < 0.05$) than those obtained using the corresponding MBP fusion proteins (Table 6).

The ELISA results obtained from the infected sheep using MAP2698c protein as a MBP fusion, factor Xa-cleaved, or His-tagged protein were analyzed by one-way ANOVA with a Bonferroni correction for multiple comparisons. The OD values obtained using factor Xa-cleaved *M. avium* subsp. *paratuberculosis* proteins and His-tagged *M. avium* subsp. *paratuberculosis* proteins were significantly higher than those obtained with MBP fu-

TABLE 3 Antigenicity evaluation of MBP fusion proteins

Protein	Protein concn (µg/ml)	OD ₄₅₀ (n) for:		P ^c	AUC _{ROC}
		Unexposed samples (23)	Exposed samples (23)		
MAP0068 ^a	5	0.133	0.218	0.002	0.71
MAP0184c ^{a,b}	10	0.215	0.355	0.002	0.745
MAP0187c	5	0.236	0.31	0.118	
MAP0435c	20	0.278	0.291	0.797	
MAP0508 ^{a,b}	6	0.3	0.501	0.004	0.713
MAP0516c ^{a,b}	5	0.257	0.439	<0.001	0.758
MAP0540	7	0.298	0.36	0.238	
MAP0810	10	0.187	0.241	0.111	
MAP0834c ^{a,b}	5	0.141	0.216	0.002	0.742
MAP1027c	10	0.176	0.234	0.083	
MAP1272c ^{a,b}	10	0.26	0.451	0.005	0.703
MAP1293 ^{a,b}	5	0.186	0.292	0.012	0.717
MAP1297 ^{a,b}	10	0.172	0.321	0.002	0.744
MAP1339	5	0.204	0.274	0.067	
MAP1560	10	0.252	0.273	0.51	
MAP1586	5	0.285	0.311	0.515	
MAP1589c	10	0.183	0.269	0.02	0.677
MAP1653 ^b	10	0.195	0.263	0.045	0.683
MAP1834c ^{a,b}	9	0.262	0.44	0.002	0.765
MAP1889c	5	0.215	0.203	0.652	
MAP2058c ^{a,b}	6	0.265	0.442	0.005	0.703
MAP2281c	7	0.29	0.281	0.773	
MAP2411	10	0.265	0.298	0.481	
MAP2450c ^{a,b}	8	0.202	0.293	0.045	0.677
MAP2487c	8	0.307	0.331	0.576	
MAP2705c	10	0.208	0.228	0.541	
MAP2864c ^{a,b}	8	0.336	0.455	0.035	0.658
MAP2872c ^{a,b}	8	0.166	0.289	<0.001	0.765
MAP3007 ^{a,b}	7	0.211	0.357	0.003	0.74
MAP3200 ^b	10	0.216	0.339	0.008	0.699
MAP3268	10	0.272	0.229	0.386	
MAP3393c	10	0.261	0.291	0.497	
MAP3567 ^{a,b}	1	0.137	0.234	0.002	0.728
MAP3577	7	0.194	0.215	0.465	
MAP3651c ^{a,b}	5	0.208	0.323	0.01	0.695
MAP3701c	10	0.166	0.198	0.274	
MAP3864 ^b	10	0.208	0.289	0.048	0.642
MAP3968 ^{a,b}	7	0.154	0.277	0.002	0.719
MAP4125	10	0.284	0.292	0.847	
MAP4340 ^b	10	0.194	0.296	0.006	0.732

^a Proteins able to produce significantly higher OD values for serum samples obtained from sheep with paucibacillary infections than for serum samples from unexposed sheep.

^b Proteins able to produce significantly higher OD values for serum samples that were low reactors in the Institut Pourquier ELISA than for serum samples from unexposed sheep.

^c Mean OD values for the exposed group are significantly higher than those for the unexposed group ($P < 0.05$).

sion proteins ($P < 0.001$). The OD values obtained with factor Xa-cleaved *M. avium* subsp. *paratuberculosis* proteins and His-tagged *M. avium* subsp. *paratuberculosis* proteins were similar ($P > 0.05$).

DISCUSSION

This study investigated the antigenicity of *M. avium* subsp. *paratuberculosis* proteins in either MBP-fused, factor Xa-cleaved, or His-tagged forms, using sera that were known to have *M. avium*

subsp. *paratuberculosis*-specific antibodies, as detected by the commercial Institut Pourquier ELISA (32). The proportion of antigenic proteins increased from 58% to 62% after factor Xa cleavage. Sufficient amounts of MBP fusion proteins were available for 22 proteins to be cleaved using factor Xa. Ten of 22 proteins were successfully purified and evaluated in both MBP fusion and factor Xa-cleaved forms. Four proteins were evaluated in the His-tagged form to compare the expression systems.

Factor Xa cleaves proteins after the arginine residue in its preferred cleavage site Ile-(Glu or Asp)-Gly-Arg in the fusion link between MBP and the target protein. However, reports suggest that it quite often cleaves at secondary sites, depending on the conformation of the protein substrate (37, 38). The most common secondary site is Gly-Arg, usually in a partially unfolded protein (39). The probable cause of the appearance of multiple bands on SDS-PAGE gels in this study may be proteolysis or cleavage of target proteins at such secondary sites. Furthermore, on screening of the amino acid sequences of proteins that showed multiple bands on SDS-PAGE gels, several such sites were identified in proteins MAP0435c, MAP1846, and MAP1017c. MAP0435c had two Gly-Arg sites (amino acid positions 24-25 and 142-143), MAP1846 had three Gly-Arg sites (amino acid positions 68-69, 99-100, and 159-160), and MAP1017c had one Gly-Arg site (amino acid positions 168-169).

The protein cleavage experiment showed that 1 unit of factor Xa was sufficient to cleave 100 µg of MBP fusion protein after as little as 18 h of incubation at RT (22 to 23°C), and this finding was consistent with the findings from another study (40). However, factor Xa cleavage was not complete even after 40 h of incubation for proteins MBP-MAP0068c, MBP-MAP0593c, MBP-MAP1297, MBP-MAP1846c, and MBP-MAP3701c. The probable causes of incomplete cleavage may be an inaccessible factor Xa recognition site or alteration of the factor Xa recognition site during cloning (41). This process of protein cleavage, and particularly the need for such long cleavage times, is not well understood, and further investigation may be useful.

Factor Xa is a heterodimer protease composed of two disulfide-linked polypeptide chains with apparent molecular masses of 17 and 42 kDa (42). In this study, no obvious bands of these sizes were observed, which may be due to too low a concentration to be detected in SDS-PAGE analysis. Thus, removal of factor Xa using *p*-aminobenzamidinium resin was not necessary. ELISA results for MBP fusion proteins were analyzed using OD values obtained for MBP fusion proteins without subtracting the OD values obtained for MBP-LacZ to allow for a comparative analysis between the MBP fusion proteins and the factor Xa-cleaved forms. This was justified, as levels of seroreactivity to MBP-LacZ were not significantly different between infected and uninfected samples.

The MBP fusion proteins evaluated in this study were previously reported to be differentially regulated under different *in vitro* stress conditions (8–10). More than one-half of the 50 MBP fusion proteins evaluated were found to be antigenic, suggesting that the majority of these proteins are expressed by *M. avium* subsp. *paratuberculosis* *in vivo* and are recognized by the host immune response. A recent study on the antigenicity of some of these proteins expressed as His-tagged recombinant proteins showed that 66% of the proteins (18/27 proteins) were detected by sera from *M. avium* subsp. *paratuberculosis*-infected sheep (13). Of those 27 proteins, 22 were included in this study. The proteins that were found to be antigenic in both studies are MAP0516c,

TABLE 4 Antigenicity evaluation of MBP fusion and factor Xa-cleaved proteins

Protein	Protein concn (µg/ml)	Data for MBP fusion proteins				Data for factor Xa-cleaved proteins			
		OD ₄₅₀ (n) for:		P ^c	AUC _{ROC}	OD ₄₅₀ (n) for:		P ^c	AUC _{ROC}
		Unexposed samples (23)	Exposed samples (23)			Unexposed samples (23)	Exposed samples (23)		
MAP0593c	20	0.262	0.301	0.295		0.182	0.253	0.005	0.75
MAP1017c	10	0.203	0.266	0.046	0.687	0.206	0.314	0.005	0.753
MAP1588c ^b	10	0.167	0.243	0.027	0.702	0.153	0.206	0.033	0.662
MAP1846c ^b	10	0.158	0.239	0.022	0.702	0.178	0.23	0.029	0.708
MAP1885c	10	0.237	0.261	0.659		0.166	0.169	0.864	
MAP2280c ^b	10	0.196	0.279	0.027	0.702	0.239	0.344	0.002	0.786
MAP2698c	4	0.207	0.268	0.15		0.197	0.239	0.007	0.7
MAP3190 ^{a,b}	10	0.222	0.327	0.007	0.708	0.165	0.224	0.03	0.668
MAP3538 ^b	10	0.172	0.247	0.022	0.703	0.188	0.264	0.001	0.792
MAP3555 ^{a,b}	10	0.272	0.448	0.002	0.736	0.203	0.311	0.002	0.738

^a Proteins able to produce significantly higher OD values for serum samples obtained from sheep with paucibacillary infections than for serum samples from unexposed sheep.

^b Proteins able to produce significantly higher OD values for serum samples that were low reactors in the Institut Pourquier ELISA than for serum samples from unexposed sheep.

^c Mean OD values for the exposed group are significantly higher than those for the unexposed group ($P < 0.05$).

MAP0834c, MAP1846c, MAP2450c, MAP3200, and MAP3555. The proteins that were found to be nonantigenic in both studies are MAP1339, MAP1885c, MAP1889c, MAP2705c, MAP3701c, MAP3577, and MAP4125. The proteins MAP0435c, MAP0593c, MAP1027c, MAP2281c, and MAP2411 were found to be antigenic by Kawaji et al. (13) but were not antigenic in this study. Similarly, proteins MAP0068, MAP2007, MAP2864c, and MAP3864 were found to be antigenic in this study but not by Kawaji et al. (13). Disagreement in the results for the remaining proteins may be attributed to factors such as different types of antigen (His-tagged or MBP fusion proteins) and the animals tested.

The antigenicity evaluation results for MBP fusion proteins MAP2698c and MAP3567 were in agreement with the findings from a previous study on the evaluation of their His-tagged recombinant forms (12). Another recent study reported protein MAP1272c NlpC/P60 to be strongly antigenic in cattle infected with *M. avium* subsp. *paratuberculosis* (29). Evaluation of this protein in sheep infected with *M. avium* subsp. *paratuberculosis* in this study also showed strong immunoreactivity, suggesting that this protein may be of potential use in the diagnosis of JD in both sheep and cattle. This is the first antigenicity evaluation of the remaining 28 MBP fusion proteins.

Twelve of the 50 MBP fusion proteins included in this study were previously identified to have a high number of conformational B cell epitopes, and so they may be useful for detection of *M. avium* subsp. *paratuberculosis*-specific serum antibodies in infected hosts (43). Antigenicity evaluation of these proteins in this study revealed 75% (9/12 proteins) to be antigenic.

TABLE 5 Antigenicity evaluation of His-tagged proteins

Protein	Protein concn (µg/ml)	OD ₄₅₀ for:				AUC _{ROC}
		Unexposed samples	Exposed samples	P ^a	AUC _{ROC}	
MAP1272c	5	0.140	0.318	<0.0001	0.90	
MAP2487c	4	0.160	0.297	0.004	0.70	
MAP2698c	2	0.149	0.285	0.003	0.685	
MAP3567	0.5	0.271	0.443	0.009	0.751	

^a Mean OD values for the exposed group are significantly higher than those for the unexposed group ($P < 0.05$).

The proteins found to be antigenic were from clusters such as amino acid metabolism (four proteins), antioxidant enzymes (four proteins), fatty acid metabolism (six proteins), hypothetical proteins (three proteins), proteolysis (two proteins), two-component response regulators (two proteins), and cellular processes (three proteins), as well as cell wall synthesis, cell division, ATP biosynthesis, signal recognition, and a putative virulence factor (one protein each). These clusters of *M. avium* subsp. *paratuberculosis* proteins recognized by the host immune system may be involved in the ability of *M. avium* subsp. *paratuberculosis* to evade host defense mechanisms, which may be augmented by *in vivo* expression. Antioxidant enzymes are important for protecting *M. avium* subsp. *paratuberculosis* from the oxidative stress response mounted by the host (10, 44, 45); similarly, proteins involved in fatty acid metabolism play important roles in intracellular survival, growth of mycobacteria, and pathogenicity in mycobacterial infections (46). Proteins and lipoproteins associated with cell wall synthesis and lipid membranes are critical for protecting mycobacteria from damage (47).

In conclusion, this study has identified several stress-regulated *M. avium* subsp. *paratuberculosis* proteins as being antigenic in infected sheep. Proteins that were found to be able to detect paucibacillary infections and Institut Pourquier ELISA low reactors may be of potential use in early diagnosis. The proteins that were identified as antigenic in this study (MBP fusion or cleaved) and

TABLE 6 Comparison of antigenicity of MBP fusion, His-tagged, and factor Xa-cleaved proteins^a

Protein	Data by protein type					
	MBP fusion		His tagged		Factor Xa cleaved	
	P	AUC _{ROC}	P	AUC _{ROC}	P	AUC _{ROC}
MAP1272c	0.002	0.728	<0.0001	0.90	NE ^b	NE
MAP2487c	0.331	NE	0.004	0.697	NE	NE
MAP2698c	0.150	NE	0.003	0.685	0.007	0.70
MAP3567	0.005	0.703	0.009	0.751	NE	NE

^a P and AUC_{ROC} values are for comparisons between infected and uninfected groups of sheep. Mean OD values for the exposed group are significantly higher than those for the unexposed group ($P < 0.05$).

^b NE, not examined.

by Kawaji et al. (His-tagged) (13) may be of potential use in the diagnosis and control of JD. The antigenicity results for cleaved or His-tagged *M. avium* subsp. *paratuberculosis* proteins were marginally superior to those for MBP fusion forms. Further evaluation of these proteins using a larger panel of serum samples without the bias of prior positive ELISA results is required to select potentially useful proteins for diagnosis.

ACKNOWLEDGMENTS

This work was supported by Meat and Livestock Australia and by the Cattle Council of Australia, the Sheepmeat Council of Australia, and WoolProducers Australia through Animal Health Australia.

REFERENCES

- Morris CA, Hickey SM, Henderson HV. 2006. The effect of Johne's disease on production traits in Romney, Merino and Merino × Romney-cross ewes. *N. Z. Vet. J.* 54:204–209.
- Ott SL, Wells SJ, Wagner BA. 1999. Herd-level economic losses associated with Johne's disease on US dairy operations. *Prev. Vet. Med.* 40:179–192.
- Hope AF, Kluver PF, Jones SL, Condrion RJ. 2000. Sensitivity and specificity of two serological tests for the detection of ovine paratuberculosis. *Aust. Vet. J.* 78:850–856.
- Nielsen SS, Toft N. 2008. Ante mortem diagnosis of paratuberculosis: a review of accuracies of ELISA, interferon- γ assay and faecal culture techniques. *Vet. Microbiol.* 129:217–235.
- Yokomizo Y, Merkal RS, Lyle PA. 1983. Enzyme-linked immunosorbent assay for detection of bovine immunoglobulin G1 antibody to a protoplasmic antigen of *Mycobacterium paratuberculosis*. *Am. J. Vet. Res.* 44:2205–2207.
- Yokomizo Y, Yugi H, Merkal RS. 1985. A method for avoiding false-positive reactions in an enzyme-linked immunosorbent assay (ELISA) for the diagnosis of bovine paratuberculosis. *Nihon Juigaku Zasshi* 47:111–119.
- Whittington RJ, Marshall DJ, Nicholls PJ, Marsh IB, Reddacliff LA. 2004. Survival and dormancy of *Mycobacterium avium* subsp. *paratuberculosis* in the environment. *Appl. Environ. Microbiol.* 70:2989–3004.
- Gumber S, Taylor DL, Marsh IB, Whittington RJ. 2009. Growth pattern and partial proteome of *Mycobacterium avium* subsp. *paratuberculosis* during the stress response to hypoxia and nutrient starvation. *Vet. Microbiol.* 133:344–357.
- Gumber S, Whittington RJ. 2009. Analysis of the growth pattern, survival and proteome of *Mycobacterium avium* subsp. *paratuberculosis* following exposure to heat. *Vet. Microbiol.* 136:82–90.
- Kawaji S, Zhong L, Whittington RJ. 2010. Partial proteome of *Mycobacterium avium* subsp. *paratuberculosis* under oxidative and nitrosative stress. *Vet. Microbiol.* 145:252–264.
- Gumber S, Taylor DL, Whittington RJ. 2009. Evaluation of the immunogenicity of recombinant stress-associated proteins during *Mycobacterium avium* subsp. *paratuberculosis* infection: implications for pathogenesis and diagnosis. *Vet. Microbiol.* 137:290–296.
- Gurung RB, Purdie AC, Begg DJ, Whittington RJ. 2012. *In silico* screened *Mycobacterium avium* subsp. *paratuberculosis* (MAP) recombinant proteins upregulated under stress conditions are immunogenic in sheep. *Vet. Immunol. Immunopathol.* 149:186–196.
- Kawaji S, Gumber S, Whittington RJ. 2012. Evaluation of the immunogenicity of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) stress-associated recombinant proteins. *Vet. Microbiol.* 155:298–309.
- Banasure KD, Basagoudanavar SH, Chaudhury P, Tiwari V, Parihar NS, Goswami PP. 2001. Identification and characterization of a gene encoding a 35-kDa protein from *Mycobacterium avium* subspecies *paratuberculosis*. *FEMS Microbiol. Lett.* 196:195–199.
- Bannantine JP, Paulson AL, Chacon O, Fenton RJ, Zinniel DK, McVey DS, Smith DR, Czuprynski CJ, Barletta RG. 2011. Immunogenicity and reactivity of novel *Mycobacterium avium* subsp. *paratuberculosis* PPE MAP1152 and conserved MAP1156 proteins with sera from experimentally and naturally infected animals. *Clin. Vaccine Immunol.* 18:105–112.
- Bannantine JP, Paustian ML. 2006. Identification of diagnostic proteins in *Mycobacterium avium* subspecies *paratuberculosis* by a whole genome analysis approach. *Methods Mol. Biol.* 345:185–196.
- Cho D, Sung N, Collins MT. 2006. Identification of proteins of potential diagnostic value for bovine paratuberculosis. *Proteomics* 6:5785–5794.
- Hughes V, Bannantine JP, Denham S, Smith S, Garcia-Sanchez A, Sales J, Paustian ML, McLean K, Stevenson K. 2008. Immunogenicity of proteome-determined *Mycobacterium avium* subsp. *paratuberculosis*-specific proteins in sheep with paratuberculosis. *Clin. Vaccine Immunol.* 15:1824–1833.
- Koets AP, Rutten V, Hoek A, Bakker D, van Zijderveld F, Muller KE, van Eden W. 1999. Heat-shock protein-specific T-cell responses in various stages of bovine paratuberculosis. *Vet. Immunol. Immunopathol.* 70:105–115.
- Leroy B, Viart S, Trincherio N, Roupie V, Govaerts M, Letesson JJ, Huygen K, Wattiez R. 2009. Use of *Mycobacterium avium* subsp. *paratuberculosis* specific coding sequences for serodiagnosis of bovine paratuberculosis. *Vet. Microbiol.* 135:313–319.
- Muller J, Michal I, Fishman Y, Hovav AH, Barletta RG, Bercovier H. 2002. The immunogenicity of *Mycobacterium paratuberculosis* 85B antigen. *Med. Microbiol. Immunol.* 190:179–187.
- Olsen I, Boysen P, Kulberg S, Hope JC, Jungersen G, Storset AK. 2005. Bovine NK cells can produce gamma interferon in response to the secreted mycobacterial proteins ESAT-6 and MPP14 but not in response to MPB70. *Infect. Immun.* 73:5628–5635.
- Bannantine JP, Hansen JK, Paustian ML, Amonsin A, Li LL, Stabel JR, Kapur V. 2004. Expression and immunogenicity of proteins encoded by sequences specific to *Mycobacterium avium* subsp. *paratuberculosis*. *J. Clin. Microbiol.* 42:106–114.
- Sachdev D, Chirgwin JM. 2000. Fusions to maltose-binding protein: control of folding and solubility in protein purification. *Methods Enzymol.* 326:312–321.
- Bannantine JP, Rosu V, Zanetti S, Rocca S, Ahmed N, Sechi LA. 2008. Antigenic profiles of recombinant proteins from *Mycobacterium avium* subsp. *paratuberculosis* in sheep with Johne's disease. *Vet. Immunol. Immunopathol.* 122:116–125.
- Pediaditakis P, Monga SPS, Mars WM, Michalopoulos GK. 2002. Differential mitogenic effects of single chain hepatocyte growth factor (HGF)/scatter factor and HGF/NK1 following cleavage by factor Xa. *J. Biol. Chem.* 277:14109–14115.
- Liu D, Zou L, Li W, Wang L, Wu Y. 2009. High-level expression and large-scale preparation of soluble HBx antigen from *Escherichia coli*. *Biotechnol. Appl. Biochem.* 54:141–147.
- Pasek M, Boeggeman E, Ramakrishnan B, Qasba PK. 2010. Galectin-1 as a fusion partner for the production of soluble and folded human β -1,4-galactosyltransferase-T7 in *E. coli*. *Biochem. Biophys. Res. Commun.* 394:679–684.
- Bannantine JP, Lingle CK, Stabel JR, Ramyar KX, Garcia BL, Raeber AJ, Schacher P, Kapur V, Geisbrecht BV. 2012. MAP1272c encodes an NlpC/P60 protein, an antigen detected in cattle with Johne's disease. *Clin. Vaccine Immunol.* 19:1083–1092.
- Bannantine JP, Stabel JR, Bayles DO, Geisbrecht BV. 2010. Characteristics of an extensive *Mycobacterium avium* subspecies *paratuberculosis* recombinant protein set. *Protein Expr. Purif.* 72:223–233.
- Ellis TM, Norris RT, Martin P, Casey RH, Hawkins CD. 1998. Evidence for freedom from Johne's disease in cattle and goats in Western Australia. *Aust. Vet. J.* 76:630–633.
- Gumber S, Eamens G, Whittington RJ. 2006. Evaluation of a Pourquier ELISA kit in relation to agar gel immunodiffusion (AGID) test for assessment of the humoral immune response in sheep and goats with and without *Mycobacterium paratuberculosis* infection. *Vet. Microbiol.* 115:91–101.
- Perez V, Marin JFG, Badiola JJ. 1996. Description and classification of different types of lesion associated with natural paratuberculosis infection in sheep. *J. Comp. Pathol.* 114:107–122.
- Begg DJ, de Silva K, Di Fiore L, Taylor DL, Bower K, Zhong L, Kawaji S, Emery D, Whittington RJ. 2010. Experimental infection model for Johne's disease using a lyophilised, pure culture, seedstock of *Mycobacterium avium* subspecies *paratuberculosis*. *Vet. Microbiol.* 141:301–311.
- Burton AB, Wagner B, Erb HN, Ainsworth DM. 2009. Serum interleukin-6 (IL-6) and IL-10 concentrations in normal and septic neonatal foals. *Vet. Immunol. Immunopathol.* 132:122–128.

36. Zweig MH, Campbell G. 1993. Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin. Chem.* 39:561–577.
37. Eaton D, Rodriguez H, Vehar GA. 1986. Proteolytic processing of human factor VIII: correlation of specific cleavages by thrombin, factor Xa, and activated protein C with activation and inactivation of factor VIII coagulant activity. *Biochemistry* 25:505–512.
38. Nagai K, Perutz MF, Poyart C. 1985. Oxygen binding properties of human mutant hemoglobins synthesized by *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* 82:7252–7255.
39. Huang X, Crute BE, Sun C, Tang YY, Kelley JJ, III, Lewis AF, Hartman KI, Laue TM, Speck NA, Bushweller JH. 1998. Overexpression, purification, and biophysical characterization of the heterodimerization domain of the core-binding factor β subunit. *J. Biol. Chem.* 273:2480–2487.
40. Ludeman JP, Pike RN, Bromfield KM, Duggan PJ, Cianci J, Le Bonniec B, Whisstock JC, Bottomley SP. 2003. Determination of the P₁' , P₂' and P₃' subsite-specificity of factor Xa. *Int. J. Biochem. Cell Biol.* 35:221–225.
41. He M, Jin L, Austen B. 1993. Specificity of factor Xa in the cleavage of fusion proteins. *J. Protein Chem.* 12:1–5.
42. Himmelspach M, Pfleiderer M, Fischer BE, Plaimauer B, Antoine G, Falkner FG, Dörner F, Schlokot U. 2000. Recombinant human factor X: high yield expression and the role of furin in proteolytic maturation in vivo and in vitro. *Thromb. Res.* 97:51–67.
43. Gurung RB, Purdie AC, Begg DJ, Whittington RJ. 2012. *In silico* identification of epitopes in *Mycobacterium avium* subsp. *paratuberculosis* proteins that were upregulated under stress conditions. *Clin. Vaccine Immunol.* 19:855–864.
44. Zahrt TC, Deretic V. 2002. Reactive nitrogen and oxygen intermediates and bacterial defenses: unusual adaptations in *Mycobacterium tuberculosis*. *Antioxid. Redox Signal.* 4:141–159.
45. Zhu X, Tu ZJ, Coussens PM, Kapur V, Janagama H, Naser S, Sreevatsan S. 2008. Transcriptional analysis of diverse strains *Mycobacterium avium* subspecies *paratuberculosis* in primary bovine monocyte derived macrophages. *Microbes Infect.* 10:1274–1282.
46. Dyer DH, Lyle KS, Rayment I, Fox BG. 2005. X-ray structure of putative acyl-ACP desaturase DesA2 from *Mycobacterium tuberculosis* H37Rv. *Protein Sci.* 14:1508–1517.
47. He Z, De Buck J. 2010. Localization of proteins in the cell wall of *Mycobacterium avium* subsp. *paratuberculosis* K10 by proteomic analysis. *Proteome Sci.* 8:21. doi:10.1186/1477-5956-8-21.