

1936

EC1415 Nebraska Emergency Mash Formula #1936

Follow this and additional works at: <http://digitalcommons.unl.edu/extensionhist>

"EC1415 Nebraska Emergency Mash Formula #1936" (1936). *Historical Materials from University of Nebraska-Lincoln Extension*. 2531.
<http://digitalcommons.unl.edu/extensionhist/2531>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

585
E7
#1415 c.1

EC 1415

Nebraska
COOPERATIVE EXTENSION WORK
IN AGRICULTURE AND HOME ECONOMICS
U. of N. Agr. College & U. S. Dept. of Agr. Cooperating
W. H. Brokaw, Director, Lincoln

Extension
Circular
1415

NEBRASKA EMERGENCY MASH FORMULA #1936

	No. 36	No. 36M (With Skimmilk)
	Lbs.	Lbs.
1. Shorts.....	320*	420
2. Bran.....	200	200
3. Pulverized oats or barley.....	200	200
4. Alfalfa meal (must be green and high in protein)....	100	100
5. Soybean oil meal.....	75	25
6. Meat scraps or fish meal.....	75	25
7. Limestone (pulverized).....	20	20
8. Salt.....	10	10
	1000	1000

RECEIVED

OCT 19 1972

*If some corn is available, replace 120 pounds shorts with cormeal.

Oyster shell or limestone grit constantly available.

C. Y. THOMPSON
LIBRARY

Feed grain, wheat, oats and barley; depending upon price.

Hopper feed mash and grain to growing turkeys and pullets. Hopper feed only mash to laying hens.

Wetting mash in extremely hot weather will keep up consumption and maintain growth and production.

Feed the same mash to growing pullets, turkeys and laying hens.

Feed requirements for turkeys, pullets and laying hens.

	1st Mo.	2nd Mo.	3rd Mo.	4th Mo.	5th Mo.	6th Mo.	7th Mo.
Young Turkeys	1.02#	3.07#	7.48#	11.6#	14.82#	18.0#	20.4#

Leghorn pullets require approximately 24 pounds of feed up to laying age. Heavy breed pullets require approximately 28 pounds of feed up to laying age. Laying hens use approximately seven pounds of feed per month (grain and mash). Chickens use approximately equal portions of grain and mash. Turkeys use about one and one-half times as much mash as grain (45# mash and 30# of grain).

Rigid culling is recommended--it does not pay to feed the scrubs.

Use waste-proof, wind-proof and filth-proof feeders. Provide shade and keep feeders and waterers in shady places.

Get rid of mites--they sap the profits. Treat houses and fixtures with wood preservative.

Lice are almost as bad as mites. Treat hens with nicotine sulfate or sodium fluoride.

For details regarding these recommendations see County Agricultural Agent or write to the College of Agriculture.

16964fr