

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of Nebraska-
Lincoln Extension

Extension

2-1951

EC1496 Revised 1951 Poultry Rations and Feeding

W. F. Aubol

Follow this and additional works at: <http://digitalcommons.unl.edu/extensionhist>

Aubol, W. F., "EC1496 Revised 1951 Poultry Rations and Feeding" (1951). *Historical Materials from University of Nebraska-Lincoln Extension*. 2652.

<http://digitalcommons.unl.edu/extensionhist/2652>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Nebr. A. Un 3/3.4:1496

S Feb.

85 1951

E7

#1496R

C.1

E.C. 1496

Revised -


POULTRY RATIONS

and FEEDING

UNIVERSITY OF NEBRASKA
LIBRARY

APR 27 1951

DOCUMENTS
ROOM



EXTENSION SERVICE

UNIVERSITY OF NEBRASKA COLLEGE OF AGRICULTURE

AND U. S. DEPARTMENT OF AGRICULTURE

COOPERATING

W. V. LAMBERT, DIRECTOR

POULTRY RATIONS AND FEEDING

Extension Poultry Husbandman
W. F. Aubol

Efficient feeding is essential if the poultry enterprise is to prove successful. By efficient feeding is meant supplying feeds that will maintain health, growth and production at prices justifying their use. The feeding of poultry is probably the most important single factor in the maintenance of health and production. This will be reflected in profitableness of the flock.

Since 60 to 70 percent of the poultryman's total production cost is attributable to feed, the efficient manager will adjust feeding practices to fluctuations in prices of ingredients yet maintain egg production in order to secure maximum returns.

When grain is being fed in small quantities it may be fed on top of the mash in the feeders. When fed in larger quantities it should be placed in separate feeders. The proportion of grain to mash that is consumed can be controlled by the number of feeders used for each.

Quantity Buying

Many flock owners do not have flocks large enough to justify buying ready-mixed feed in quantities of one ton or more, or large amounts of ingredients for home mixing of mash.

Savings can often be made in buying feed if several producers pool their orders to enable them to make quantity purchases.

Feed that is stored for an excessive period of time will often deteriorate and lose some of its value. When buying quantities of feed in excess of needs for a reasonable period of time, precautions should be taken to prevent deterioration with a loss in feeding value.

Complete Chick Starting-Mash

	Lbs.
Yellow cornmeal.....	400
Soybean meal.....	224
Ground Oats	100
Shorts or Bran	100
Meat Scraps	50
Alfalfa Meal (17% Protein +)	50
*Fish Meal (or fish soluble blend)	25
Dried Whey	25
**Mineral Mixture No. 45	25
***Act. animal sterol (1500 IC units vitamin D per gm.)	1
Antibiotic Feed Supplement (Follow Manufacturers Directions)	
Vitamin Concentrate (Follow Manufacturers Di- rections)	
	<u>1000</u>
Average Protein.....	20.8%

The ration shown above can be used as a complete start-
ingration. Chicks are started on this ration and continue
until they reach four to six weeks of age when they are
gradually changed to an all-purpose mash. One hundred
chicks will require about 200 pounds of this mash through
the first six weeks.

*See page seven.

**Mineral mixture Nos. 45 & 50	No. 45 Lbs.	No. 50 Lbs.
Ground limestone.....	66.2	33.2
Iodized salt	33.0	33.0
Steamed bone meal	--	33.0
Manganese sulphate.....	.8	.8
	<u>100.0</u>	<u>100.0</u>

***See page seven.

Nebraska Broiler Ration No. 970

	Lbs.
Yellow cornmeal.....	500
Soybean meal.....	317
Meat scraps	50
Alfalfa meal (17% prot. or better)	50
*Fish soluble blend or fish meal	25
**Mineral Mix No. 50	25
Dried Whey	25
***Act. animal sterol (1500 IC units vitamin D per gm.)	1
Antibiotic Feed Supplement (Follow Manufacturers Directions)	
Vitamin Concentrate (Follow Manufacturers Di- rections)	1000
Average Protein.....	23%

Broiler ration No. 970 is complete and can be used to start the chicks and its use can be continued throughout the growing period. A small amount of whole oats may be fed in addition to this mash beginning with the eighth week. Oats should not be fed in amounts in excess of 10% of mash consumed. One hundred broilers require 1,100 to 1,200 pounds for the 12-week growing period.

This formula has given very high growth rates in recent experiments of broiler production at the University of Nebraska. It is especially well adapted to use here in Nebraska since it is made up of 50% corn, which is usually low priced for poultry feed; and also provides a very high energy feed which will produce rapid growth.

*See page seven.

**See page three.

***See page seven.

All-Purpose or Laying Mash

	Lbs.
Yellow cornmeal.....	310
Wheat middlings.....	200
Soybean oil meal.....	200
Whole oats.....	100
Alfalfa meal (17% protein +).....	50
Meat and bone scrap (50%).....	50
*Fish meal or fish soluble blend.....	27
Dried whey.....	25
**Mineral Mixture No. 45.....	30
***Act. animal sterol (1500 IC units vitamin D per gm.)	2
Antibiotic feed supplement (Follow Manufacturers Directions)	
Vitamin Concentrate (Follow Manufacturers Di- rections)	1000
Average Protein.....	21.5%

The all-purpose mash can be used to replace the chick starter for pullets after they have reached four to six weeks of age. Pullets fed all-purpose mash should be given some oats from the eighth week throughout the growing season. While on the range, pullets may be fed equal parts of mash and a mixture of corn and oats.

When this mash is fed to laying hens home-grown grain may be used in varying proportions but not in excess of an amount equal to the mash consumed. The grain may consist chiefly of oats with enough corn added to maintain the the body weight of the hens. One hundred laying hens require about 400 pounds of this mash per month in addition to about 400 pounds of grain.

*See page seven.

**See page three.

***See page seven.

Nebraska 49-X Protein Supplement 26% Protein

	Lbs.
Yellow cornmeal.....	150
Shorts	300
Alfalfa meal (17% protein +).....	100
Meat scraps.....	100
*Fish soluble blend (or fish meal)	50
Soybean meal.....	170
Corn gluten meal	100
**Mineral mix No. 45	30
***Act. animal sterol (1500 IC units vitamin D per gm.)	2
	1000

Oats and corn can be fed with the 26% protein supplement given above. Each of the grains can be fed in an amount equal to the mash. Wheat if available can be used to replace one-half of the oats. If skim milk or buttermilk is available, it may be fed at the rate of three gallons per hundred hens per day to replace about four pounds of the supplement. One hundred laying hens will require about 275 pounds of this supplement and 500 pounds of grain per month.

Suggested Feeding Program:

	Lbs.	Protein
100 lbs. 49-X.....	26	
100 lbs. whole yellow corn.....	10	
100 lbs. whole oats or wheat or half of each	12	
	48	
Protein level entire ration		16%

*See page seven.

**See page three.

***See page seven.

34% Protein Concentrate

	Lbs.
Shorts.....	100
Alfalfa meal (17% protein +)	100
Meat scraps	100
*Fish soluble blend (or fish meal).....	50
Soybean meal	170
Corn gluten meal.....	100
**Mineral mix No. 45.....	30
***Act. animal sterol (1500 IC units vitamin D per g.)	2
	650

Hens eat whole grains readily and are equipped to grind these grains efficiently. For this reason it is usually more economical for a farmer to buy, or have mixed a concentrate so that a larger amount of homegrown grains can be used.

Suggested Feeding Program:

	Lbs.	Protein
100 lbs. 34% concentrate.....	34	
200 lbs. whole yellow corn.....	20	
200 lbs. whole oats or wheat or mixture	24	
of oats and wheat	24	
	78	

Protein level of entire ration	15.6%
--------------------------------	-------

*Fish soluble blend may be prepared by mixing 40% fish solubles with 60% alfalfa meal. Fish meal may be used as a substitute but the cost is frequently prohibitive. Several commercial products will give comparable results.

**See page three.

***Vitamin D is also found in fish liver oils. When chickens have access to adequate sunshine as in summer it is unnecessary to include Vitamin D in the ration.

The requirements for 100 laying hens are as follows:

1. Thirty-two linear feet of feeder space (four feeders four feet long).
2. Twenty-five to twenty-seven pounds of feed per day.
3. One large round fountain or a three-foot trough for water.
4. Seven to eight gallons of water per day.
5. Oyster shell, free choice.
6. Grit (road gravel), free choice.

The requirements for 100 broilers are as follows:

1. Twenty-four linear feet of feeder space (three feeders four feet in length).
2. Four one-gallon or two 2-gallon waterers.

The requirements for starting 100 chicks (first 6 weeks) are as follows:

1. Two feeders four feet in length.
2. Two one-gallon waterers.

