

1987

EC87-1511 Field Crop Insect Control Guide for Nebraska : Alfalfa, Soybeans, Wheat, Range and Pasture

D. L. Keith

F. P. Baxendale

University of Nebraska-Lincoln, fbaxendale1@unl.edu

L. L. Peters

Follow this and additional works at: <http://digitalcommons.unl.edu/extensionhist>

Keith, D. L.; Baxendale, F. P.; and Peters, L. L., "EC87-1511 Field Crop Insect Control Guide for Nebraska : Alfalfa, Soybeans, Wheat, Range and Pasture" (1987). *Historical Materials from University of Nebraska-Lincoln Extension*. 4829.
<http://digitalcommons.unl.edu/extensionhist/4829>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

AGRI
S
85
E7

FIELD CROP INSECT CONTROL GUIDE FOR NEBRASKA
ALFALFA, SOYBEANS, WHEAT, RANGE AND PASTURE

RECEIVED

FEB 27 1987

D.L. Keith, F.P. Baxendale, L.L. Peters, J.F. Witkowski,
J.B. Campbell, A.F. Hagen, S.D. Danielson
Extension Entomology Specialists
K.J. Jarvi, IPM Extension Assistant
R.C. Seymour, IPM Extension Assistant
J.A. Kalisch, Extension Technologist (Entomology)

UNIVERSITY OF NEBRASKA
LIBRARIES

Insect control suggestions in this circular are based on University of Nebraska test results, data from surrounding states, USDA recommendations, previous experience, and label registrations. These suggestions are designed to guide Nebraska farmers when they select an insect control program. NebGuides and other publications containing additional information on insect identification, damage, and life cycles are referenced under insect headings and are available by mail order (Write - Bulletins, 104 ACB, University of Nebraska-Lincoln, NE 68583-0918) or from Cooperative Extension Service Offices.

INDEX TO THIS PUBLICATION

Subject	Page No.	Subject	Page No.
Abbreviations	4	Grasshoppers	
Alfalfa Insects	4	Alfalfa	8
Alfalfa Weevil	4	Range/Pasture	16
Adults	5	Soybeans	10
Larvae (Worms)	4	Wheat	14
Aphids/Alfalfa	5	Green Cloverworm/Soybeans	10
Army Cutworms/Wheat	13	Greenbugs/Wheat	14
Armyworms		Hessian Fly/Wheat	15
Range and Pasture	16	Insecticide	
Wheat	13	Field Re-Entry Periods	3
Bean Leaf Beetle/Soybeans	9	Policy For Making Pesticide Suggestions	1
Black Grass Bugs/Range and Pasture	16	Restricted Use	2
Blister Beetle/Alfalfa	9	Sprayed by Mistake?	3
Chemigation	2	Toxicity	1
Chewing Insects/Soybeans	11	Pale Western Cutworm/Wheat	13
Chinch Bugs	15	Potato Leafhopper/Alfalfa	7
Conservation Tillage/Insect Pests	2	Protect Bees	2
Container Disposal	3	Range and Pasture Insects	15
Control Decision Guidelines/ Economic Thresholds	2	Russian Wheat Aphid	14
Cutworms		Soil Cutworms/Soybeans	11
Alfalfa	6	Soybean Insecticide Restrictions	12
Soybeans	11	Soybean Insects	9
Fall Armyworms/Wheat	13	Spider Mites/Soybeans	12
		Stored Grain, Insect Prev/Control	2
		Webworms/Alfalfa	6
		Wheat Insects	13
		Wireworms/Soybeans	11

POLICY FOR MAKING PESTICIDE SUGGESTIONS

The choice of a pesticide should not be based entirely on its cost. Several other factors need to be considered in the decision, including efficacy against the target pest or pest combination, formulation of the pesticide, label restrictions, safety to non-target species (including humans) and environmental conditions at the time of application.

THE USER IS RESPONSIBLE FOR THE EFFECTS OF PESTICIDE RESIDUES ON CROPS AND LIVESTOCK, AS WELL AS PESTICIDE DRIFT AND CONTAMINATION. This publication does not supersede label information. Always read and carefully follow the instructions on the container label. For current information, contact your local Cooperative Extension Service Office.

The use of trade names in this circular is not to be considered an endorsement by the Nebraska Cooperative Extension Service, and no discrimination is intended.

TOXICITY OF INSECTICIDES

(NebGuides G85-758, G84-715, G79-460, G79-472, G79-473, G79-479)

All insecticides are poisonous and must be used with caution. Always store them in their original containers out of the reach of children, unauthorized personnel, and livestock. Skull and crossbones and the words Danger/Poison appear in red on the label of highly toxic materials. Liquid formulations of these products are not recommended for farmer application. However, farmers should be able to safely use granular formulations of these chemicals if they use proper precautions, as indicated on the label. Moderate and low toxicity pesticides are marked with the signal words Warning and Caution, respectively.



Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Leo E. Lucas, Director of Cooperative Extension Service, University of Nebraska, Institute of Agriculture and Natural Resources.



<R> RESTRICTED USE <R>

Several insecticides listed in this circular are classified RESTRICTED USE by the Environmental Protection Agency. These compounds are marked with a <R>. Pesticides may be classified as Restricted Use based on their persistence, toxicity or potential environmental hazards. To purchase and use these products, EPA certification is required. Your local Cooperative Extension office will have a listing of the dates and locations where certification training can be obtained. Remember that the status of a formulation can change at any time. When purchasing a chemical, be certain to ask the dealer if the attached label is up to date.

CONTROL DECISION GUIDELINES/ECONOMIC THRESHOLDS

Economic thresholds are flexible guidelines. They indicate the level of insect abundance that can be tolerated before control actions should be taken. THEY ARE NOT HARD RULES THAT APPLY TO EVERY SITUATION. Used conscientiously, they should be helpful in making treatment decisions. Many variables can affect your decision to treat including insect abundance, anticipated value of the crop, relative effectiveness of controls, and cost of the pesticide and its application. Timing and accuracy of application, plus the effects of weather, also determine the ultimate degree of control.

CHEMIGATION

The term "chemigation" refers to the injection and application of chemicals through irrigation systems. Proper equipment needs, procedures for calibration and other instructions for application through center pivot systems are provided in NebGuides - G84-703 (Applying Insecticides Through Center Pivots) and G73-43 (Anti-Pollution Devices for Applying Chemicals Through Irrigation Systems, Revised August, 1984). The Nebraska Chemigation Act takes effect January 1, 1987. Consult your local Cooperative Extension Service office or Natural Resources District office before chemigating.

INSECT PREVENTION AND CONTROL IN FARM-STORED GRAIN

Properly managed stored grain should have few insect problems during the first year of storage in Nebraska. If grain is to be stored for more than one year, additional effort is required to maintain quality. All bins should be inspected regularly for moisture content, temperature, mold development, and insect pests. Push a sharp pointed stick or rod into the grain to see if hard, compacted areas are developing. Check grain temperature at several locations. Moisture content should also be monitored, even if the grain was dry when binned. Collect several grain samples with a grain probe. Screen the samples and look for evidence of insects. Take samples from the surface and as deep as possible into the grain. Turn on the aeration fan for a few minutes and smell the air. Does it smell normal or does it have a spoiled or musty odor? In addition to following sound sanitation practices, using good aeration management, and making regular inspections, insecticides and/or fumigation may be needed. For more detailed information on insect control and a listing of currently recommended bin sprays, grain protectants, and fumigants, see EC 84-1534, Revised April, 1986, "Pest Management of Farm-Stored Grain," available at your local Cooperative Extension Service Office.

INSECT PESTS IN CONSERVATION TILLAGE SYSTEMS

Modifications of the crop environment in conservation tillage systems could alter the relative importance of Nebraska's more traditional insect pests and possibly create conditions where incidental organisms achieve pest status. Cooler soil temperatures and slower drying in reduced tillage fields may delay planting in the spring and slow seed germination, leaving seeds and young plants susceptible to soil insects for longer periods. Improper placement and incorporation of soil insecticides may create additional problems where crop residues are heavy. Elimination of deep plowing and the resulting increased surface debris may permit certain insects to overwinter in greater numbers. Other conservation tillage practices should result in a reduction in insect numbers. While the use of reduced tillage raises some questions regarding pest control, anticipated problems should not be a barrier to the development of new techniques. It is highly unlikely that insects will be damaging in all fields, and the beneficial effects of conservation tillage appear to outweigh insect control considerations.

PROTECT BEES

Honeybees do not observe manmade boundaries but will collect nectar and/or pollen wherever they can, including field crops such as corn, sorghum, soybeans and alfalfa. If bee colonies are nearby when fields are sprayed during flowering (pollen shed stage for corn and sorghum), they may be killed in substantial numbers. To avoid injury to important pollinators, try to observe the following precautions: 1) Treat only if pests are abundant enough to justify it; 2) if possible, treat only when crops are not in bloom; 3) be certain that honeybee colonies are not directly sprayed with any insecticide; 4) check the crop for heavy concentration of blooming weeds; 5) spray only those parts of fields that have significant pest infestations; 6) if a reasonable choice is available, select an insecticide formulation that is less toxic to bees; and 7) time the application later in the evening when bees are not actively foraging.

SPRAYED BY MISTAKE?

Gardens, particularly plantings of sweet corn, are often placed in or adjacent to crop fields that may be sprayed with an insecticide. The produce is safe to eat IF THE INSECTICIDE IS REGISTERED FOR USE ON THE VEGETABLE OR FRUIT AND THE SPECIFIED WAITING PERIOD HAS ELAPSED. We do not recommend using vegetables or fruit that have been treated with a pesticide which is not labeled for that vegetable or fruit. The following are some preharvest intervals (waiting periods). Check appropriate labels for any others. If you have questions regarding accidental treatments, determine the specific pesticide formulation used, the application rate, and time of spraying. Then, by checking the pesticide label, an informed decision can be made concerning use of the crop.

Minimum Number of Days Between Application and Harvest For Some Selected Crops

Insecticide	Tomatoes	Peppers	Sweet Corn	Cucumbers	Broccoli	Apples
Ambush 2E	NR	3	1	NR	1	***
Counter 15G	NR	NR	30	NR	NR	NR
Cygon 400	7	0	NR	3	7	28
Diazinon AG500	1	5	0	7	5	NR
Diazinon 14G	**	**	**	**	**	NR
Dipel 2X	0	0	0	0	0	0
Di-Syston 8EC	30	NR	28	NR	14	NR
Di-Syston 15G	30	NR	40	NR	14	NR
Dyfonate 4EC	*	*	*	NR	*	NR
Dyfonate 20G	NR	NR	30	NR	*	NR
Furadan 4F	NR	NR	7	NR	NR	NR
Furadan 15G	NR	21	**	**	NR	NR
Imidan 50WP	NR	NR	14	NR	NR	7
Lannate 1.8L	1	3	0	1-3	3	8
Lorsban 4E	NR	NR	35	NR	30	NR
Malathion EC	1-5	3	5	1	3	3
Metasystox-R	NR	**	7-21	**	7	NR
Nudrin 1.8L	1	3	0	1-3	1-3	8
Parathion 8E (ethyl)	10	15	12	15	7	14
Pennac-M	15	NR	3	NR	NR	14
Pounce 3.2EC	0	3	1	NR	1	***
Pydrin 2.4EC	1	7	1	3	3	21
Sevin 80S	0	0	0	0	3	1
Thimet 20G	NR	NR	*	NR	NR	NR

NR = Not Registered

* = At or prior to planting time application only

** = Registered, preharvest interval not indicated on label

*** = Do not apply after petal fall

SOME SUGGESTED FIELD RE-ENTRY PERIODS

Re-entry periods may be listed on the label. FOLLOW LABEL DIRECTIONS AND DO NOT ENTER FIELDS AFTER TREATMENT UNTIL THE RE-ENTRY PERIOD HAS PASSED.

Aastar 15G - 7 days	Guthion 50WP - 24 hrs
Ambush 2E - When spray is dry	Imidan 50WP - None
Comite 6.5EC - 48 hrs	Lannate 1.8L, 90SP - When spray is dry
Counter 15G - 7 days	Larvin 3.2F - When spray is dry
Cygon 400 - 48 hrs	Lorsban 4E - 24 hrs
Diazinon AG500 - When spray is dry	15G - Not stated on label
Diazinon 14G - Not stated on label	Malathion EC - 0 days
Dipel 10G, ES - 0 days	Metasystox-R - 48 hrs
Di-Syston 8EC - 24 hrs, 15G - 7 days	methyl parathion - 48 hr
Dyfonate 20G - 24 hrs	Nudrin 1.8L, 90SP - When spray is dry
4EC - Not stated on label	Pay-Off 2.5EC - When spray is dry
Dylox 80SP - 0 days	Pennac-M - When spray is dry
EPN 4EC, 5EC - 24 hrs	Pounce 3.2EC - When spray is dry
Ethion EC - 24 hrs	Pydrin 2.4EC - When spray is dry
ethyl parathion - 48 hrs	Sevin - All formulations, 0 days
Furadan 4F - 14 days	Thimet 20G - 7 days
15G - None stated on label	

CONTAINER DISPOSAL

Proper disposal of insecticide containers is very important. Serious accidents have occurred when "empty" containers have not been disposed of safely. Suggested methods of disposal are:

Paper Bags: Be certain that all contents have been emptied into applicators or tanks. Burn paper containers in open fields where: 1) regard is given to wind direction in relation to people, domestic animals, and water supplies; 2) where such burning is not in violation of Federal, State or local ordinances; and 3) provisions are made to avoid contamination of surface water.

Metal, Glass, or Plastic Containers: Thoroughly rinse containers at least 3 times with water and dump rinse material into tanks to be used with regular applications. Recycle 5 gallon or larger metal drums where possible after complete rinsing. Containers that cannot be recycled should be punctured, crushed, and buried in a landfill or 24 inches below the soil surface in a location that will not result in contamination of water, crops, man, or animals.

ABBREVIATIONS

AI/A - Active Ingredient Per Acre	Form. - Formulation	LS - Liquid Solution
E - Emulsifiable	G - Granular	Oz - Ounce
EC - Emulsifiable Concentrate	L - Liquid	S - Soluble
ES - Emulsifiable Suspension	lb - Pound	SP - Soluble Powder
F - Flowable	LC - Liquid Concentrate	WP - Wettable Powder

ALFALFA INSECTS

This guide deals only with pests of alfalfa grown for forage. Insects which attack alfalfa grown for seed are discussed in another circular (EC 87-1537).

Protect Pollinators From Insecticides

Protection of honeybees and other pollinators requires communication and cooperation among producers, beekeepers and commercial pesticide applicators. Insecticides, when carelessly applied to legumes or other flowering crops, can destroy substantial numbers of honeybees and result in disastrous losses for beekeepers. The following are suggestions to help reduce bee losses:

1. Ensure through careful inspection that pest densities have reached treatment thresholds before applying an insecticide.
2. Harvest rather than spray if insects are threatening and alfalfa is beginning to bloom.
3. When insecticides are needed, apply when bees are not actively foraging. Late evening applications after bees have returned to the hive and early morning treatments before bees become active outside the hive are less hazardous than mid-day applications.
4. Avoid spraying alfalfa during bloom or if a field has numerous blossoming weeds, such as dandelions or mustards.
5. If insecticides must be applied to blooming alfalfa, notify local beekeepers at least 48 hours in advance so bees can be moved or confined during application.

ALFALFA WEEVIL
(NebGuide G73-30)

Check alfalfa in April and May for the first signs of larval feeding in the tips of alfalfa stems. Earliest damage appears as tiny shotholes on terminal foliage and buds. Look for small, greenish or yellowish larvae, about 1/8 inch long, with dark brown heads and a light stripe down the middle of the back. When fully grown, the larvae are about 3/8 inch long. Plan to treat for alfalfa weevil when an average of 20 or more larvae are caught per sweep of a standard 15 inch sweep net, or if 30-40 percent of the tips are damaged early in May. As the alfalfa matures, the treatment threshold may need to be adjusted upward to 50-60 percent tip feeding. After bud stage, it is probably more profitable to harvest the alfalfa than to treat with a insecticide. Resistant alfalfa varieties are available which will reduce (but not eliminate) damage by the alfalfa weevil.

SPRAYING BLOOMING ALFALFA CAN BE EXTREMELY HAZARDOUS TO BEES.
COORDINATE WITH LOCAL BEEKEEPERS BEFORE APPLYING INSECTICIDES.

REGISTERED FOR LARVAL ALFALFA WEEVIL CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus)	1.0-1.5 lb	No waiting period. Highly toxic to bees. Avoid application during bloom stage. Wait 3 days before harvest.
carbofuran <R> (Furadan 4F)	0.25-1.0 lb	Apply only to pure stands of alfalfa. Do not apply more than once per cutting or twice per year. Do not move bees into treated areas within 7 days of application. See label for harvest restrictions.
chlorpyrifos (Lorsban 4E)	0.5-1.0 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation system.
diazinon AG500, 50W	1.0-1.5 lb	Do not apply during bloom. Wait 10 days before harvest.

malathion ULV 9.33	1.2 lb. (16 oz form.)	Wait 5 days before harvest. Apply when day temperatures are expected to exceed 65° F. Do not apply during bloom. Do not apply to seed alfalfa.
malathion 57EC	1.0-1.25 lb	No waiting period. During bloom, apply only in the evening or early morning.
methidathion <R> (Supracide 2E)	0.5-1.0 lb	Do not apply during bloom. Wait 10 days before harvest. One application per cutting.
methomyl <R> (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.9 lb	Wait 7 days before harvest. Do not apply to dormant or semi-dormant alfalfa when temperature is 50° F or lower. Do not apply when alfalfa is in bloom.
methoxychlor 2EC	1.0-1.5 lb	Wait 7 days before harvest.
methyl parathion <R> (PennCap-M)	0.5-0.75 lb	Do not spray alfalfa during bloom to avoid injury to bees. Do not apply within 15 days of harvest or grazing. May be applied through sprinkler irrigation system.
<R> parathion 4EC, 8E, 8F	0.25-0.5 lb	Wait 15 days before harvest.
phosmet (Imidan 50WP)	1.0 lb	Only 1 application per cutting. Do not graze or harvest for 7 days.

STUBBLE TREATMENTS

Following the cutting and removal of the hay, examine the stubble in several areas for evidence of continued feeding. Sift through windrows and under litter. Check in and around crowns for larvae, cocoons and adult weevils. Adults have long snouts, are light brown to gray with a dark stripe down the back and are about 3/8 inch long. If normal regrowth is not forthcoming after 5-7 days and adult weevils are numerous, a stubble treatment may be necessary.

REGISTERED FOR ADULT ALFALFA WEEVIL CONTROL
(Stubble Treatments)

Insecticide	Rate AI/Acre	Restrictions and Comments
carbofuran <R> (Furadan 4F)	0.5-1.0 lb	Apply only to pure stands of alfalfa. Do not apply more than once per cutting or twice per year. See label for harvest restrictions.
chlorpyrifos (Lorsban 4E)	0.5-1.0 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation system.
methidathion <R> (Supracide 2E)	0.5 lb	Wait 10 days before harvest. One application per cutting.
methyl parathion <R> (PennCap-M)	0.5-0.75 lb	Wait 15 days before harvest. One application per cutting. May be applied through overhead sprinkler irrigation system.
<R> parathion 4EC, 8E, 8F	0.25-0.5 lb	Wait 15 days before harvest.
phosmet (Imidan 50WP)	1.0 lb	Do not apply more than once per cutting.

APHIDS

Aphids are small (about 1/8 inch long) whitish, yellow, green or blue-green insects which suck plant nutrients from alfalfa stems and leaves. Their feeding reduces plant vigor, causes stunting, and may kill cells around the point of entry, either through mechanical destruction or by injection of a toxic saliva. Typically, early symptoms of aphid feeding are yellowish spots on leaves which later merge into larger yellow areas. Eventually, damaged leaves may drop and plants die. In Nebraska, several species of aphids (including the pea aphid, spotted alfalfa aphid and blue alfalfa aphid) infest alfalfa, but fortunately, weather and natural enemies normally hold their numbers below economic threshold levels. Resistant varieties are an important tool in aphid management in alfalfa. Check the latest listing of resistant varieties recommended for your area.

SPRAYING BLOOMING ALFALFA CAN BE EXTREMELY HAZARDOUS TO BEES.
COORDINATE WITH LOCAL BEEKEEPERS BEFORE APPLYING INSECTICIDES.

REGISTERED FOR APHID CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbofuran <R> (Furadan 4F)	0.25-1.0 lb	Apply only to pure stands of alfalfa. Do not apply more than once per cutting or twice per year. Do not move bees into treated areas within 7 days of application. See label for harvest restrictions.
chlorpyrifos (Lorsban 4E)	0.5-1.0 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation system.
diazinon AG500, 50W	0.375-0.5 lb	Wait 7 days before harvest.
dimethoate (Cygon 400)	0.25-0.50 lb	Wait 10 days before harvest or grazing. One application per cutting. Do not apply during bloom.
malathion 57EC	1.0-1.25 lb	No waiting period. During bloom, apply only in the evening or early morning.
methidathion <R> (Supracide 2E)	0.5-1.0 lb	Do not apply during bloom. Wait 10 days before harvest.
methomyl <R> (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.45-0.9 lb	Wait 7 days before harvest. Do not apply to dormant or semi-dormant alfalfa when temperature is 50° F or lower. Do not apply when alfalfa is in bloom.
methyl parathion <R> (PennCap-M)	0.5-0.75 lb	Do not apply during bloom. Wait 15 days before harvest or grazing. Not registered for spotted alfalfa aphid. May be applied through overhead sprinkler irrigation system.
<R> parathion 4EC, 8E, 8F	0.25-0.50 lb	Wait 15 days before harvest.
phosmet + dimethoate (Imidan 50WP)+(Cygon 400)	1.0 lb + 0.125-0.25 lb	Do not apply more than 1 time per cutting. Do not graze or cut for hay within 10 days. Do not apply to alfalfa in the bloom period.

WEBWORMS

Fully developed webworm larvae are yellowish or greenish to nearly black in color. They are 1 to 1 1/4 inches long with dark and light stripes extending down the center of the back and 3 small dark spots on either side of each segment. Infestations are often the result of migrations from weedy areas, so webworms are more commonly a problem in older fields or where stands are thin and weedy. However, webworms also occasionally defoliate established fields and reduce stands of newly seeded alfalfa. When larvae feed, they spin silken webs in the tops of plants and tie leaves together. No Nebraska threshold data are available for webworms, but Texas A & M University entomologists recommend treatment when 25 to 30 percent of plant terminals are infested and the crop is more than 2 weeks from harvest. Otherwise, early cutting is suggested.

REGISTERED FOR WEBWORM CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0-1.5 lb	Wait 3 days before harvest.
<R> parathion 4EC, 8E, 8F	0.25-0.50 lb	Wait 15 days before harvest.
trichlorfon (Dylox 80SP)	0.25-1.0 lb	Do not apply more than 3 times per cutting. No waiting period.

CUTWORMS

Several species of night-flying millers, including the variegated and army cutworm moths, deposit eggs in alfalfa. When larvae emerge and begin to feed, they are about 1/8 inch long, but eventually reach a length of an inch or longer. Cutworms feed at night, usually climbing onto the plants to feed on leaves, and they hide in the soil by day. In newly seeded alfalfa, food reserves in the roots are limited and a small amount of feeding

is enough to kill individual plants. Established alfalfa is not likely to suffer stand loss, but may be delayed and yields reduced if cutworms are numerous. Spray in evening or early morning for best control.

REGISTERED FOR CUTWORM CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus)	1.5 lb	Wait 3 days before harvest.
chlorpyrifos (Lorsban 4E)	0.5 - 1.0 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation systems.
diazinon AG500	2.0-4.0 lb	Wait 7 days before harvest.
methyl parathion <R> (PennCap-M)	0.5-1.0 lb	Do not apply during bloom. Wait 15 days before harvest or grazing. May be applied through overhead sprinkler irrigation system.
<R> parathion 4EC, 8E, 8F	0.25-0.50 lb	Wait 15 days before harvest.
trichlorfon (Dylox 80SP)	0.5-1.0 lb	Do not apply more than 3 times per cutting. No waiting period.

POTATO LEAFHOPPER

These small (1/8 inch long), green, wedge-shaped insects frequently become abundant in alfalfa, but only occasionally cause damage. Normally, our harvesting practices interrupt the leafhopper life cycle in the egg stage. Damage, caused by injection of a toxin as the leafhopper feeds, is expressed as yellow or purplish triangular areas on individual leaflets. If leafhoppers are abundant, leaves may yellow and drop. In Nebraska, potato leafhoppers are usually more important on the second and third cuttings. Certain alfalfa varieties are somewhat resistant to potato leafhopper damage. The following treatment thresholds are recommended for potato leafhoppers:

Average height of alfalfa (inches)	Average number of potato leafhoppers per sweep of a 15" diameter insect net
0-3	0.2
3-6	0.5
6-12	1.0
12 or more*	1.5

*If alfalfa exceeds 12 inches or is approaching bloom, consider an early harvest rather than treatment.

REGISTERED FOR POTATO LEAFHOPPER CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.25 lb	Wait 3 days before harvest.
carbofuran <R> (Furadan 4F)	0.5-1.0 lb	Apply only to pure stands of alfalfa. Do not apply more than once per cutting or twice per year. Do not move bees into treated areas within 7 days of application. See label for harvest restrictions.
chlorpyrifos (Lorsban 4E)	0.5-1.0 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation systems.
diazinon AG500, 50W	0.375-0.5 lb	Wait 7 days before harvest.
dimethoate (Cygon 400)	0.25-0.5 lb	Wait 10 days before harvest. Only 1 application per cutting. Do not apply during bloom.
malathion 57EC	1.0-1.25 lb	No waiting period. During bloom, apply only in the evening or early morning.
methidathion <R> (Supracide 2E)	0.5-1.0 lb	Do not apply during bloom. Wait 10 days before harvest.

methyl parathion <R> (PennCap-M)	0.5-0.75 lb	Do not spray alfalfa during bloom to avoid injury to bees. Do not apply within 15 days of harvest or grazing. May be applied through sprinkler irrigation system.
<R> parathion 4EC, 8E, 8F	0.25-0.50 lb	Wait 15 days before harvest.
phosmet (Imidan 50WP)	1.0 lb	Only 1 application per cutting. Do not graze or harvest for 7 days.
trichlorfon (Dylox 80SP)	0.5-1.0 lb	Do not apply more than 3 times per cutting. No waiting period.

GRASSHOPPERS (NebGuide G86-791)

Grasshoppers are best controlled while they are small and are still feeding in field margins. If hoppers average 21-40 in field margins or 8-14 per square yard in the alfalfa, an insecticide application should be considered. Look for increased grasshopper activity in weedy alfalfa and fields surrounded by weedy waste areas or pasture. Hoppers become especially abundant during dry seasons. When establishing new stands of alfalfa, field margins should be treated before plants begin to emerge if grasshoppers are present at threshold levels. Refer to the following table for treatment thresholds for grasshoppers in crop fields and field margins.

TREATMENT THRESHOLDS FOR GRASSHOPPERS IN FIELDS AND FIELD MARGINS (NON-CROP AREAS)

Classification	Number of nymphs or adult hoppers per square yard		Treatment Necessary?
	Field	Field Margin	
Non-economic	0 to 2	5 to 10	Usually not
Light	3 to 7	11 to 20	Questionable (depends on hopper size, species and crop)
Moderate	8 to 14	21 to 40	Probably
Abundant	15 or more	41 or more	Yes

REGISTERED FOR CONTROL OF GRASSHOPPERS IN NON-CROP AND WASTE AREAS Rates are active ingredient per acre.

acephate (Orthene 75S)	0.125-0.5 lb
carbaryl (Sevin 4 Oil, XLR Plus, 80S)	0.5-1.5 lb
diazinon AG500	0.5-1.5 lb
<R> fenvalerate (Pydrin 2.4EC)	0.05-0.1 lb
malathion ULV 9.33	8-12 fl oz form.

If grasshoppers have already invaded the alfalfa, treatment is probably justified when there are 8-14 hoppers per square yard.

REGISTERED FOR GRASSHOPPER CONTROL IN ALFALFA

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, 4 Oil, XLR Plus)	0.5-1.5 lb	Wait 3 days before harvest.
carbofuran <R> (Furadan 4F)	0.125-0.25 lb	Apply only to pure stands of alfalfa. Do not apply more than once per cutting or twice per year. Do not move bees into treated areas within 7 days of application. See label for harvest restrictions.
chlorpyrifos (Lorsban 4E)	0.25-0.5 lb	Do not apply more than once per cutting or 4 times per year. See label for harvest restrictions. Some phytotoxic symptoms may occur on young plants. May be applied through overhead sprinkler irrigation system.
diazinon AG500, 50WP	0.5 lb	Wait 7 days before harvest.
dimethoate (Cygon 400)	0.25-0.50 lb	Wait 10 days before harvest. Only 1 application per cutting. Do not apply during bloom.
malathion 57EC	1.0-1.25 lb	No waiting period. During bloom, apply only in the evening or early morning.
malathion ULV 9.33	0.6 lb	No waiting period. Do not apply to alfalfa in bloom. Do not apply to seed alfalfa.

methyl parathion <R> (PennCap-M)	0.25-0.75 lb	Do not spray alfalfa during bloom to avoid injury to bees. Do not apply within 15 days of harvest or grazing. May be applied through sprinkler irrigation system. Use higher rates if majority of insects are large or weather is cool.
<R> parathion 4EC, 8E, 8F	0.25-0.50 lb	Wait 15 days before harvest.

BLISTER BEETLES

Blister beetles feed in clusters on the foliage and flowers of numerous weeds and agricultural crops. Several species are commonly found in alfalfa and can sometimes be abundant in July and August. Adults are elongate, cylindrical, soft-bodied beetles which range in length from 1/2 to 1 inch. Beetles are typically black, gray, spotted or striped with conspicuous heads and wing covers which do not cover the tip of the abdomen. When large aggregations of beetles are feeding, fields take on a ragged, stunted appearance. The larvae of some blister beetles are predaceous on the eggs of grasshoppers and in this capacity are considered beneficial. The body fluids of most blister beetles contain cantharidin, a chemical substance capable of producing large, watery blisters if allowed to come in contact with the skin. When consumed along with hay or feed, blister beetles can seriously affect livestock health. Horses are particularly sensitive with as few as 2-5 ingested beetles capable of causing colic, abortion, lethargy or urinary tract infections. Alfalfa producers should check fields for the presence of blister beetles before harvest, particularly for cuttings made late in the season. If beetles are numerous, an insecticide application should be considered. The use of a hay "crimper" may accentuate the problem if beetles are crushed along with the hay rather than moving out of the field following cutting.

SPRAYING BLOOMING ALFALFA CAN BE EXTREMELY HAZARDOUS TO BEES.
COORDINATE WITH LOCAL BEEKEEPERS BEFORE APPLYING INSECTICIDES.

REGISTERED FOR BLISTER BEETLE CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0 lb	Wait 3 days before harvest.
<R> parathion 4EC, 8E, 8F	0.25-0.5 lb	Wait 15 days before harvest.

SOYBEAN INSECTS (NebGuides G78-397 and G84-703)

Most insects that damage soybeans can be placed in two large groups, chewing insects and sucking insects. Chewing insects are the most common soybean pests and will damage cotyledons, leaf tissue, flowers and pods. Bean leaf beetles, green cloverworms, and grasshoppers are the most common chewing insect pests of soybeans in Nebraska. Other chewing insects include woollybear caterpillars, thistle caterpillars, soil cutworms, variegated cutworms, webworms, blister beetles, corn earworms, western bean cutworms, imported longhorn weevils, and Mexican bean beetles.

Sucking insects and mites injure soybean plants by probing leaves, stems, and pods with their piercing mouthparts. Damage is produced when plant juices are removed and/or toxins injected causing plants to appear stippled, mottled, and to take on a yellowed or discolored appearance. Stink bugs also occasionally attack pods and may discolor the developing bean within the pod. Thrips feed on leaves and flowers, but are not considered a serious threat. Spider mites may be of concern primarily during hot, dry years.

Do not use leftover soil insecticides on soybeans as crop damage may result. Certain organophosphates may produce a cross reaction and damage beans when planted in fields where herbicides such as Sencor and Lexone (metribuzin) were applied. Further, soybeans are unlikely to need protection from soil insects.

CHEWING INSECTS

Treatment decision-making

Treatment decisions are based on levels of defoliation, number of insects per unit area (usually number per foot of row), and stage of plant development. In general, soybeans can tolerate 30% defoliation before bloom and 20% defoliation between bloom and pod fill without significant yield loss. More detailed information can be found in NebGuide G78-397.

BEAN LEAF BEETLE

Beetles are about 1/4 inch long and vary in color from yellow to tan or red. They have a black outside border and usually 2 black spots on each wing cover. Bean leaf beetles feed on soybean cotyledons, leaves, and pods. Generally, soybeans are tolerant of leaf feeding.

Scout for bean leaf beetles in soybeans by examining 5 feet of row at each of 5 locations. Count the beetles and estimate damage at each site. Average the number of beetles at the sites to determine if control is justified.

TREATMENT THRESHOLDS FOR BEAN LEAF BEETLES

Growth Stage	Threshold
Seedling	At least 5 beetles present and 1 cotyledon destroyed per foot of row (newly emerged plants) or defoliation reaches 30 percent (plants with true leaves).
Post-seedling but before bloom	At least 10 beetles per foot of row and when defoliation reaches 30 percent.
Bloom to maturity	At least 10 beetles per foot of row and when defoliation reaches 20 percent or when pod damage reaches 5 to 10 percent.

SPRAYING FLOWERING SOYBEANS CAN BE EXTREMELY HAZARDOUS TO BEES.
COORDINATE WITH LOCAL BEEKEEPERS BEFORE APPLYING INSECTICIDES.

REGISTERED TO CONTROL BEAN LEAF BEETLE*
Rates are active ingredient per acre.

	acephate (Orthene 75S)	0.5-1.0 lb
<R>	azinphos methyl (Guthion 2S)	0.375-0.5 lb
	carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0 lb
	chlorpyrifos (Lorsban 4E)	0.5-1.0 lb
	dimethoate (Cygon 400)	0.5 lb
<R>	fenvalerate (Pydrin 2.4EC)	0.05-0.1 lb
<R>	methomyl (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.25-0.5 lb
<R>	methyl parathion (PennCap-M)	1.0 lb
<R>	permethrin (Ambush 2E, 25W, Pounce 3.2EC, 25WP)	0.05-0.1 lb
	thiodicarb (Larvin 3.2F)	0.45-0.75 lb
*See Restrictions and Comments on page 12.		

GREEN CLOVERWORM

Cloverworms are green caterpillars with a narrow, white stripe down each side. When fully grown, they are about 1 1/4 inch long. High numbers can strip soybeans of foliage. Control cloverworms when 12 or more half-grown (about 1/2 inch long) worms are found per foot of row and 20% defoliation occurs during bloom through pod fill.

REGISTERED FOR CONTROL OF GREEN CLOVERWORM*
Rates are active ingredient per acre.

	acephate (Orthene 75S)	0.5-1.0 lb
<R>	azinphos methyl (Guthion 2S)	0.375-0.5 lb
	Bacillus thuringiensis (Dipel, Thuricide)	See label
	carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0 lb
	chlorpyrifos (Lorsban 4E)	0.25-0.5 lb
<R>	fenvalerate (Pydrin 2.4EC)	0.05-0.1 lb
	malathion ULV 9.33	0.6 lb
	malathion 57EC	1.875 lb
<R>	methomyl (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.25-0.5 lb
<R>	methyl parathion (PennCap-M)	0.5 lb
<R>	parathion 4EC, 8EC	0.5 lb
<R>	permethrin (Ambush 2E & 25W, Pounce 3.2EC, 25WP)	0.05-0.1 lb
	thiodicarb (Larvin 3.2F)	0.25-0.4 lb
<R>	tralomethrin (Scout 0.3EC)	0.0141-0.0164 lb
*See Restrictions and Comments on page 12.		

GRASSHOPPERS
(NebGuide G86-791)

Grasshoppers breed in waste vegetation (roadsides, borrow pits, fence rows, benches, terraces) and pastures, moving into border rows of soybeans in July. Hoppers are best controlled in these staging areas before they invade soybeans. Plan to treat if hoppers average 20 or more per square yard in field margins or 15 or more per square yard in the soybean field itself. Refer to grasshopper recommendations in alfalfa section for a listing of insecticides registered for control of grasshoppers in non-crop and waste areas. The following are suggested for control of hoppers once they have moved into the field.

REGISTERED FOR CONTROL OF GRASSHOPPERS IN SOYBEANS*
Rates are active ingredient per acre.

	acephate (Orthene 75S)	0.25-0.5 lb
	carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0 lb
<R>	carbofuran (Furadan 4F)	0.125-0.25 lb
	chlorpyrifos (Lorsban 4E)	0.25-0.5 lb
	dimethoate (Cygon 400)	0.5 lb
<R>	fenvalerate (Pydrin 2.4EC)	0.1-0.2 lb
	malathion ULV 9.33	0.6 lb
<R>	methyl parathion (PennCap-M)	0.25-0.5 lb
<R>	parathion 4EC, 8EC	0.5 lb
<R>	tralomethrin (Scout 0.3EC)	0.015-0.0181 lb
*See Restrictions and Comments on page 12.		

SOIL CUTWORMS

Soil cutworms are occasional pests of seedling soybeans. The most common species involved is the black cutworm. This pest is a greasy black or gray worm with a brown head that may be over one inch long when feeding is completed. Black cutworms feed primarily at night and will hide under debris or in the soil during the day. Young black cutworms feed on above-ground portions of plants, while older cutworms cut plants at or below the soil surface. Controls are justified when 20 percent of the plants are cut, stands have gaps of one foot or more, and cutworms are present.

REGISTERED FOR CONTROL OF SOIL CUTWORMS*
Rates are active ingredient per acre.

	carbaryl (Sevin 80S, XLR Plus, Sevimol 4)	1.0 lb
	chlorpyrifos (Lorsban 4E)	0.5-1.0 lb
	(Lorsban 15G) (planting time)	8.0 oz/1,000 ft row
	diazinon AG500	(preplant only) 2.0-4.0 lb
<R>	fenvalerate (Pydrin 2.4EC)	0.1-0.2 lb
	thiodicarb (Larvin 3.2F)	0.5-0.75 lb
<R>	tralomethrin (Scout 0.3EC)	0.015-0.0181 lb
*See Restrictions and Comments on page 12.		

WIREWORMS AND SEED DAMAGING INSECTS*

Planter box seed treatments of diazinon and/or lindane.

*See Restrictions and Comments on page 12.

OTHER FOLIAGE FEEDERS

Treat when defoliation reaches 30% before bloom and 20% between bloom and pod fill.

SPRAYING FLOWERING SOYBEANS CAN BE EXTREMELY HAZARDOUS TO BEES.
COORDINATE WITH LOCAL BEEKEEPERS BEFORE APPLYING INSECTICIDES.

REGISTERED FOR CHEWING INSECTS ON SOYBEANS*

Insect	Insecticide	Rate AI/Acre
Blister Beetles	carbaryl (Sevin 80S, XLR Plus)	1.0 lb
Loopers	acephate (Orthene 75S)	0.5-1.0 lb
	Bacillus thuringiensis (Dipel, Thuricide)	See label
	<R> fenvalerate (Pydrin 2.4EC)	0.1-0.2 lb
	<R> methomyl (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.5-1.0 lb
	<R> permethrin (Ambush 2E, 25WP, Pounce 3.2EC)	0.05-0.10 lb
	thiodicarb (Larvin 3.2F)	0.5-0.75 lb
	<R> tralomethrin (Scout 0.3EC)	0.015-0.0181 lb
Saltmarsh Caterpillar	carbaryl (Sevin 80S, XLR Plus)	2.0 lb
	chlorpyrifos (Lorsban 4E)	0.50-1.0 lb
	<R> fenvalerate (Pydrin 2.4EC)	0.05-0.10 lb
	<R> methomyl (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.5 lb
Thistle Caterpillar	carbaryl (Sevin 80S, XLR Plus)	2.0 lb
Variegated Cutworm	carbaryl (Sevin 80S, XLR Plus)	1.0 lb
	chlorpyrifos (Lorsban 4E)	0.5-1.0 lb
	<R> fenvalerate (Pydrin 2.4EC)	0.1-0.2 lb
	thiodicarb (Larvin 3.2F)	0.5-0.75 lb
	<R> tralomethrin (Scout 0.3EC)	0.015-0.0181 lb
Webworms	carbaryl (Sevin 80S, XLR Plus)	1.0 lb
	<R> methyl parathion (PennCap-M)	0.25 lb
	<R> parathion 4EC, 8EC	0.25 lb

Woollybear Caterpillar	chlorpyrifos (Lorsban 4E)	0.50-1.0 lb
	<R> fenvalerate (Pydrin 2.4EC)	0.05-0.10 lb
	<R> permethrin (Ambush 2E, Pounce 3.2EC)	0.05-0.10 lb
	thiodicarb (Larvin 3.2F)	0.25-0.4 lb

*See Restrictions and Comments below.

SPIDER MITES

Control if mites are abundant on undersides of leaves and lower leaves are starting to drop.

REGISTERED FOR CONTROL OF SPIDER MITES*

Rates are active ingredient per acre.

chlorpyrifos (Lorsban 4E)	0.25-0.5 lb
dimethoate (Cygon 400)	0.5 lb

*See Restrictions and Comments below.

SOYBEAN INSECTICIDE RESTRICTIONS

In some cases, harvest intervals and other label restrictions prevent the use of certain pesticides even if registered for the problem. When choosing an insecticide, be certain that the use of the product does not result in illegal application; danger to the user, environment, or beneficial organisms; or residues that exceed tolerances. Use the following table as a guide when choosing a chemical control.

Insecticide	Restrictions and Comments
acephate (Orthene 75S)	Apply in 10 to 50 gallons of water with ground equipment or in 2 to 10 gallons with aerial equipment. Apply as needed but do not apply within 14 days of harvest. Do not graze or cut vines for hay or forage.
azinphos methyl <R> (Guthion 2S)	Apply as necessary. Do not apply within 45 days of harvest. Do not graze or feed treated vines to livestock.
Bacillus thuringiensis (Dipel, Thuricide)	No waiting period.
carbaryl (Sevin 80S, Sevimol 4, XLR Plus)	No waiting period. May be applied through overhead sprinkler irrigation system.
carbofuran <R> (Furadan 4F)	Do not apply within 21 days of harvest. Do not make more than two foliar applications per season. Do not graze or feed foliar treated forage to livestock or cut for hay or silage. Do not use Furadan 4F as a foliar application if Furadan 10G, 15G or 4F was applied to beans at planting time.
chlorpyrifos (Lorsban 4E)	For soil treatment, (except pivot application) use at least 10 gallons of spray per acre. For foliar treatment, apply in sufficient water to ensure thorough coverage. Do not apply more than 3 lbs AI (6 pints formulation) per acre per season. Do not apply last treatment within 28 days before harvest nor apply last two treatments closer than 14 days apart. Do not allow livestock to graze in treated areas nor otherwise feed treated soybean forage to meat or dairy animals within 14 days after application. Do not feed straw from treated soybeans to meat or dairy animals within 28 days after application. On determinate soybeans do not apply more than one application after pod set. May be applied through overhead sprinkler irrigation system.
(Lorsban 15G)	Do not make more than one application per season.
diazinon AG500	Apply a minimum of 5 gallons of water per acre for ground applications. No harvest restrictions.
dimethoate (Cygon 400)	Do not apply within 21 days of harvest. Do not feed or graze within 5 days of last application.
fenvalerate <R> (Pydrin 2.4EC)	Do not feed or graze livestock on treated plants. Do not exceed 0.8 lb (42.64 fl oz formulation) per acre per season.
malathion 57EC	No waiting period.
malathion ULV 9.33	Do not apply within 7 days of harvest or grazing.
methomyl <R> (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	Do not apply within 14 days of harvest. If applied at under 0.5 lb AI, 3 days forage and 7 days hay; if applied at 0.5 lb AI or over, 10 days forage and 12 days hay.

methyl parathion <R> (PennCap-M)	Do not apply within 20 days of harvest or grazing. Do not apply more than 2 times per season. May be applied through overhead sprinkler irrigation system.
<R> parathion 4EC, 8EC	Do not apply within 20 days of harvest. Do not apply more than twice per growing season.
permethrin <R> (Pounce 3.2EC, 25WP)	Apply a minimum of 1 gallon of finished spray per acre by aircraft and 5 gallons with ground equipment. Do not make more than 2 applications per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage. Do not plant rotational crops within 60 days of last application.
<R> (Ambush 2E, 25WP)	Apply by air or ground. Do not make more than 2 applications per season. Can be applied up to 60 days before harvest. Do not graze treated areas or harvest for forage or hay.
thiodicarb (Larvin 3.2F)	Do not feed forage, hay, or straw to livestock. Do not apply less than 28 days before harvest.
tralomethrin <R> (Scout 0.3EC)	Do not graze or harvest treated soybean forage, straw or hay for livestock feed. Do not apply more than 0.12 AI/A in one growing season. Do not apply within 21 days of harvest.

WHEAT INSECTS

Wheat can be damaged by various insect pests. The principal insects that attack the wheat plant are the pale western and army cutworms, armyworms, grasshoppers, greenbugs and other aphids, the Hessian fly, and various seed feeding insects such as true and false wireworms. Much of the loss to the wheat crop from insect pests can be prevented by proper cropping practices. Delayed seeding, for example, is an important control measure for the Hessian fly and wheat curl mite. Timeliness of application is important when chemicals are used for insect control. Check appropriate NebGuides for additional information.

WIREWORMS AND OTHER SEED DESTROYING INSECTS

Drill box seed treatment with lindane and/or diazinon. Follow label directions for amounts and restrictions.

PALE WESTERN AND ARMY CUTWORMS (NebGuide G74-130)

Sporadic outbreaks of pale western and army cutworms have occurred in Nebraska following periods of drought. Thin stands on lighter soils are more subject to attack. The pale western cutworm is gray, about 1 inch long and feeds beneath the soil surface, destroying plants and causing serious damage. One pale western cutworm per foot of drill row is considered an economic level. The army cutworm is mottled brown, up to 1 1/4 inches long and "grazes" the wheat above ground. Army cutworms are not considered serious pests unless conditions are dry and worms average 2-3 per foot of row. Spray early morning or evening for best results.

REGISTERED FOR PALE WESTERN AND ARMY CUTWORM CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
endosulfan (Thiodan 3EC)	0.5 lb	Do not apply after heads begin to form. Do not feed treated forage to livestock. Army cutworm only.
<R> parathion 4EC, 8EC	0.5 lb	Wait 15 days before harvest.
trichlorfon (Dylox 80SP)	1.0 lb	Wait 21 days before harvest.

ARMYWORMS (True, Fall, Wheathead, Yellowstriped) (NebGuide G82-615)

Four species of armyworm occur in Nebraska. They are generally associated with field crops and small grains but also feed on pasture grasses. These pests hide in the soil or beneath debris by day, emerging at night to feed on foliage. Spray in early morning or evening for best results.

REGISTERED FOR ARMYWORM CONTROL

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 80S, XLR Plus, 4 Oil)	1.25-1.78 lb	No waiting period--forage. 21 days for grain.

methomyl <R> (Lannate 1.8L, 90SP, Nudrin 1.8L, 90SP)	0.225-0.45 lb	Wait 7 days before harvest, 10 days before grazing.
methyl parathion <R> (PennCap-M)	0.5-0.75 lb	Wait 15 days before harvest or grazing.
<R> parathion 4EC, 8EC	0.5 lb	Wait 15 days before harvest.
trichlorfon (Dylox 80SP)	1.0 lb	Wait 21 days before harvest.

GRASSHOPPERS (NebGuide G86-791)

Prevent damage to wheat by controlling grasshoppers when there are 20 or more adults per square yard in field margins, or 8 or more in the field itself. Time of day, temperature, wind, plant density and height of vegetation affect grasshopper activity and should be considered in making counts. For grasshopper control recommendations in non-crop and waste areas, refer to page 8.

REGISTERED FOR GRASSHOPPER CONTROL IN WHEAT

Insecticide	Rate AI/Acre	Restrictions and Comments
carbaryl (Sevin 5% bait)	20-40 lb form.	No waiting period.
(Sevin 80S, Sevimol 4, XLR Plus)	0.5-1.5 lb	No waiting period.
carbofuran <R> (Furadan 4F)	0.25 lb 0.25-0.5 fl oz/ 1,000 lin. ft	See label. Apply in seed furrow using microtube applicator.
dimethoate (Cygon 400)	0.375 lb	Wait 14 days for grazing, 60 days for harvest. Maximum of 2 applications per season.
malathion 57EC	1.0 lb	Wait 7 days before harvest.
malathion ULV 9.33	0.6 lb	Wait 7 days before harvest or forage use.
methyl parathion <R> (PennCap-M)	0.25-0.75 lb	Wait 15 days before harvest or grazing.
<R> parathion 4EC, 8EC	0.5 lb	Wait 15 days before harvest.

NOTE: Thimet 20G at a rate of 1.2 oz per 1,000 feet (minimum 8-inch row spacing) and Di-Syston 15G* (disulfoton) at a rate of 1.7 oz per 1,000 feet can be used in the seed furrow at planting time. Only the border 30-40 feet normally benefit from treatment. Wait at least 45 days before grazing after applying Thimet 20G and 75 days after applying Di-Syston 15G*. Do not make any later applications after planting time treatment. Do not mix granules with wheat seed in drill box.

*Do not plant any food or feed crop in rotation after a field treatment with disulfoton (Di-Syston 15G) unless it is a registered use for disulfoton.

GREENBUGS AND OTHER APHIDS

Wheat infested by greenbugs develops yellowed or dead areas during late fall or early spring. Examination of these areas should reveal the small green aphids feeding on the undersides of leaves.

The Russian Wheat Aphid (RWA) is a new and serious threat to wheat production. Begin to watch fields in March and continue through heading. Like the greenbug, it injects a potent salivary toxin into plants, causing leaves to roll and turn purple with yellow or white lengthwise stripes. Heavily infested plants appear flattened, with young tillers lying parallel to the ground. RWA is difficult to control due to its habit of feeding in rolled up leaves of new growth. Feeding by RWA on flag leaves can cause premature death and distorted white heads. Damage often appears in circular, purple or reddish-brown areas in a field. The RWA is a small aphid, with a greenish spindle-shaped body and short antennae. Its "double-tailed" appearance and reduced cornicles ("tailpipes") help to distinguish it from greenbugs and other aphids commonly found in wheat. Contact your local Cooperative Extension Service Office for help in distinguishing these aphid species. Heavy infestations of RWA can reduce yields by 50% or more. While no insecticides currently are registered for this new pest, most of the following could be used under their "aphid" labeling. CAUTION: Several other species of aphids occur in wheat -- the greenbug, oat-bird cherry aphid, English grain aphid and corn leaf aphid. The last three are rarely damaging. Therefore identification is critical. Don't make a control decision until you are sure which aphids you have in your fields.

Table I. Treatment thresholds for greenbugs at various growth stages of wheat. Kansas Cooperative Extension Service.

Stage and Development of Plants	Number of greenbugs per linear foot of drill row
Seedlings, thin stands, less than 3 tillers	50 (fewer if plants are very small)
3 to 6 inch wheat, 3 tillers or more	100 - 300
6 to 10 inch wheat	300 - 500

REGISTERED FOR CONTROL OF GREENBUGS AND OTHER APHIDS

Insecticide	Rate AI/Acre	Restrictions and Comments
dimethoate (Cygon 400)	0.25-0.375	Wait 60 days before harvest, 14 days to graze. Maximum 2 applications per season.
disulfoton <R> (Di-Syston 8EC)*	0.5 lb	Wait 21 days before harvest. Do not graze.
<R> (Di-Syston 15G)*	1.0 lb	Planting time application. Wait 30 days to graze.
malathion 57EC	1.0 lb	Wait 7 days before harvest or forage use.
methyl parathion <R> (PennCap-M)	0.25-0.5 lb	Wait 15 days before harvest or grazing.
<R> parathion 4EC, 8EC	0.5 lb	Wait 15 days before harvest.
phorate <R> (Thimet 20G)	1.0 lb	Apply in seed furrow at planting. Wait 45 days to graze.

*Do not plant any food or feed crop in rotation after a field treatment with disulfoton unless it is a registered use for disulfoton.

HESSIAN FLY (NebGuide G73-46)

Injury caused by Hessian fly is not conspicuous at first, but can be devastating. Hessian fly numbers appear to be increasing in Nebraska perhaps due to recent increased rainfall. Wheat infested in the fall is stunted. Leaves of plants take on a dark bluish-green color, are distinctly thickened and stand more erect than those of uninfested plants. Control consists of cultural methods. In eastern and central Nebraska, delay planting until after the summer generation of flies has died to reduce fall infestations. Fly-free dates for planting wheat in your area can be obtained from your local Extension office. In addition, destroying wheat stubble will help reduce the fly population. However, in western Nebraska, because of wind erosion, this practice may not always be practical. The planting of fly-resistant wheat varieties is also important in reducing the overall fly infestation. Two systemic insecticides, disulfoton (<R> Di-Syston 15G*) and phorate (<R> Thimet 20G) are registered for Hessian fly control, but the economics and effectiveness of preventive treatments in Nebraska have not been demonstrated.

*Do not plant any food or feed crop in rotation after a field treatment with disulfoton unless it is a registered use for disulfoton.

CHINCH BUGS (NebGuide G86-806)

Adults leave overwintering sites and migrate to small grains when temperatures reach 70° F or above. Heavy spring infestations in wheat may reduce yields somewhat, particularly if wheat is under stress from other causes. Usually either a partial or full generation occurs in wheat fields prior to migration to corn or sorghum. Controls are seldom necessary, but 1/2 pint per acre of <R> parathion applied aerially can be used at least 15 days prior to harvest. Damage to fall-sown wheat seldom occurs because chinch bugs are then moving to overwintering sites.

WHITE GRUBS

White grubs occasionally damage fall-sown wheat in Nebraska. In 1986, some reports were received from Chase, Perkins and Keith counties in southwestern Nebraska. Oklahoma entomologists suggest treatment when grubs average 3-4 per square foot. Treatment must be applied at or before planting. Once plants have emerged from the soil, little can be done. Phorate (<R> Thimet 20G) is federally registered as a planting time treatment, applied in-furrow at a rate of 1.2 oz formulation per 1,000 feet of drill row (minimum 8 inch row spacings). Diazinon 14G and AG500 are both registered for preplant treatment at a rate of 1-2 lbs AI/A disked into the upper 2-4 inches of soil.

RANGE AND PASTURE INSECTS

Range and pastures in Nebraska are attacked by several insect pests, with grasshoppers being the most important. Grasshoppers usually affect large areas while areas damaged by other

species are usually much smaller. It may not be economical to spray large areas of range and pasture without supplemental financial assistance. Small infested areas may be sprayed to control the insects and prevent their spread. Leafhoppers, western harvester ants, blister beetles, and army cutworms may also cause injury to range and pasture grass. However, there are no insecticides currently registered for control of these insects on range or pasture.

RECOMMENDED FOR CONTROL OF RANGE AND PASTURE INSECTS

Insect	Insecticide	Rate	Restrictions and Comments
Grasshoppers	acephate	0.094-0.125 lb	Minimum of 1/2 gallon of solution by (Orthene 75S) air or 10 to 20 gallons per acre by ground. Do not apply when lactating dairy cattle are present. Do not pasture or feed to lactating dairy cattle within 21 days of after application. Do not remove treated hay within 21 days after application. Remove meat animals from treated areas at least one day prior to slaughter if they were present at application or grazed treated areas within 21 days after application. Do not make more than one application per season.
	carbaryl (Sevin XLR Plus, 80S, 4 Oil, 20% Bait)	0.5-1.5 lb	No restrictions.
	diazinon AG500	0.375-0.5 lb	Apply in water or oil (minimum one gallon of oil per acre). Dairy and beef cattle and sheep (1) may be grazed immediately following application, (2) may be fed green forage immediately following cutting, and (3) may be fed hay provided grass is not cut for hay for 21 days if the spray is a water solution, or for 30 days if the spray is an oil solution. Do not spray livestock directly with either air or ground equipment. Do not repeat application for 30 days.
	<R> ethyl parathion 4EC, 8EC	0.5-0.75 lb	Do not apply within 15 days of harvest or grazing.
	malathion 57EC	1-1.5 lb	May be applied on the day of harvest or grazing.
	malathion ULV 9.33	8-12 oz form.	May be applied on the day of harvest or grazing.
	<R> methyl parathion (PennCap-M)	0.25-0.5 lb	Do not apply within 15 days of harvest or grazing.
	naled (Dibrom 8EC)	0.5-0.75 lb	See label for restrictions.
	Armyworms carbaryl (Sevimol 4, 80S, XLR Plus)	1-1.5 lb	No restrictions.
	<R> ethyl parathion 4EC, 8EC	0.5-0.75 lb	Do not apply within 15 days of harvest or grazing.
Black Grass Bugs (Labops sp.)	malathion 57EC	1.5 lb	0 days to harvest.
	<R> methyl parathion (PennCap-M)	0.25-0.5 lb	Do not apply within 15 days of harvest
	carbaryl (Sevin XLR Plus, 80S, Sevimol 4)	1-1.5 lb	No restrictions.
	<R> ethyl parathion 4EC, 8EC	0.5-0.75 lb	Do not apply within 15 days of harvest or grazing.
	malathion ULV 9.33	8-12 oz form.	May be applied on the day of harvest or grazing.