

# Beef palatability and its relationship with protein degradation and muscle fibre type profile in *longissimus thoracis* in Alentejana breed from divergent growth pathways

P. Costa<sup>1†</sup>, J. A. Simões<sup>2</sup>, S. P. Alves<sup>1</sup>, J. P. C. Lemos<sup>1</sup>, C. M. Alfaia<sup>1</sup>, P. A. Lopes<sup>1</sup>, J. A. M. Prates<sup>1</sup>, J. F. Hocquette<sup>3,4</sup>, C. R. Calkins<sup>5</sup>, V. Vleck<sup>6</sup> and R. J. B. Bessa<sup>1,2</sup>

<sup>1</sup>CIISA – Faculdade de Medicina Veterinária, Universidade de Lisboa, Avenida da Universidade Técnica, Pólo Universitário do Alto da Ajuda, 1300-477 Lisboa, Portugal; <sup>2</sup>L-INIAV Santarém, Instituto Nacional dos Recursos Biológicos, Quinta da Fonte Boa, 2005-048 Vale de Santarém, Portugal; <sup>3</sup>INRA, UR 1213, Unité de Recherches sur les Herbivores (URH), Theix, F-63122 Saint-Genès Champanelle, France; <sup>4</sup>Clermont Université, VetAgro Sup, UMR1213, Herbivores, BP 10448, F-63000 Clermont-Ferrand, France; <sup>5</sup>Department of Animal Science, University of Nebraska, Lincoln, NE 68583-0908, USA; <sup>6</sup>CIPER – Faculdade de Móticidade Humana, Universidade de Lisboa, Estrada da costa, Cruz Quebrada-Dafundo, 1499-002 Lisboa, Portugal

(Received 1 March 2016; Accepted 3 June 2016; First published online 5 July 2016)

*The traditional beef production in the South of Portugal is based on a discontinuous growth (DG) system that requires lower external inputs and could enhance meat quality and financial returns to cattle producers. This system allows farmers to take advantage of the bull's compensatory growth when the pasture is abundant and finishes the cattle on concentrates for 2 to 3 months before slaughter. The fast gain rate before slaughter could be a valuable strategy to improve tenderness and to reduce its inconsistency in beef production. Therefore, the aim of this study was to evaluate the effects of production system (continuous growth (CG) v. DG) on longissimus thoracis muscle properties from Alentejana bulls. In total, 40 Alentejana male calves were allocated to two distinct feeding regimes: in the CG system, animals were fed concentrate plus hay and slaughtered at 18 months of age, whereas in the DG system, animals were fed on hay until 15 months of age and then fed the same diet provided to the CG group until 24 months of age. The DG system had a positive impact on meat tenderness ( $P < 0.001$ ) and global acceptability ( $P < 0.001$ ). DG bulls had greater fibre cross-sectional area (CSA) of glycolytic fibres ( $P < 0.05$ ) and relative area of the muscle (RA) occupied by type IIX fibres ( $P < 0.01$ ) and greater levels of  $\alpha$ -actinin ( $P < 0.05$ ) and myosin light chain 2 ( $P < 0.01$ ) proteins, and  $\text{pH}_{24\text{h}}$  ( $P < 0.01$ ) than CG bulls. The latter had greater CSA of type I ( $P < 0.05$ ) and type IIA ( $P < 0.01$ ) and greater RA of type IIA ( $P < 0.05$ ) and oxidative ( $P < 0.05$ ) than CG bulls. The compensatory growth production system had a positive impact on meat tenderness and global acceptability, overcoming the negative effects of slaughter of the bulls at a later age. The DG beef system could be a worthwhile strategy of beef production in Mediterranean areas due to the low-quality pasture in summer.*

**Keywords:** bulls, compensatory growth, meat sensory characteristics, muscle fibre traits, muscle protein fragments

## Implications

The traditional Mediterranean beef is based on a discontinuous growth (DG) production system. This requires both lower external input and provides an enhanced financial return to cattle producers than continuous systems, and contributes to environmental sustainability. This study shows that DG production could be a valuable strategy for the improvement of the palatability of Portuguese beef. The increased growth rates that are obtained before slaughter in DG animals could overcome the negative effects of age on tenderness and other markers of consumer acceptability.

## Introduction

Traditionally, beef production in the South of Portugal is based on a DG production system, whereby the calves that are born in the summer are weaned at the beginning of, and then spend all of, spring grazing on pasture. Afterwards, the cattle are maintained on low-quality pasture, supplemented with hay in order to ensure at least the weight maintenance until the following spring. The farmers thus take advantage of the bull's compensatory growth phase during its second spring when pasture is abundant. Finally, the cattle are fed on concentrate for 2 to 3 months before slaughter at 24 months of age. This system requires lower external input and could enhance both meat quality and thus cattle producers' financial return. However, some beef producers are

<sup>†</sup> E-mail: paulocosta@fmv.utl.pt

reluctant to use this system because, in order to obtain commercially acceptable carcasses, animals have to be slaughtered 6 months later than those produced via more intensive systems.

It is widely accepted that meat quality varies with fibre type profile (Lefaucheur, 2010). This is mainly due to indirect effects of muscle fibre type on muscle components such as sarcoplasmic proteins, muscle enzymes, intramuscular fat (IMF) levels and connective tissues (Lefaucheur, 2010). Beef palatability is determined by tenderness, juiciness and flavour (Calkins and Hodgen, 2007; Hocquette *et al.*, 2014). Meat final tenderness depends mainly on the *postmortem* changes affecting the muscle contractile system. The increase in meat tenderness that results from ageing is a consequence of proteolytic degradation of structural (cytoskeletal) components of muscle by endogenous proteases which depend on fibre profile. The ageing rate of glycolytic fibres is generally faster than that of oxidative fibres (Ouali and Talmant, 1990; Whipple and Koohmaraie, 1992; Lefaucheur, 2010).

As the proteolytic enzymes in the living muscles have a decisive role in *postmortem* protein degradation and meat tenderness (Fishell *et al.*, 1985), it is expected that a restrictive/re-feeding strategy before slaughter would lead to greater *postmortem* proteolysis and improved ultimate tenderness (Therkildsen, 2005). The ensuing high energy intake during re-feeding leads to an increase in protein turnover, with the increase in protein synthesis larger than protein degradation leading to improved meat tenderness. In a previous study from our research team, it was found that Alentejana bulls produced on DG had consistently low Warner-Bratzler shear force (WBSF) values than their counterparts produced on continuous growth (CG). Therefore, the practice of employing fast gain rates before slaughter, as in a DG production system, could be a valuable strategy to improve tenderness and to reduce its inconsistency in beef meat production. Thus, the effect of growth path on muscle fibre composition and protein degradation and how these effects relate to beef palatability in *longissimus thoracis* (Lt) of Alentejana bulls will be evaluated in the present study.

## Material and methods

### *Animals and meat samples*

This experiment is fully described in Costa *et al.* (2015). In brief, 40 purebred Alentejana male calves ( $9.0 \pm 0.46$  (mean  $\pm$  SD) months of age,  $239 \pm 45$  kg live weight) were randomly allocated to each of the two distinct feeding regimes. In the CG production system, the animals were fed *ad libitum* on concentrates plus hay throughout the trial and slaughtered at 18 months of age. In the DG production system, animals were fed *ad libitum* on hay only from 9 to 15 months of age. They were then fed the same diet provided to the CG group (concentrates plus hay) until 24 months of age. After being stunned with a captive bolt, the cattle were dressed with carcasses suspended by the Achilles tendon. The carcasses were split

along the spine and the kidney knob and channel fat from both sides were removed. During the first hour *postmortem*, biopsies of about  $1 \text{ cm}^3$ , for the purpose of histological analysis, were obtained from the superficial/middle layer of the Lt muscle at the level of the 9th vertebrae and on the right side of the carcass. The samples were immediately frozen by immersion in isopentane cooled by liquid nitrogen, and were kept at  $-80^\circ\text{C}$  until analysis.

After 24 h chilling at  $6^\circ\text{C}$ , muscle pH was assessed using a Hanna Instruments Hi 9023 device (Póvoa do Varzim, Portugal). The carcasses were then stored at  $4^\circ\text{C}$  until 7 days *postmortem* (cold carcass). They were then jointed and two steaks of 2.5 cm thickness were taken from the space between the 8th and 12th vertebrae of the Lt muscle. The steaks were vacuum packed and frozen at  $-30^\circ\text{C}$  until subsequent analysis.

### *Muscle histology*

Serial transverse muscle sections ( $10 \mu\text{m}$  thick) were cut in a cryostat at  $-24^\circ\text{C}$  and stained for myosin adenosine triphosphatase (ATPase) at pH 4.45 (Brooke and Kaiser, 1970). Fibres were classified as type I, type IIA and type IIX. It has been reported that myosin heavy chain-IIb (MHC-IIb) is not expressed in cattle limb and trunk muscles (Maccatrozzo *et al.*, 2004; Toniolo *et al.*, 2005). However, Picard and Cassar-Malek (2009) found the MHC-IIb isoform in *semi-tendinosus* and Lt muscles of the Blonde d'Aquitaine breed. It is possible that some of the muscle fibres classified here as type IIX are in fact IIb.

The succinate dehydrogenase (SDH) protocol of Sheehan and Hrapchak (1987) was used to classify fibres as oxidative or glycolytic. The combination of ATPase and SDH classifications in serial cuts has previously been used to distinguish between slow oxidative, fast oxidative-glycolytic and fast glycolytic types (Picard *et al.*, 1998; Oury *et al.*, 2010). Here, the ATPase and SDH classifications were performed separately. The percentage of each fibre and the mean surface area for 200 fibres were obtained from two randomly selected areas of the serial sections using an image analysis software program (version 10.5.0, VectorWorks, Nemetschek, 2003). Type IIC fibres were also identified by staining at pH 4.2 (Picard *et al.*, 1998) but are not included in the 'Results' section of this paper because they accounted for  $<5\%$  of the total fibre population of each section.

### *Muscle protein fragments*

Myofibrils from Lt muscle at 7 days *postmortem* were obtained according to the method of Parrish *et al.* (1973), using 50 mM Tris, 5 mM ethylenediaminetetraacetic acid (EDTA) and 100 mM KCl, pH 7.6 buffer as the isolation medium. The final residue was suspended in 100 mM KCl, 1 mM  $\text{NaN}_3$ . The sample suspension was then adjusted to 2 mg protein/ml with the same KCl- $\text{NaN}_3$  solution and stored at  $-20^\circ\text{C}$  until analysis. Myofibrillar proteins from the suspensions were precipitated with acetone and evaporated to dryness (Savant™ SC210 P1 SpeedVac-Thermo Fisher Scientific, Waltham, MA, USA). Dried samples were dissolved

in the following buffers: 5 mM Tris, 0.5 mM EDTA, 1.25% SDS, 1% of 2-mercaptoethanol, 0.1% urea, pH 8.0 and 8 M urea, 2 M thiourea, 3% SDS, 75 mM dithiothreitol (DTT) and 25 mM Tris, pH 6.8 (Fritz and Greaser, 1991) in a 1 : 1 proportion. The samples were heated at 50°C for 20 min before electrophoresis. Horizontal electrophoresis was performed using PhastSystem equipment (Amersham Biosciences; Buckinghamshire, UK) using a PhastGel Homogeneous 12.5% and a PhastGel Gradient 4% to 15%. The samples (5 µg) were run in the stacking gel at 1.0 mA for 1 V h and in the separation gel at 3.0 mA, 80 V h at 15°C. Gels were stained with Coomassie Blue R-350 and molecular weights of the proteins were estimated by running standard proteins of known weight in each gel. The quantification of major proteins and the identification of their respective bands separated in each gel were performed using a Biotec-Fischer HI-CAM densitometer (Reiskirchen, Germany).

#### *Trained sensory panel analysis*

The meat trained panel that was used for this study had a minimum of 3 years experience in performing sensory evaluation of meat. The panellists were recruited from 80 FMV/CIISA employees and graduate students who had previously been selected and trained according to Cross *et al.* (1978), Meilgaard *et al.* (2006) and American Meat Science Association (2015). In brief, the candidates were first pre-screened to determine their level of interest, availability, dependability, health (including dentures, allergies, use of medication), work experience, gender, age, smoking and food preferences. Those who passed the pre-screening stage were then invited to participate in several screening exercises with the objective of selection of those having interest and motivation in sensory evaluation, normal sensory acuity and ability to discriminate and reproduce results. In all, 26 candidates were selected for training to improve their ability to categorise and recall sensory information accruing from meat tasting sessions.

For the present study, five males and seven female panellists (average  $\pm$  SD age  $38.9 \pm 7.76$  years) were recruited. For each session of sensory analysis, 10 frozen steaks were thawed at 4°C for 24 h and cooked in a plate grill (65/70 FTES Electric Griddle; Modular Catering Equipment, Treviso, Italy) at 250°C, until they reached an internal temperature of 70°C, which was monitored by an internal thermocouple (Lufft C120; München, Germany). Cooking losses (CL) were determined by calculating the difference in weight before and after thermal processing. The steaks were trimmed of any external connective tissue, cut into  $\sim 2 \times 2 \times 2$  cm<sup>3</sup> samples and maintained at 60°C until tasting. The tasting involved two cores of each sample being tasted, as soon as possible, by each panellist. The sensory analysis was performed in four sessions (eight to nine samples per session).

The beef attributes evaluated were as follows: tenderness (defined as the opposite of the force required to bite through the sample with the molars), juiciness (the amount of liquid drained from the sample during the two initial chews),

off-flavour (the intensity with which an undesirable flavour, not associated with beef, is recognised), flavour (the intensity with which the beef sample is recognised as distinctly bovine rather than deriving from any other species) and global acceptability (the perception of palatability, taking into account the aforementioned attributes). The scale applied in the sensory analysis was structured into 8 points, with 1 being extremely tough, dry, weak and less desirable and 8 being extremely tender, juicy, strong and more desirable for tenderness, juiciness, flavor and overall acceptability, respectively. The off-flavour was structured into nine points, representing 0 (absence) and 8 (extremely intense).

#### *Shear force measurements*

After thawing for 24 h at 4°C, the steaks were grilled to an internal temperature of 70°C (model HD8704; Delta OHM, Caselle di Selvazzano, Italy). Then, after cooling for 1 h, the steaks were cut, in the direction of the fibres, into eight to 10 cores each with cross-sectional area (CSA) of about 1 cm<sup>2</sup> and a length of 2 cm. Shear force expressed in kg was measured in a texturometer (TA-tx2i Texture Analyser; Stable Micro Systems, Surrey, UK) equipped with a Warner-Bratzler shear device, using specific software (Texture Expert Exceed; Stable Micro Systems).

#### *Intramuscular fat content and myofibril fragmentation index*

The IMF content of fresh Lt muscle samples was determined by hydrolysis with 4 M HCl followed by Soxhlet extraction with petroleum ether for 6 h (Association of Official Analytical Chemists, 2000). The myofibril fragmentation index (MFI) was determined for frozen Lt muscle samples 7 days *postmortem*, according to Culler *et al.* (1978).

#### *Statistical analysis*

Sensory scores, meat traits, muscle protein fragments and muscle fibre profile were analysed using the Proc Mixed procedure of SAS software, version 9.2 (SAS Institute, Cary, NC, USA), considering the animal as the experimental unit. The model included the fixed effect of production system (CG *v.* DG). All data were reported as mean  $\pm$  standard error. Differences between groups were examined for statistical significance at the 95% confidence level using the PDIF option. Pearson's correlation coefficients between sensory panel scores, fibre profile and meat quality traits were calculated using STATISTICA (StatSoft Inc., OK, USA, 2004) software. In order to explore the relationship between meat traits, a principal component analysis (PCA) was carried out using STATISTICA software, with the variables for PCA being standardised to a mean of 0 and variance of 1. The use of multivariate statistical methods, such as PCA for complex samples as meat, makes it possible to identify the most important directions of variability in a multivariate data matrix and to present the results in a graphical plot. The PCA transforms the original variables into new axes, or principal components (PCs) that are orthogonal, so that the data presented in those axes are uncorrelated with each other. PCA therefore expresses as much as possible of the total

variation in the data in only a few PCs and each successively derived PC expresses decreasing amounts of the variation (Destefanis *et al.*, 2000).

## Results

### Muscle fibre characteristics

The production system was not a source of variation in muscle fibre type distribution (Table 1). In contrast, the fibre CSA was 575  $\mu\text{m}^2$  and around 1100  $\mu\text{m}^2$  greater in CG for type I ( $P = 0.01$ ) and type IIA ( $P < 0.01$ ), respectively, than in DG bulls. Glycolytic fibre CSA was about 1100  $\mu\text{m}^2$  greater ( $P = 0.02$ ) in DG than in CG bulls. The CG bulls also had a greater proportion of type IIA ( $P = 0.01$ ) and oxidative fibres ( $P = 0.02$ ) than DG. The relative muscle area (RA) occupied by IIX fibres was greater ( $P < 0.01$ ) in DG than in CG.

### Muscle protein fragments

Details of the Lt muscle protein fragments at 7 days *post-mortem* are presented in Table 2. DG bull meat had greater levels of  $\alpha$ -actinin ( $P = 0.04$ ) and myosin light chain 2 (MLC2) ( $P = 0.001$ ) than bulls on the CG system. The production system was not a significant source of variation for the remaining protein fragments ( $P > 0.05$ ). However, CG bulls had a tendency to present greater kDa70 ( $P < 0.08$ ) and less troponin T ( $P < 0.07$ ) levels than DG bulls.

### Sensory panel scores and meat quality traits

Sensory panel scores for, and meat quality characteristics of, the Lt muscle are presented in Table 3. Tenderness ( $P < 0.001$ ) and global acceptance ratings ( $P < 0.001$ ) were 0.5 points greater in meat from DG than CG bulls. As compared with CG, a tendency ( $P < 0.06$ ) to greater juiciness and flavour was observed in DG cattle. With respect to

**Table 1** Muscle fibre profile of longissimus thoracis from Alentejana bulls produced from to continuous (CG) and discontinuous growth (DG) production systems

	CG (n = 17)	DG (n = 17)	SEM	P-value
Fibre type distribution (%)				
I	31.9	30.8	1.4	0.58
IIA	30.7	28.8	1.2	0.27
IIX	37.4	40.4	1.3	0.11
Oxidative	44.5	47.4	1.4	0.16
Cross-sectional area ( $\mu\text{m}^2$ )				
I	3614	3039	155	0.01
IIA	5489	4322	246	0.002
IIX	6007	5522	294	0.26
Glycolytic	6780	7938	342	0.02
Oxidative	4040	3697	199	0.24
Relative area of the muscle (%)				
I	23.1	21.4	1.4	0.41
IIA	33.0	28.1	1.3	0.01
IIX	43.9	50.4	1.6	0.006
Oxidative	33.6	28.3	1.5	0.02

**Table 2** Protein fragments of longissimus thoracis muscle from Alentejana bulls produced according to continuous (CG) and discontinuous growth (DG) production systems

	CG (n = 17)	DG (n = 17)	SEM	P-value
MHC	83.91	86.48	5.38	0.74
Protein C	49.75	50.35	2.92	0.88
$\alpha$ -Actinin	46.60	52.00	1.81	0.04
kDa70	39.52	44.56	1.95	0.08
Tropomyosin	40.52	43.28	2.11	0.36
Actin	120.97	123.21	6.41	0.81
Troponin T	57.73	47.76	3.73	0.07
kDa36	36.25	40.93	2.08	0.13
kDa34	37.81	41.44	2.45	0.31
kDa31	41.70	37.08	2.12	0.14
kDa30	55.73	54.16	3.65	0.76
MLC1	57.45	63.35	3.66	0.27
Troponin I	68.10	60.62	3.69	0.15
Troponin C	35.58	35.47	2.59	0.97
MLC2	63.28	93.13	5.90	0.001

MLC, myosin light chain; MHC, myosin heavy chain.

**Table 3** Sensory panel scores and meat quality traits of longissimus thoracis muscle from Alentejana bulls produced according to continuous growth (CG) and discontinuous growth (DG) production systems

	CG (n = 17)	DG (n = 17)	SEM	P-value
Trained sensory panel				
Tenderness	4.89	5.46	0.11	0.0002
Juiciness	3.48	3.72	0.09	0.06
Off-flavour	0.10	0.14	0.04	0.55
Flavour	4.41	4.66	0.09	0.05
Global	4.61	5.16	0.09	<0.0001
Meat quality traits				
IMF (g/100 g) <sup>a</sup>	1.63	1.33	0.15	0.16
pH <sub>24h</sub> <sup>a</sup>	5.67	5.41	0.05	0.001
MFI <sub>7d</sub> <sup>a</sup>	34.3	28.9	2.7	0.17
WBSF (kg) <sup>a</sup>	7.73	6.96	0.43	0.23
CL (%)	35.4	35.7	0.9	0.83

IMF = intramuscular fat; MFI<sub>7d</sub> = myofibril fragmentation index after 7 days of ageing; WBSF = Warner-Bratzler shear force; CL = cooking loss.

<sup>a</sup>Published in Costa *et al.* (2015).

meat quality traits, except for pH<sub>24h</sub> ( $P < 0.01$ ), the production system had no significant influence on the IMF content, MFI after 7 days of ageing (MFI<sub>7d</sub>), WBSF and CL of Lt muscle.

### Correlations between fibre type profile, meat quality and sensory panel scores

Pearson's correlation coefficients for the relationship between muscle histological traits, meat quality and sensory scores are presented in Tables 4 and 5. As regards the CG samples, tenderness was positively correlated with IIX fibre type frequency ( $r = 0.51$ ,  $P = 0.04$ ), juiciness ( $r = 0.54$ ,  $P = 0.04$ ) and global acceptability ( $r = 0.88$ ,  $P < 0.001$ ). Furthermore, global acceptability was positively correlated

**Table 4** Pearson's correlation coefficients between fibre profile, meat quality and sensory panel scores of longissimus thoracis muscle from Alentejana bulls produced according to a continuous growth production system

	WBSF	Tenderness	Juiciness	Off-flavour	Flavour	Global
I (%)	0.06	-0.21	-0.15	-0.16	0.31	0.02
IIA (%)	-0.15	-0.32	-0.09	-0.09	0.12	-0.25
IIX (%)	-0.08	0.51*	0.23	0.25	-0.41	0.22
Oxidative (%)	-0.08	-0.25	-0.37	-0.06	0.11	-0.35
Glycolytic (%)	0.09	0.17	0.25	-0.02	0.09	0.13
CL (%)	0.06	-0.33	-0.64*	-0.06	-0.25	-0.61*
IMF (g/100 g)	0.10	0.08	-0.07	-0.28	-0.16	-0.03
pH <sub>24h</sub>	0.18	-0.04	0.07	-0.18	-0.16	0.08
MFI <sub>7d</sub>	0.16	0.29	0.01	0.21	0.46	0.30
WBSF (kg)	1.00	0.05	0.11	-0.08	0.08	0.10
Tenderness	0.05	1.00	0.54*	0.38	0.07	0.88***
Juiciness	0.11	0.54*	1.00	0.19	0.41	0.72***
Off-flavour	-0.08	0.07	0.19	1.00	0.25	0.40
Flavour	0.08	0.07	0.41	0.25	1.00	0.39
Global	0.10	0.88***	0.72***	0.40	0.39	1.00

WBSF = Warner-Bratzler shear force; CL = cooking loss; IMF = intramuscular fat content; MFI<sub>7d</sub> = myofibril fragmentation index after 7 days of ageing.

\* $P < 0.05$ , \*\*\* $P < 0.001$ .

**Table 5** Pearson's correlation coefficients between fibre profile, meat quality traits and sensory panel scores of longissimus thoracis muscle from Alentejana bulls produced according to a discontinuous growth production system

	WBSF	Tenderness	Juiciness	Off-flavour	Flavour	Global
I (%)	-0.67**	0.02	-0.22	-0.08	-0.16	0.03
IIA (%)	0.35	0.03	0.41	-0.01	0.37	0.09
IIX (%)	0.35	-0.05	-0.16	0.09	-0.18	-0.11
Oxidative (%)	-0.75**	-0.03	0.02	-0.14	-0.08	0.05
Glycolytic (%)	0.08	-0.14	-0.34	-0.08	-0.34	-0.16
CL (%)	-0.07	-0.03	-0.41	-0.08	0.18	-0.02
IMF (g/100 g)	-0.23	-0.13	0.06	-0.15	-0.06	-0.10
pH <sub>24h</sub>	0.00	0.33	0.22	0.73**	0.14	0.42
MFI <sub>7d</sub>	0.45	0.42	0.11	0.32	-0.22	0.32
WBSF (kg)	1.00	0.14	0.21	0.24	0.26	0.17
Tenderness	0.14	1.00	0.55*	-0.27	0.20	0.98***
Juiciness	-0.21	0.55*	1.00	-0.06	0.48	0.60*
Off-flavour	0.24	-0.27	-0.02	1.00	0.01	-0.40
Flavour	0.26	0.20	0.48	0.04	1.00	0.28
Global	0.17	0.98***	0.60*	-0.40	0.28	1.00

WBSF = Warner-Bratzler shear force; CL = cooking loss; IMF = intramuscular fat content; MFI<sub>7d</sub> = myofibril fragmentation index after 7 days of ageing.


\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

with juiciness ( $r = 0.72$ ,  $P = 0.002$ ). Negative correlations were recorded between CL and juiciness ( $r = -0.64$ ,  $P = 0.01$ ) and global acceptability ( $r = -0.61$ ,  $P = 0.02$ ).


As for the DG production system samples, noticeable negative correlations were observed between WBSF and type I ( $r = -0.67$ ,  $P = 0.006$ ) and oxidative ( $r = -0.75$ ,  $P = 0.001$ ) fibres. Moreover, a positive correlation was observed between pH<sub>24h</sub> and off-flavour ( $r = 0.73$ ,  $P = 0.001$ ). The increase of tenderness was associated with greater juiciness ( $r = 0.55$ ,  $P = 0.03$ ). In addition, the global acceptability was positively correlated with tenderness ( $r = 0.98$ ,  $P < 0.001$ ) and juiciness ( $r = 0.60$ ,  $P = 0.02$ ).

#### Principal component analysis

The PCA for the CG and DG production systems are presented in Figures 1 and 2, respectively. The PC1 in CG bulls explained 29.4% of the total variation. Two groups of variables lying on the PC1 were distinguished. The first group included several muscle protein fragments in the right side of the plot. In opposite direction, tenderness, global acceptability, juiciness, IIX fibres and IMF formed a distinguished group lying in the left hand side of the plot. The PC2 explained 20.2% of the variance and it was influenced by MLC2, CL and actin and by troponin T, juiciness and kDa36 in the opposite extreme of the plot.



**Figure 1** Projection of the variables in the plane defined by the first two principal components (PC) for *longissimus thoracis* muscle of Alentejana bulls produced according to continuous growth production system. Ox = oxidative; Gly = glycolytic; CL = cooking loss; IMF = intramuscular fat content; MFI<sub>7d</sub> = myofibril fragmentation index after 7 days of ageing; WBSF = Warner-Bratzler shear force; MLC = myosin light chain.



**Figure 2** Projection of the variables in the plane defined by the first two principal components (PC) for *longissimus thoracis* muscle of Alentejana bulls produced according to discontinuous growth production system. Ox = oxidative; Gly = glycolytic; CL = cooking loss; IMF = intramuscular fat content; MFI<sub>7d</sub> = myofibril fragmentation index after 7 days of ageing; WBSF = Warner-Bratzler shear force; MLC = myosin light chain.

In DG bulls, PC1 and PC2 explained 29.4% and 14.3% of the variance, respectively. The PC1 was mainly characterised by a group of variables that include muscle protein fragments in the left side and flavour, juiciness, off-flavour and MFI<sub>7d</sub> in the right side of the plot. The PC2 was mainly affected by fibre

type I and oxidative, and off-flavour in one direction, whereas in the opposite direction it was mainly characterised by pH<sub>24h</sub>, tropomyosin, WBSF, global acceptability and tenderness.

**Discussion**

*Muscle fibre characteristics*

Restricted feeding has been reported to increase the proportion of red fibres in *longissimus dorsi* muscle (Seideman and Crouse, 1986; Nicastro and Maiorano, 1994) and to reduce fibre hypertrophy in skeletal muscle of bovines (Yambayamba and Price, 1991). However, when the effect of feed restriction was studied at similar BW and after 3 to 4 months of *ad libitum* fed, no differences in fibre size and proportions were found between older restricted bovines and their younger and unrestricted counterparts (Yambayamba and Price, 1991). Cassar-Malek *et al.* (2004) reported that a DG path affected muscle metabolism but not histological profile. The authors hypothesised that feed restriction (from 9 to 12 months of age) was not severe enough to induce fibre type transitions. In this study, the average daily gain (ADG) from 9 to 15 months of age was around 1.5 kg/day greater ( $P < 0.0001$ ) in CG than in DG Alentejana bulls. In contrast, in the period from 15 to 18 months of age, the ADG was almost 0.5 kg greater ( $P = 0.0004$ ) in DG than in CG bovines. ADG for DG bulls were around 1.3 and 1.1 kg/day from 18 to 21 months of age and from 21 to 24 months of age, respectively (Costa *et al.*, 2015). The fibre type frequency was not affected by growth path (Table 1). However, the fibre CSA of types I and IIA in DG bulls were significantly lower than in the CG bulls while the CSA of glycolytic fibres was greater in DG bulls. Consequently, the RA occupied by type IIX fibres was greater, while the IIA fibre RA was lower in DG in relation to CG. The RA of oxidative fibres was significantly higher in CG than in DG. It appears that the long re-feeding period (9 months) did not compensate the effects of the restrictive period (6 months) on muscle fibre profile of Alentejana bulls.

*Meat quality traits*

Beef tenderness, juiciness and flavour are among the most important eating quality attributes for consumers (Calkins and Hodgen, 2007; Hocquette *et al.*, 2014). Tenderness is considered to be a major influence on the consumers' acceptance, and repetitive purchase, of a particular type of beef (Calkins and Hodgen, 2007). It depends on the architecture and integrity of the skeletal muscle cells and on events that modify those cells (Huff Lonergan *et al.*, 2010). The impact of growth rate on tenderness could be related to its effect on the structure and crosslinking of the collagen matrix (Allingham *et al.*, 1998), as well as on the proteolytic activity and glycolysis rate of the muscle myofibrils (Oddy *et al.*, 2001).

After 7 days of ageing, α-actinin and MLC2 were the only protein fragments that were significantly influenced by growth path (i.e. greater in DG bulls) (Table 2). MLC2 has a regulatory role in muscle contraction (Kamm and Stull,

2011), whereas  $\alpha$ -actinin anchors myofibrillar actin filaments in the Z-disc. These differences in muscle protein degradation fragments between DG and CG bulls result from different disruption of muscle cell integrity. This disruption is influenced by compensatory growth (Therkildsen, 2005) and could help to explain the differences found on meat sensory tenderness between the two groups (Table 3). Meat tenderness is influenced by *postmortem* activity of muscle proteolytic enzymes. As these enzymes are also involved in protein turnover in the living muscles, it could be expected that feed restriction, followed by re-alimentation, could have the potential to improve tenderness (Therkildsen, 2005). A fast growth, as observed during compensatory phase, increases protein turnover and the ratio between protein synthesis and degradation in connective tissue and myofibrils, making these younger and more liable to breakdown during conditioning and cooking. After the feed restriction period (from 9 to 15 months of age) during which the ADG was around 100 g/day, the DG bulls had a compensatory growth phase in which the ADG was of almost 1700 g/day (Costa *et al.*, 2015). However, the production system had no significant impact on WBSF, contrary to the results of previous studies that used *longissimus* muscle (Muir *et al.*, 2001; Therkildsen, 2005; Hansen *et al.*, 2006) but used a compensatory period of <5 months. Despite not observed for Lt muscle, *supraspinatus*, *triceps brachii*, *semitendinosus* and *biceps femoris* muscles from DG bulls exhibited lower WBSF values than CG bulls (Costa *et al.*, 2015).

The longer (9 months) re-feeding period that was implemented here may have partially diluted the effects of compensatory growth, particularly on Lt muscle. The long re-feeding period is necessary to produce commercially acceptable carcasses but could be responsible for a decrease on meat tenderness perceived by the consumer. Studies regarding the best feed manage to finish Alentejana bulls, produced according to DG, are warranted in order to obtain homogeneous high meat quality.

The fibre proportions influences muscle *postmortem* conversion into meat, and may therefore affect meat quality (Ozawa *et al.*, 2000). As regards the association between muscle fibre profile and meat sensory panel scores, Maltin *et al.* (1998) reported a positive correlation between tenderness and proportion of slow-twitch oxidative fibres. They noted a negative correlation between tenderness and frequency of fast-twitch glycolytic fibres. Using a different fibre classification system, Calkins *et al.* (1981) reported that  $\alpha$ W muscle fibre content was negatively correlated with marbling and tenderness while at the same time the  $\alpha$ R content was positively correlated to these attributes. Contrary to these results, we observed a positive correlation between tenderness and the proportion of IIX fibres in CG but not in DG bulls. Interestingly, increasing the proportion of type I and oxidative fibres was associated with lower WBSF in the latter group. According to Xiong *et al.* (2007), muscles with a greater proportion of type II fibres are more susceptible to early *postmortem* proteolytic degradation than are muscles that mainly comprise type I fibres. This can be due to a higher

calpain/calpastatin ratio in fast-twitch glycolytic than in slow-twitch oxidative muscles (Ouali and Talmant, 1990). It is commonly reported that muscle fibre composition influences many aspects of meat quality, including colour, water holding capacity, tenderness, juiciness and flavour. However, identifying the specific relationships between meat quality and myofibre characteristics, including fibre types and CSA, remains a difficult task (Lee *et al.*, 2010; Lefaucheur, 2010). As expected, increasing tenderness and juiciness were associated with higher global acceptability which is in accordance with a previous study from our research team (Costa *et al.*, 2012). No significant correlations were observed between IMF, MFI<sub>7d</sub> and WBSF and sensory panel scores in both groups, CG and DG. The IMF has an important role in meat palatability. However, the effect on sensory attributes depends on its level in meat. A minimum value of 2.5% IMF has been proposed as necessary for sensory acceptability of pork (Madeira *et al.*, 2013). The low level of IMF in Lt muscle from experimental groups (<2 g/100 g of fresh beef; Costa *et al.*, 2015) could have not been enough to affect the trained sensory panel scores. It is possible that at 24 months of age, part of the effects of compensatory growth on myofibrillar proteins were diluted and no longer observed, which could justify the lack of relationship between beef sensory attributes and MFI<sub>7d</sub> in DG bovines. The absence of this relationship was not expected for CG bulls. In general, the correlation between tenderness and WBSF is significant across muscles. However, this is not always observed and sometimes the tenderness perceived by a sensory panel and the instrumental WBSF are not in accordance (Costa *et al.*, 2012).

The PCA could be a very effective procedure to obtain a synthetic judgement of meat quality (Destefanis *et al.*, 2000). The PCA for CG bulls (Figure 1) and DG bulls (Figure 2) showed distinct associations among meat traits, suggesting that growth pathway has an important effect on Alentejana beef quality. The tenderness, global acceptance, juiciness, flavour, WBSF, MFI<sub>7d</sub>, IIX and IIA fibres formed a distinguishable group, suggesting a relationship among these variables in DG bulls. The close association between tenderness and global acceptability, also observed in CG bulls, was expected (Costa *et al.*, 2012) and confirms the decisive role of tenderness in beef acceptability.

## Conclusions

Taking advantage of compensatory growth in bovine production could provide an economic opportunity for Portuguese cattle breeders. The compensatory growth production system appears to have a positive impact on meat tenderness and global acceptability, overcoming the negative effects of age at slaughter on collagen solubility and tenderness. In addition, compensatory growth affects muscle fibre characteristics. It also influences pH<sub>24h</sub> and  $\alpha$ -actinin and MLC2 protein levels. However, a significant relationship between global palatability of meat and its muscle fibre characteristics has not yet been established.

## Acknowledgements

This research was supported by the Fundação para a Ciência e a Tecnologia (FCT) through financial support from FCT grants (PTDC/CVT/65707/2006 and PTDC/CVT/111744/2009) and individual fellowships to P. C. (SFRH/BPD/111051/2015), S.P.A. (SFRH/BPD/76836/2011) and V. V. (SFRH/BPD/104394/2014). P. A. L. is a researcher from the FCT 'Ciência 2008' programme and Incentivo 2014 project (AGR/UI0276/2014). We are grateful to Eng. António Sequeira for his decisive role on animal management during the trial. The authors are grateful to EZN slaughterhouse staff, particularly to Paula Menezes, for her help with carcass data and meat sample collection. The authors acknowledge Luisa Cristina Roseiro and Carlos Santos for help on histological analysis.

## References

- Allingham PG, Harperand GS and Hunter RA 1998. Effect of growth path on the tenderness of the semitendinosus muscle of Brahman-cross steers. *Meat Science* 48, 65–73.
- American Meat Science Association 2015. Research guidelines for cookery, sensory evaluation, and instrumental tenderness measurements of meat. AMSA research guidelines for cookery and evaluation, version 1.0. AMSA, Champaign, IL, USA.
- Association of Official Analytical Chemists 2000. Official methods of analysis. AOAC, Arlington, VA, USA.
- Brooke MH and Kaiser KK 1970. Muscle fiber types: how many and what kind? *Archives of Neurology* 23, 369–379.
- Calkins CR, Dutson TR, Smith GC, Carpenterand ZL and Davis GW 1981. Relationship of fiber type composition to marbling and tenderness of bovine muscle. *Journal of Food Science* 46, 708–710.
- Calkins CR and Hodgen JM 2007. A fresh look at meat flavor. *Meat Science* 77, 63–80.
- Cassar-Malek I, Hocquette JF, Jurie C, Listrat A, Jailler R, Bauchart D, Briand Y and Picard B 2004. Muscle-specific metabolic, histochemical and biochemical responses to a nutritionally induced discontinuous growth path. *Animal Science* 79, 49–59.
- Costa P, Lemos JP, Lopes PA, Alfaia CM, Costa ASH, Bessa RJB and Prates JAM 2012. Effect of low- and high-forage diets on meat quality and fatty acid composition of Alentejana and Barrosã beef breeds. *Animal* 6, 1187–1197.
- Costa P, Simões JA, Costa ASH, Lemos JPC, Navas D, Hocquette JF, Calkins CR and Bessa RJB. 2015. Repercussions of growth path on carcass characteristics, meat colour and shear force in Alentejana bulls. *Animal* 9, 1414–1422.
- Cross H, Moen RR and Stanfield MS 1978. Training and testing of judges for sensory analysis of meat quality. *Food Technology* 32, 48–54.
- Culler RD, Smith GC and Cross HR 1978. Relationship of myofibril fragmentation index to certain chemical, physical and sensory characteristics of bovine longissimus muscle. *Journal of Food Science* 43, 1177–1180.
- Destefanis G, Barge MT, Brugiapaglia A and Tassone S 2000. The use of principal component analysis (PCA) to characterize beef. *Meat Science* 56, 255–259.
- Fishell VK, Aberle ED, Judge MD and Perry TW 1985. Palatability and muscle properties of beef as influenced by preslaughter growth rate. *Journal of Animal Science* 61, 151–157.
- Fritz JD and Greaser ML 1991. Changes in titin and nebulin in postmortem bovine muscle revealed by gel electrophoresis, western blotting and immunofluorescence microscopy. *Journal of Food Science* 56, 607–610.
- Hansen S, Therkildsen M and Byrne DV 2006. Effects of a compensatory growth strategy on sensory and physical properties of meat from young bulls. *Meat Science* 74, 628–643.
- Hocquette JF, Van Wezemael L, Chriki S, Legrand I, Verbeke W, Farmer L, Scollan ND, Polkinghorne R, Rødbotten R, Allen P and Pethick DW 2014. Modelling of beef sensory quality for a better prediction of palatability. *Meat Science* 97, 316–322.
- Huff Lonergan E, Zhang W and Lonergan SM 2010. Biochemistry of postmortem muscle – lessons on mechanisms of meat tenderization. *Meat Science* 86, 184–195.
- Kamm KE and Stull JT 2011. Signaling to myosin regulatory light chain in sarcomeres. *The Journal of Biological Chemistry* 286, 9941–9947.
- Lee SH, Joo ST and Ryu YC 2010. Skeletal muscle fiber type and myofibrillar proteins in relation to meat quality. *Meat Science* 86, 166–170.
- Lefaucheur L 2010. A second look into fiber typing – relation to meat quality. *Meat Science* 84, 257–270.
- Maccatrozzo L, Patrino M, Toniolo L, Reggiani C and Mascarello F 2004. Myosin heavy chain 2B isoform is expressed in specialized eye muscles but not in trunk and limb muscles of cattle. *European Journal of Histochemistry* 48, 357–366.
- Madeira MS, Costa P, Alfaia CM, Lopes PA, Bessa RJB, Lopes PA, Lemos JCP and Prates JAM 2013. The increased intramuscular fat promoted by dietary lysine restriction in lean but not in fatty pig genotypes improves pork sensory attributes. *Journal of Animal Science* 91, 3177–3187.
- Maltin CA, Sinclair KD, Warriss PD, Grant CM, Porter AD, Delday MI and Warkup CC 1998. The effects of age at slaughter, genotype and finishing system on the biochemical properties, muscle fiber type characteristics and eating quality of bull beef from suckled calves. *Animal Science* 66, 341–348.
- Meilgaard MC, Civille GV and Carr BT 2006. Sensory evaluation techniques, 4th edition. CRC Press, Boca Raton, FL, USA.
- Muir PD, Smith NB, Dobbie PM, Smith DR and Bown MD 2001. Effects of growth pathway on beef quality in 18-month-old Angus and South Devon × Angus pasture-fed steers. *Animal Science* 72, 297–308.
- Nicastro FM and Maiorano G 1994. Histological characteristics of chianina breed related to meat quality. Expression regulation and role of proteinases in muscle development and meat quality. *Sciences des Aliments* 14, 423–430.
- Oddy VH, Harper GS, Greenwood PL and McDonagh MB 2001. Nutritional and developmental effects on the intrinsic properties of muscles as they relate to the eating quality of beef. *Australian Journal of Experimental Agriculture* 41, 921–942.
- Ouali A and Talmant A 1990. Calpains and calpastatin distribution in bovine, porcine and ovine skeletal muscles. *Meat Science* 28, 331–348.
- Oury MP, Dumont R, Jurie C, Hocquette JF and Picard B 2010. Specific fiber composition and metabolism of the rectus abdominis muscle of bovine Charolais cattle. *BMC Biochemistry* 11, 1–12.
- Ozawa S, Mitsuhashi T, Mitsumoto M, Matsumoto S, Itoh N, Itagaki K, Kohno Y and Dohgo T 2000. The characteristics of muscle fiber types of longissimus thoracis muscle and their influences on the quantity and quality of meat from Japanese Black steers. *Meat Science* 54, 65–70.
- Parrish FC, Young RB, Miner BE and Andersen LD 1973. Effect of postmortem conditions on certain chemical, morphological and organoleptic properties of bovine muscle. *Journal of Food Science* 38, 690–695.
- Picard B and Cassar-Malek I 2009. Evidence for expression of IIb myosin heavy chain isoform in some skeletal muscles of Blonde d'Aquitaine bulls. *Meat Science* 82, 30–36.
- Picard B, Duris MP and Jurie C 1998. Classification of bovine muscle fibers by different histochemical techniques. *The Histochemical Journal* 30, 473–477.
- Seideman SC and Crouse JD 1986. The effects of sex condition, genotype and diet on bovine muscle fiber characteristics. *Meat Science* 17, 55–72.
- Sheehan DC and Hrapchak BB 1987. Theory and practice of histotechnology. Battelle Memorial Institute, Columbus, OH, USA.
- Therkildsen M 2005. Muscle protein degradation in bull calves with compensatory growth. *Livestock Production Science* 98, 205–218.
- Toniolo L, Maccatrozzo L, Patrino M, Caliaro F, Mascarello F and Reggiani C 2005. Expression of eight distinct MHC isoforms in bovine striated muscles: evidence for MHC-2B presence only in extraocular muscles. *Journal of Experimental Biology* 208, 4243–4253.
- Whipple G and Koohmaraie M 1992. Effects of lamb age, muscle type, and 24-hour activity of endogenous proteinases on postmortem proteolysis. *Journal of Animal Science* 70, 798–804.
- Xiong YL, Mullins OE, Stika JF, Chen J, Blanchard SP and Moody WG 2007. Tenderness and oxidative stability of post-mortem muscles from mature cows of various ages. *Meat Science* 77, 105–113.
- Yambayamba EP and Price M 1991. Fiber-type proportion and diameters in the longissimus muscle of beef heifers undergoing catch-up (compensatory) growth. *Canadian Journal of Animal Science* 71, 1031–1035.