

October 2006

# Fast and efficient generation of influenza A virus like particles from synthetic genes

Theresa Schinko

*University of Natural Resources and Applied Life Sciences, Vienna, Austria, [theresa.schinko@boku.ac.at](mailto:theresa.schinko@boku.ac.at)*

Harutha Thaisuchat

*University of Natural Resources and Applied Life Sciences, Vienna, Austria, [Harutha.Thaisuchat@boku.ac.at](mailto:Harutha.Thaisuchat@boku.ac.at)*

Hendrik Viljoen

*University of Nebraska - Lincoln, [hviljoen1@unl.edu](mailto:hviljoen1@unl.edu)*

Nisha Padhye

*University of Nebraska - Lincoln*

Reingard Grabherr

*University of Natural Resources and Applied Life Sciences, Vienna, Austria, [reingard.grabherr@boku.ac.at](mailto:reingard.grabherr@boku.ac.at)*

Follow this and additional works at: <http://digitalcommons.unl.edu/cbmeviljoen>



Part of the [Chemical Engineering Commons](#)

---

Schinko, Theresa; Thaisuchat, Harutha; Viljoen, Hendrik; Padhye, Nisha; and Grabherr, Reingard, "Fast and efficient generation of influenza A virus like particles from synthetic genes" (2006). *Hendrik J. Viljoen Publications*. 1.  
<http://digitalcommons.unl.edu/cbmeviljoen/1>

This Article is brought to you for free and open access by the Chemical and Biomolecular Research Papers -- Faculty Authors Series at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Hendrik J. Viljoen Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Poster Presentation

Open Access

## Fast and efficient generation of influenza A virus like particles from synthetic genes

Theresa Schinko<sup>\*1</sup>, Haruthai Thaisuchat<sup>1</sup>, Hendrik Viljoen<sup>2</sup>, Nisha Padhye<sup>2,3</sup> and Reingard Grabherr<sup>1</sup>

Address: <sup>1</sup>Institute of Applied Microbiology, University of Natural Resources and Applied Life Sciences, Vienna, Austria, <sup>2</sup>Department of Chemical Engineering, University of Nebraska, Lincoln, NE 68588 USA and <sup>3</sup>Megabase Research Products, 2726N.48th Street, Lincoln, NE 68504 USA

\* Corresponding author

from The 4th Recombinant Protein Production Meeting: a comparative view on host physiology  
Barcelona, Spain. 21–23 September 2006

Published: 10 October 2006

*Microbial Cell Factories* 2006, **5**(Suppl 1):P39 doi:10.1186/1475-2859-5-S1-P39

© 2006 Schinko et al; licensee BioMed Central Ltd.

### Background

With the recent emergence of the bird flu in many European countries, molecular biologists are challenged more than ever to advance the present methods of vaccine development. We have developed a method that is based on the following key elements: safety, efficacy and rapidity. Influenza A virus like particles (VLP) were generated in insect cells by co-transfection of especially designed plasmids. In order to produce VLPs, four recombinant baculoviruses were generated each containing two influenza genes under control of the *Autographa californica* multiple nuclear polyhedrosis virus (*AcMNPV*) p<sub>H</sub> and p<sub>10</sub> promoters for high level expression in *Sf9* insect cells. VLPs contained 8 of 10 influenza A virus proteins of strain PR8, missing NS1 and NS2. Alternatively, VLPs were generated, by assembly of just three proteins, HA, NA and M1, which are responsible for induction of the immune system *in vivo*. All influenza genes have been produced from synthetic oligonucleotides, using a rapid thermocycler, the PCRJet<sup>®</sup>. Synthetic genes of new emerging influenza A variants can be produced accurately and rapidly by using this technology. Recombinant baculoviruses were generated using the Bac-to-Bac system by homologous recombination between a transfer vector and the baculoviral shuttle vector (bacmid) in DH10Bac cells. By optimizing DNA synthesis and gene transfer into *Sf9* cells, we anticipate major improvements in flexibility, speed and yield of influenza vaccine production as compared to available technologies.

### Results

Co-transfection of bacmids resulted in the generation of influenza A virus like particles in the supernatant of *Sf9* cells. VLPs were purified by means of Sucrose gradient centrifugation and the expected results were confirmed by Electron microscopy, Western Blot analysis and hemagglutination assays.