

1-1-2004

Hospital Universitario, “La Paz”, Madrid, SPAIN

Michael G. Zeece

University of Nebraska - Lincoln, mzeece1@unl.edu

Gautam Sarath

University of Nebraska - Lincoln, Gautam.sarath@ars.usda.gov

John P. Markwell

University of Nebraska - Lincoln, jmarkwell2@unl.edu

N. Palmer

University of Nebraska-Lincoln, nathan.palmer@ars.usda.gov

J. Steckelberg

University of Nebraska - Lincoln

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/biochemistrysarath>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#)

Zeece, Michael G.; Sarath, Gautam; Markwell, John P.; Palmer, N.; Steckelberg, J.; and Schultz, J., "Hospital Universitario, “La Paz”, Madrid, SPAIN" (2004). *Gautam Sarath Publications*. 3.
<http://digitalcommons.unl.edu/biochemistrysarath/3>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Gautam Sarath Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Michael G. Zeece, Gautam Sarath, John P. Markwell, N. Palmer, J. Steckelberg, and J. Schultz

521 Microarray Assessment of IgE-Binding Proteins in Lentils (*Lens culinaris L.*)

M. G. Zeece¹, G. Sarath², J. Markwell³, N. Palmer¹, J. Steckelberg¹, J. Schultz⁴, C. Pascual⁵; ¹Food Science, University of Nebraska, Lincoln, NE, ²USDA-ARS, Lincoln, NE, ³Biochemistry, University of Nebraska, Lincoln, NE, ⁴University of Nebraska, Lincoln, NE, ⁵Hospital Universitario, "La Paz", Madrid, SPAIN.

RATIONALE: Proteomic technology has been shown to have distinct advantages for allergen characterization and identification. Therefore, we have used 2-D electrophoresis and protein microarrays to characterize the profile of allergens and assess their level in lentils.

METHODS: Lentil varieties were extracted and centrifuged to obtain 2S, 7S and 11S fractions. The profile of IgE-binding proteins in these fractions was assessed using lentil-allergic sera, by immunoblotting of 2-D electrophoretic separations. The level of IgE-binding was also assessed using protein microarrays. Arrays of covalently attached proteins were first incubated with lentil-allergic sera, followed by incubation with fluorescently-labeled anti-IgE detection antibody. Laser-induced fluorescence was used to detect IgE-binding.

RESULTS: Immunoblotting of electrophoretic separations showed several regions of IgE-binding that were consistently observed for all varieties. Major IgE-binding was observed for isoforms of 31 and 48 kDa spots. These proteins are of particular interest because of their low abundance, but high IgE-binding. Comparatively less IgE-binding was found for 12 and 14 kDa spots. Microarray analysis showed that IgE-binding of the 11S fraction was significantly greater than all other fractions. Microarrays enabled quantitative assessment of IgE-binding.

CONCLUSIONS: The application of proteomic technologies resulted in characterization of IgE-binding protein profile and determination of their level in lentils. Microarrays represent a promising tool for allergen investigation, detection and diagnosis.

Funding: University of Nebraska

SUNDAY