

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Proceedings of the North American Prairie
Conferences

North American Prairie Conference

1989

Correlations Between Insects and Birds in Tallgrass Prairie Riparian Habitats

Lawrence J. Gray

Department of Biology, Ottawa University, Ottawa, Kansas

Follow this and additional works at: <http://digitalcommons.unl.edu/napcproceedings>



Part of the [International and Area Studies Commons](#)

Gray, Lawrence J., "Correlations Between Insects and Birds in Tallgrass Prairie Riparian Habitats" (1989). *Proceedings of the North American Prairie Conferences*. 18.

<http://digitalcommons.unl.edu/napcproceedings/18>

This Article is brought to you for free and open access by the North American Prairie Conference at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Proceedings of the North American Prairie Conferences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

CORRELATIONS BETWEEN INSECTS AND BIRDS IN TALLGRASS PRAIRIE RIPARIAN HABITATS

Lawrence J. Gray

Department of Biology, Ottawa University, Ottawa, Kansas 66067

Abstract. Previous observations in riparian habitats of Kings Creek, Konza Prairie Research Natural Area, Kansas indicated that emerging aquatic insects represent a concentrated source of food for insectivorous birds, particularly the flycatcher and gleaner guilds. This hypothesis was tested by concurrent measurements of net insect emergence (total emergence minus adults returning to the stream) and densities of birds at six sampling sites from June to August 1987 and May to June 1988. Significant positive correlations with emergence were found for flycatchers ($r = 0.93$) and gleaners ($r = 0.91$), the two insectivore guilds containing the majority of individuals along Kings Creek. Observations also showed that flycatcher and gleaner populations rapidly respond to temporal changes in insect emergence.

Key Words. fly catcher, gleaner, bird, aquatic insect, tallgrass prairie, stream, Kansas

INTRODUCTION

Streams in the prairie and desert regions of western North America support riparian ecosystems that are of critical importance as habitats for local and regional avifaunas (Knopf *et al.* 1988). In addition to providing cover and water, these habitats provide a concentrated source of insects as food for avian insectivores. Along desert streams, this insect food appears to consist mainly of emerging aquatic insects, such as midges and mayflies. Jackson and Fisher (1986), for example, found that the annual biomass of emerging aquatic insects from Sycamore Creek, Arizona, was over 23 g/m².

Emergence biomass from Kings Creek, a tallgrass prairie stream in the Konza Prairie Research Natural Area (KPRNA), averages 20.3 mg/m²/day in the summer (Gray, unpublished). Field observations during previous summers had suggested that certain species, particularly flycatchers, were found in greater abundance along those stream reaches with the highest rates of emergence. Thus the objective of this study was to determine if riparian habitats of tallgrass prairie streams, like desert streams, serve as an important feeding area for insectivorous birds by testing the hypothesis that insectivore densities should be positively correlated with emergence biomass.

Previous studies of birds in the KPRNA primarily have dealt with typical grassland species (Zimmerman 1982, Finck 1984) rather than riparian species. Published information on riparian species is confined to a listing of seasonal occurrence and breeding status (Zimmerman 1985).

METHODS

The Kings Creek catchment (1,637 ha) lies entirely within the boundaries of KPRNA. The headwaters are typically ephemeral, except near springs and seeps, whereas intermittent and perennial reaches occur downstream. Riparian vegetation in the headwaters is composed of shrubs, small trees, grasses, and sedges. A relatively abrupt transition to a gallery forest of oak (*Quercus* L. spp.), hackberry (*Celtis* L. spp.), and elm (*Ulmus* L. spp.) occurs along lower stream reaches (Gurtz *et al.* 1982). In this study, three sampling sites were along third- and fourth-order channels in the prairie/shrub vegetation type, and three sites were located in fourth- and fifth-order channels bordered by gallery forest. Two of the prairie/shrub sites had flow during April and May but then dried completely by late June. The other four sites had flow until late

June; discharge then declined until only isolated pools remained in July and August. No scouring floods occurred at any site. Discharge was measured by using a meter stick, stopwatch, and fluorescein dye.

Aquatic insect adults emerging from the stream were collected in traps constructed from inverted 15-liter plastic buckets similar in design to that used by Jackson and Fisher (1986). Each trap collected adults emerging from an area of 452 cm². Traps were emptied each 24-hour period using an aspirator. The number of traps placed at each site varied from 10 to 25, depending on flow conditions and habitat area. Adults were preserved in 80% ethanol, sorted by taxon, and dried at 80 C for 24 hours before weighing on an analytical balance.

Aquatic adults returning to the stream were collected by isolating a short section of channel with two drift nets (1.0 mm mesh) for a 24-hour period. Returning biomass was calculated by dividing the biomass collected in the downstream net by the channel area between nets. Net emergence equalled trap biomass minus returning biomass. Net emergence/m² was multiplied by the square meters of wetted area in 100 linear meters of channel to compensate for variations in total stream area.

Birds were censused by recording all sightings within an area of 0.2 ha for 15 minutes. For a given site and date, three replicate censuses were taken along a 250 m reach of stream channel. All censuses occurred within two hours after sunrise during fair weather. Census counts were not adjusted for undetectable (silent) birds (Emlen 1977). On the eight occasions when birds were counted at a site for a period of several hours, no birds were found in addition to those counted in the three, 15-minute censuses.

Insectivorous birds were placed into one of four feeding guilds based on the principal method of feeding as observed in the field. Gleaners actively search vegetation for stationary prey, whereas flycatchers wait for active prey (usually flying insects) to enter their field of view before moving from a perch. Sweepers feed on flying insects during flight, often ranging over a large area. Woodpeckers excavate deeply in wood and bark. Emphasis is placed here on the gleaners and flycatchers. The census method was not suitable for measuring the densities of sweepers, especially the common nighthawk (*Chordeiles minor*), the most abundant species. Woodpeckers were rare. Stream emergence and bird densities were determined for a total of 22 site-date combinations during June to August 1987 (N = 10) and late May to June 1988. Each study site was sampled at least once each year. Gallery forest sites were sampled a total of 12 times. For N = 22 (d.f. = 20), a correlation coefficient (r) > 0.42 is significant at P = 0.05; r > 0.54 insignificant at P = 0.01.

DISCUSSION

The emergence of aquatic insects from Kings Creek shows wide spatial and temporal variability (Gray, unpublished). For the sampling dates in this study, daily emergence varied from 0 to 21 g/100 m. Emergence is strongly dependent on stream flow. When stream channels dry to isolated pools, daily emergence biomass is 10 to 100 times less than that when stream flow is stable.

The most common insects emerging from Kings Creek were midges (Diptera: Chironomidae), mayflies (Ephemeroptera), and stoneflies (Plecoptera). These groups comprised 53%, 27%, and




FIG. 3. Reduction in the combined densities of flycatchers and gleaners at a gallery forest site in response to decreases in aquatic insect emergence. Stream discharge declined from 100 l/sec on 11 June to 0 l/sec by 8 August.

Until the diets of insectivorous birds in riparian habitats of the Konza prairie can be analyzed in more detail, the contribution by stream insects is unknown. Although other factors are undoubtedly influencing the distribution of gleaners and flycatchers (e.g., competition, availability of nest sites, etc.), the strong correlations between bird densities and insect emergence, and field observations of feeding behavior, suggest that these birds are dependent to some extent on stream insects. Thus riparian habitats of tallgrass prairie streams, like that of desert streams, appear to be a concentrated source of insect food for riparian insectivores.

ACKNOWLEDGEMENTS

K. Johnson assisted with the field work, and E. Finck provided advice in the design of the bird census method. This research was supported by National Science Foundation Grant BSR-8704333 and the Division of Biology, Kansas State University, Manhattan.

LITERATURE CITED

- Emlen, J.T. 1977. Estimating breeding season bird densities from transect counts. *The Auk* 94:455-468.
- Finck, E.J. 1984. Male dickcissel behavior in primary and secondary habitats. *Wilson Bulletin* 96:672-680.
- Gurtz, M.E., G.R. Marzolf, K.T. Killingbeck, D.L. Smith, and J.V. McArthur. 1982. Organic matter loading and processing in a pristine stream draining a tallgrass prairie/riparian forest watershed. Contribution No. 230, Kansas Water Resources Research Institute, Manhattan.
- Jackson, J.K. and S.G. Fisher. 1986. Secondary production, emergence, and export of aquatic insects of a Sonoran Desert stream. *Ecology* 67:629-638.
- Knopf, F.L., R.R. Johnson, T. Rich, F.B. Samson, and R.C. Szaro. 1988. Conservation of riparian ecosystems in the United States. *Wilson Bulletin* 100:272-284.
- Zimmerman, J.L. 1982. Nesting success of Dickcissels (*Spiza americana*) in preferred and less preferred habitats. *The Auk* 99:292-298.
- Zimmerman, J.L. 1985. The birds of Konza Prairie Research Natural Area, Kansas. *Prairie Naturalist* 17:185-192.