

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications: Department of Entomology

Entomology, Department of

February 2004

Fitness Characteristics of Transgenic Strains of *Cochliomyia Hominivorax*, the New World Screwworm

Margaret Allen

Dennis Berkebile

USDA-ARS, DENNIS.BERKEBILE@ars.usda.gov

Steven Skoda

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

Allen, Margaret ; Berkebile, Dennis; and Skoda, Steven, "Fitness Characteristics of Transgenic Strains of *Cochliomyia Hominivorax*, the New World Screwworm" (2004). *Faculty Publications: Department of Entomology*. 10.
<http://digitalcommons.unl.edu/entomologyfacpub/10>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Title: Fitness Characteristics of Transgenic Strains of *Cochliomyia Hominivorax*, the New World Screwworm

Authors

- [Allen, Margaret](#)
- [Berkebile, Dennis](#)
- [Skoda, Steven](#)

Submitted to: Keystone Symposia

Publication Type: Abstract

Publication Acceptance Date: October 6, 2003

Publication Date: February 5, 2004

Citation: Allen, M.L., Berkebile, D.R., Skoda, S.R. 2004. Fitness characteristics of transgenic strains of *Cochliomyia hominivorax*, the New World Screwworm [abstract]. Keystone Symposia, "Genetic Manipulation of Insects". Poster No. 201, Program Book p. 57.

Interpretive Summary: Sterile insect technique (SIT) programs are designed to eradicate pest species by release of mass-reared, sterile insects into an infested area. The first large-scale implementation of SIT was the New World Screwworm Eradication Program, which successfully eliminated the New World screwworm (NWS), *Cochliomyia hominivorax* (Coquerel) (Diptera: Calliphoridae) from the Continental US, Mexico, and most of Central America. Ionizing radiation is currently used for sterilization, but transgenic insect techniques could hypothetically replace this method, providing a safer, more cost-effective alternative. Genetic transformation methods have recently been demonstrated in NWS. Some key issues involved in mass release of sterile insects revolve around fitness. Transgenic insects must be highly productive in mass rearing, and equally capable of surviving and mating with wild insects in order to be truly effective. Therefore it is critical to analyze fitness characteristics of transgenic insects. Eight transgenic strains of *C. hominivorax* were compared with the wild-type parental laboratory strain. None of the transgenic colonies exhibited significantly lower adult fitness characteristics than the control parental colony. Overall, the presence of the transgene used to produce the tested strains did not incur a fitness cost to the colonies of laboratory-reared *C. hominivorax*. Therefore genetically manipulated strains of NWS may be effective in SIT programs.

Technical Abstract: Over the past ten years genetic transformation of insects has matured as a method to a stage at which realistic expectations concerning implementation of transgenic insect control strategies may now be considered. Sterile insect technique (SIT) programs are designed to eradicate pest species by release of mass-reared, sterile insects into an infested area. The first large-scale implementation of SIT was the New World Screwworm Eradication Program, which successfully eliminated the New World screwworm (NWS), *Cochliomyia hominivorax* (Coquerel) (Diptera: Calliphoridae) from the Continental US, Mexico, and most of Central America. Ionizing radiation is currently used for sterilization, but transgenic insect techniques could hypothetically replace this method, providing a safer, more cost-effective alternative. Genetic transformation methods have recently been demonstrated in NWS. Some key issues involved in mass release of sterile insects revolve around fitness. Transgenic insects must be highly productive in mass rearing, and equally capable of surviving and mating with wild insects in order to be truly effective. Therefore it is critical to analyze fitness characteristics of transgenic insects. Eight transgenic strains of *C. hominivorax* were compared with the wild-type parental laboratory strain (P95) in culture. Measurements of fecundity, fertility, larval biomass productivity, adult emergence, male ratio, and mating

competitiveness were analyzed. The parental strain control colony was sub-cultured and subjected to handling procedures equivalent to transgenic strains for valid comparisons of overall colony fitness. Two of the eight transgenic strains were significantly less fertile than the controls, but most strains were significantly more fecund. None of the transgenic colonies exhibited significantly lower adult fitness characteristics than the control parental colony, although one transgenic strain had a significantly higher ratio of adults emerging from pupae, and four transgenic strains had higher average pupal weight. Males of one transgenic strain were shown to mate with equal frequency when compared to males of the controls. Overall, the presence of the transgene used to produce the tested strains did not incur a fitness cost to the colonies of laboratory-reared *C. hominivorax*.

Last Modified: 08/23/2007