

December 1993

THE BEHLEN OBSERVATORY VARIABLE STAR SURVEY: FINDING CHARTS AND LIGHT CURVES FOR THE FIRST NINETY- THREE STARS

Edward G. Schmidt

University of Nebraska-Lincoln, eschmidt1@unl.edu

Darwin E. Reiswig

University of Nebraska - Lincoln

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsschmidt>



Part of the [Physics Commons](#)

Schmidt, Edward G. and Reiswig, Darwin E., "THE BEHLEN OBSERVATORY VARIABLE STAR SURVEY: FINDING CHARTS AND LIGHT CURVES FOR THE FIRST NINETY-THREE STARS" (1993). *Edward Schmidt Publications*. 11.

<http://digitalcommons.unl.edu/physicsschmidt/11>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Edward Schmidt Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

THE BEHLEN OBSERVATORY VARIABLE STAR SURVEY: FINDING CHARTS AND LIGHT CURVES FOR THE FIRST NINETY-THREE STARS

EDWARD G. SCHMIDT

Behlen Observatory, Department of Physics and Astronomy, University of Nebraska, Lincoln, Nebraska 68588-0111
and The National Science Foundation, 1800 G St. NW, Washington, DC 20550
Electronic mail: eschmidt@unlinfo.unl.edu

DARWIN E. REISWIG

Behlen Observatory, Department of Physics and Astronomy, University of Nebraska, Lincoln, Nebraska 68588-0111
Received 1993 June 28; revised 1993 August 9

ABSTRACT

In a previous paper, photometry for the first ninety-three stars in a photometric survey of variable stars was discussed. Finding charts, accurate coordinates, and light curves for those stars are presented here. The reliability of the coordinates from the *General Catalogue of Variable Stars* is discussed. The magnitude scale of the present photometry is compared with that of the *Hubble Space Telescope Guide Star Catalogue*.

1. INTRODUCTION

Previously (Schmidt 1991, hereafter referred to as Paper I; 1992) we discussed photometry of the first 93 stars observed in the Behlen Observatory variable star survey. Several parameters describing the light curves were presented and the periods and variable star classifications of some stars were rediscussed in light of the new data. Stars with unusual properties were identified.

Characterizing light curves in terms of a few parameters is useful but it is also helpful to examine the curves themselves. Many variables are difficult to identify since finding charts are not always widely available and the coordinates are sometimes inaccurate or erroneous. With modern instruments which point with arcsecond accuracy, coordinates assume even greater importance. To address these needs this paper presents finding charts, accurate coordinates, and plots of the light curves for the stars previously discussed. The original photometric data have been deposited in the IAU Archives of Unpublished Observations of Variable Stars (Breger *et al.* 1990). The archive file numbers are given in Table 1. These data will also appear in the ApJ/AJ CD-ROM series.

Although the *Hubble Space Telescope Guide Star Catalogue* (GSC; described by Lasker *et al.* 1990; Russell *et al.* 1990; Jenkner *et al.* 1990) was generated for a particular purpose, providing guide stars for the *HST*, it is a valuable record of stellar magnitudes at the epoch of the survey plates (mostly 1982 to 1984 for the northern hemisphere). For example, Schmidt *et al.* (1992) used GSC magnitudes to investigate the long term behavior of the variable U Tau. In order to utilize the GSC magnitudes, it is necessary to understand their reliability as well as the relationship of the magnitude system to standard photometric systems. In Sec. 4 we explore these issues using the comparison stars from our variable star photometry.

2. FINDING CHARTS AND IDENTIFICATION

Table 2 lists the stars from Paper I. Column 1 contains the variable star designation while column 2 is the number of the star in the GSC. The first four digits identify the GSC small region within which the star falls and the last four digits identify the star within that region. The lack of such a number indicates that the variable is not present in the GSC. The next two columns give the coordinates for the equinox of 2000. These were obtained from the GSC for variables which were found in the catalogue. For most others, at least one other star in our CCD frame was in the GSC and the coordinates were determined from it. These coordinates are accurate to 1" which is sufficient to uniquely identify the objects.

No GSC star was found in the CCD fields of KV Per and DR Lyr. Their coordinates were determined by measurements on the Palomar Observatory Sky Survey prints relative to nearby GSC stars. The coordinates of these stars are estimated to be accurate to 5" which is still adequate for identification.

Finding charts for the variables were generated from the GSC. However, many of our comparison stars and variables are too faint to be included in the catalogue. In generating the charts we also found that a significant number of stars brighter than the nominal limit for the catalogue are missing. This is not surprising in view of the stated preference of the compilers of the GSC for rejecting dubious images over including spurious objects (Lasker *et al.* 1990). Additionally, the magnitude limit of the catalogue was intentionally set at a brighter level in more crowded fields. When the variable or comparison stars are missing from the GSC we have added them to our charts using coordinates from our CCD frames. Figure 1 displays the finding charts. Each chart shows a region of the sky 10' on a side. The variable is plotted as an open circle when it is brighter than about fourteenth magnitude. Fainter variables are indicated by a solid dot within a circle.

TABLE 1. File numbers in the IAU archives of unpublished observations of variable stars.

File No.	Constellation
251E	And, Aql, Ari, Aur
252E	Cam, Cnc, Cas, Cep, Com, Cyg ^a
253E	Del, Gem, Her
254E	Leo, LMi, Lyr, Mon, Oph, Per, Psc
255E	Ser, Sex, Tau, Tri, UMa, Vir, Vul

Note to TABLE 1

^aThe star labeled CV Cyg in the archives is actually CV Del (see Schmidt 1992) while that labeled V830 Cyg is a newly identified variable which is referred to as Anon Cyg in this paper.

Each chart (except that for Anon Cyg, see below) is centered on the coordinates given in the *General Catalogue of Variable Stars* (GCVS; Kholopov 1985, 1987). As can be seen, most of the stars fall near the center of the chart. When this was not the case and when there were other grounds for suspicion (e.g., our data gave a period which

differed significantly from that in the GCVS) the original finding charts were consulted. With a single exception, the identification was confirmed.

The star we identified as V830 Cyg is not at the GCVS coordinates of that star and the field does not match the finding chart published by Tsevech & Mandel (1963). Clearly we misidentified the star and our comments regarding V830 Cyg in Paper I should be disregarded. Neither the GCVS nor the *Catalogue of Suspected Variables* (Kukarkin 1982) contains a star close to the location of the variable we observed. It is apparently a new discovery and it will be referred to as Anon Cyg in lieu of a variable star designation.

Coordinates in the GCVS are tabulated to 1^s in right ascension and 0.1' in declination. However, only 28% of the GCVS right ascensions agree with ours to within the implied $\pm 0.5^s$ while 19% of the declinations agree to within the implied $\pm 3''$. Only 10% of the stars agree to

TABLE 2. Identification and coordinates of variables.^a

Star	GSC	R.A.	(2000)	Dec	Star	GSC	R.A.	(2000)	Dec
ZZ	And	17420303	0 49 34.9	27 1 19	IL	Her	-----	17 0 48.7	30 14 16
BK	And	32350127	23 35 6.1	41 6 11	LW	Her	20801960	17 41 48.9	25 09 26
CI	And	-----	01 55 8.2	43 45 58	OX	Her	31020835	18 3 26.4	38 41 41
DE	And	-----	23 17 26.2	48 33 7	V392	Her	20822371	17 20 51.5	26 32 20
DY	And	32410213	23 58 42.3	41 29 20	V442	Her	31111372	18 12 58.5	42 3 46
FI	And	22890006	0 56 39.9	37 15 48	V458	Her	15391217	17 8 30.9	18 31 14
GM	And	22670718	0 0 3.6	35 21 46	V469	Her	15441671	17 14 10.7	19 45 29
GV	And	27630880	23 13 12.5	36 54 4	V599	Her	-----	16 36 30.7	26 32 16
HK	And	32620112	0 46 31.9	45 5 47	V677	Her	20350068	16 8 4.2	24 59 20
V793	Aql	05002405	20 15 23.0	3 21 7	V692	Her	20510030	16 22 18.2	26 22 32
SY	Ari	12200669	2 17 34.1	21 42 59	V719	Her	30800343	17 9 52.6	42 56 8
TU	Ari	12201541	2 9 4.3	21 11 28	XX	Leo	08350142	9 59 29.4	13 47 6
BH	Aur	23970244	5 12 4.3	33 57 47	AX	Leo	08590119	11 33 3.8	12 9 14
CR	Aur	29390730	6 28 58.9	43 45 45	BT	Leo	14370360	11 12 4.7	18 30 5
DN	Aur	-----	5 8 0.0	33 23 49	Y	LMi	25131084	10 15 51.5	32 51 34
GT	Aur	-----	5 36 26.9	44 35 29	DR	Lyr	-----	19 2 12.2	26 08 42
MV	Aur	24062093	5 58 12.4	30 29 19	NW	Lyr	26611099	19 15 56.3	34 27 8
TY	Cam	40852781	5 33 26.3	62 29 15	V462	Lyr	31041423	18 28 36.9	38 3 22
AS	Cnc	19441281	8 25 42.1	25 43 9	UW	Mon	48141567	7 3 39.7	-00 11 32
BD	Cas	40140388	0 9 51.4	61 30 51	V518	Mon	-----	6 51 9.1	0 37 26
BF	Cas	40142244	0 14 3.3	60 59 11	V756	Oph	04010572	17 22 30.4	1 46 48
BR	Cas	40380014	1 21 39.8	65 36 51	V768	Oph	04260413	17 31 15.8	5 41 6
NO	Cas	40151454	0 24 4.7	61 20 30	V773	Oph	09960445	17 32 29.7	9 23 38
NY	Cas	36670948	0 40 23.3	58 37 07	V822	Oph	09982401	17 45 59.1	10 30 15
OP	Cas	40200217	0 46 29.9	63 32 36	AN	Per	-----	3 08 31.5	48 32 42
MU	Cep	39900664	22 23 38.7	57 40 52	CI	Per	36931900	2 5 2.2	57 8 35
UZ	Com	25350228	13 12 26.7	30 21 16	ET	Per	36711029	1 39 22.1	53 52 19
BW	Com	14441988	12 4 16.7	18 53 12	GP	Per	28911394	4 23 19.2	44 14 12
CY	Com	19890299	12 28 20.0	24 57 19	HQ	Per	28971485	4 43 57.9	40 50 5
FV	Com	19920696	12 37 57.2	29 58 6	KN	Per	28692543	3 22 35.6	41 19 56
GU	Com	14460093	12 39 3.3	18 18 24	KV	Per	-----	1 58 33.9	57 26 44
V508	Cyg	35731794	20 34 5.9	46 52 17	V375	Per	28651164	3 26 51.3	40 13 1
V742	Cyg	-----	20 2 32.8	37 46 34	V428	Per	28510765	3 2 35.9	41 12 5
Anon	Cyg	31710197	20 55 25.6	39 23 44	SY	Psc	00250346	1 2 32.9	5 23 42
ZZ	Del	16320732	20 31 13.9	15 4 32	DF	Ser	14890371	15 15 19.1	18 39 28
CV	Del	16322095	20 30 54.2	16 32 35	U	Sex	02400493	9 57 25.4	3 40 5
AK	Gem	-----	6 54 59.6	13 42 21	V	Sex	02400749	9 56 7.7	4 51 7
DT	Gem	18681396	6 6 58.1	25 24 2	BR	Tau	12780849	4 34 42.9	21 46 22
EW	Gem	-----	6 44 49.9	24 18 9	CN	Tau	18712093	5 58 9.4	28 2 34
GQ	Gem	-----	7 10 2.8	14 47 4	IY	Tau	18691239	5 42 23.1	27 56 47
KV	Gem	13301213	6 47 12.6	15 43 34	SX	Tri	23360433	2 33 53.1	35 47 45
CW	Her	25970581	16 50 38.3	35 27 5	AP	UMa	30121664	11 10 24.3	42 48 54
DL	Her	09900684	17 20 22.5	14 30 38	BD	UMa	34521543	11 57 6.8	48 24 26
EE	Her	15080531	16 12 42.6	17 59 15	BK	UMa	30111600	10 50 18.9	42 34 8
GY	Her	30631333	16 38 18.1	37 48 4	BV	Vir	02900915	12 46 35.1	2 20 16
HO	Her	25900839	16 57 20.8	30 21 28	AW	Vul	21601036	20 29 1.7	24 48 28
					FH	Vul	16460074	20 40 19.9	22 13 25






FIG. 1. (a)–(h) Finding charts for the variables. Each chart is 10' on a side. North is up and east to the left. The magnitudes of the stars are indicated by the size of the dots according to the legend at the bottom of each panel. The brighter variables are indicated by open circles. For stars fainter than about fourteenth magnitude, the symbols are too small to distinguish from solid circles. In those cases the variable is circled.




(e) 10. ● 12.5 • 15.

(g) 10. ● 12.5 • 15.



(f) 10. ● 12.5 • 15.



(h) 10. ● 12.5 • 15.

FIG. 1. (continued)




FIG. 2. (a)–(x) Light curves for all of the variables. For each variable the upper panel presents the V magnitudes while the $(V-R)$ colors are plotted in the lower panel. The periods and epochs of maximum are from Paper I.




FIG. 2. (continued)




FIG. 2. (continued)




FIG. 2. (continued)




FIG. 2. (continued)




FIG. 2. (continued)




FIG. 3. Residuals between the GSC magnitudes and V magnitudes plotted against various quantities. In panels (a), (b), and (c) the V magnitudes are from the present photometry while those in (d) are from Landolt (1992). In (c) and (d) the solid lines indicate the fitted relationship as discussed in the text.

within these uncertainties in both coordinates. At the other extreme, the GCVS coordinates for four stars are in error by more than 8^s (or $2'$) in right ascension while 16 have declination errors larger than $1'$. We conclude that the GCVS coordinates are not accurate enough to reliably identify moderately faint variables, particularly in Milky Way fields. Lopez & Girard (1990) found a similar situation among southern variables.

3. THE LIGHT AND COLOR CURVES

In Fig. 2 we present the light and color curves. A few were published previously but all are shown here for completeness. The scatter often appears larger in $(V-R)$ than in V . Unlike photoelectric photometry, there is no ten-

dency for errors to cancel when a color is formed from CCD magnitudes. Rather, the scatter in the color increases due to the combination of the errors in two independent exposures. Additionally, we emphasized the form of the light curves in our survey and thus did not always integrate long enough in R to achieve high accuracy. Due to this larger scatter the color curves must be treated with some caution. Nonetheless, they serve to show the relation between color and magnitude and provide an indication of the nature of the stars.

4. THE GSC MAGNITUDE SCALE

There were 155 comparison stars in our program which could be identified with GSC stars. Of these, the GSC

magnitudes of 139 were based on IIaD plates exposed through a Wratten #12 filter. Only these stars will be discussed here so our conclusions pertain only to GSC V_{12} (also denoted as V_1) magnitudes.

In Figs. 3(a)–3(c) we plot the residuals of the GSC magnitudes from our V magnitudes. In Fig. 3(a) the residuals are plotted against the distance from the nearest edge of the GSC plate from which the magnitude was derived. While the average of the residuals does not depend on the distance from the plate edge, the scatter can be seen to increase near the edge. The standard deviation of the residuals is 0.25 mag for stars more than one degree from a plate edge and 0.38 for stars closer than that to an edge. Russell *et al.* (1990, their Fig. 6) show a similar comparison with data from the CCD Transit Instrument of McGraw *et al.* (1986) in their Fig. 6. However, while much of their plot shows residuals comparable with ours, we have not encountered any of the very large residuals they show at some of the plate boundaries.

In Fig. 3(b) we plot the magnitude residuals against our V magnitudes. Both the average of the residuals and the scatter appear to be relatively independent of magnitude. This is borne out by the fact that the mean of the residuals for stars brighter than $V=13$ is -0.18 with an rms scatter of 0.30 while fainter than $V=14$ the mean is -0.22 with an rms scatter of 0.32. Since the magnitudes of the GSC were calibrated with sequences of stars which ranged from ninth to fifteenth magnitude for each plate, this result is not surprising.

Figure 3(c) shows the residuals as a function of $(V-R)$ color. Since the GSC magnitudes are on the natural system of the plates, a color term is expected in this plot and the points clearly slope downward to the right. To better define the color dependence of the V_{12} magnitudes, we located 189 stars in Landolt's (1992) list of standard stars which were far enough north to fall on V_{12} plates and for which the identification with GSC stars was unambiguous. The residuals between the GSC magnitudes and Landolt's V magnitudes are plotted in Fig. 3(d) where the same trend is seen as in Fig. 3(c).

A least-squares fit to the combined data from our comparison stars and Landolt's standards gives the relation

$$m_{\text{GSC}} - V = -0.04 - 0.31(V - R)$$

which is shown by the lines in Figs. 3(c) and 3(d). It can be seen that it represents a satisfactory fit to the individual data sets. The rms scatter about this relation was ± 0.27 mag. This transformation can be used to place GSC magnitudes on the V scale and the rms scatter can be used as an indication of the accuracy of the resulting magnitudes. However, it should be remembered that much larger errors are occasionally encountered in the GSC magnitudes (Russell *et al.* 1990).

Our slope of -0.31 using $(V-R)$ corresponds to a slope of -0.18 in terms of $(B-V)$. This is in satisfactory agreement with the color term of -0.15 which was assumed in the calibration of the V_{12} magnitudes.¹

Nearby comparison stars can generally be used to investigate the long term behavior of a star. Under such a circumstance, the global scatter of the GSC magnitudes is of less interest than the consistency of magnitudes in a small region. From seventeen variable star fields with three or more comparison stars that have V_{12} magnitudes, we find a mean internal standard deviation of 0.20 mag. While this can be used as an estimate of the reliability of the magnitude scale in small areas, we again caution that significantly larger deviations may occur in isolated cases.

The research described here was supported by the National Science Foundation through Grant No. AST-8815806 and a REU supplement to that grant. The CCD camera system at Behlen Observatory was purchased with funds from NSF Grant No. AST-8504072. The reductions and analysis were carried out using the facilities of the Minnich Astronomical Computing Center which was funded by a donation from Commander Charles B. Minnich. The library at the Dominion Astrophysical Observatory was used extensively for this paper. The hospitality of the staff is appreciated.

¹Russell *et al.* (1990) plotted magnitude residuals from their calibration sequences against color (their Fig. 3) and found no color term. They then cited this as evidence that the adopted color transformations were correct. However, this is a consequence of the calibration scheme and says nothing about the validity of the adopted color terms.

REFERENCES

- Breger, M., Jaschek, & C. Dubois, P. 1990, IBVS, 3422
 Jenkner, H., Lasker, B. M., Sturch, C. R., McLean, B. J., Shara, M. M., & Russell, J. L. 1990, AJ, 99, 2082
 Kholopov, P. N. 1985, General Catalogue of Variable Stars, 4th Ed. (Nauka, Moscow), Vols. 1 and 2
 Kholopov, P. N. 1987, General Catalogue of Variable Stars, 4th Ed. (Nauka, Moscow), Vol. 3.
 Kukarkin, B. V. 1982, New Catalogue of Suspected Variable Stars (Nauka, Moscow)
 Landolt, A. U. 1992, AJ, 104, 372
 Lasker, B. M., Sturch, C. R., McLean, B. J., Russell, J. L., Jenkner, H., & Shara, M. M. 1990, AJ, 99, 2019
 Lopez, C. E., & Girard, T. M. 1990, PASP, 102, 1018
 McGraw, J. T., Cawson, M. G. M., & Keane, M. J. 1986, Proc. SPIE, 627, 60
 Russell, J. L., Lasker, B. M., McLean, B. J., Sturch, C. R., & Jenkner, H. 1990, AJ, 99, 2059
 Schmidt, E. G. 1991, AJ, 102, 1766
 Schmidt, E. G. 1992, AJ, 103, 1701
 Schmidt, E. G., Chab, J. R., & Reiswig, D. E. 1992, PASP, 104, 906
 Tsevech, V. P., & Mandel, O. E. 1963, Variable Stars, 14, 324