

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

2006 Bird Strike Committee USA/Canada, 8th  
Annual Meeting, St. Louis, MO

Bird Strike Committee Proceedings

---

August 2006

# PREY MANIPULATION AS A MANAGEMENT STRATEGY AT AN INLAND SOUTH AFRICAN AIRPORT

Ordino Kok

*University of the Free State, Bloemfontein, South Africa*

Follow this and additional works at: <http://digitalcommons.unl.edu/birdstrike2006>



Part of the [Environmental Health and Protection Commons](#)

---

Kok, Ordino , "PREY MANIPULATION AS A MANAGEMENT STRATEGY AT AN INLAND SOUTH AFRICAN AIRPORT"  
(2006). 2006 Bird Strike Committee USA/Canada, 8th Annual Meeting, St. Louis, MO. 16.  
<http://digitalcommons.unl.edu/birdstrike2006/16>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2006 Bird Strike Committee USA/Canada, 8th Annual Meeting, St. Louis, MO by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

**(15) PREY MANIPULATION AS A MANAGEMENT STRATEGY AT AN INLAND SOUTH AFRICAN AIRPORT**

*Ordino Kok, Department of Zoology and Entomology, University of the Free State, P.O.Box 339, Bloemfontein 9300 South Africa*

Prey manipulation was investigated as a means of reducing the bird hazard at the Bloemfontein airport, central South Africa. From January 1985 to December 2005 approximately 7,000 individuals representing 55 bird species which, potentially, posed a threat to aviation, were collected at the airport. Numerically the crowned plover (*Vanellus coronatus*), blacksmith plover (*V. armatus*), lesser kestrel (*Falco naumanni*), white-winged korhaan (*Eupodotis afraoides*) and double-banded courser (*Smutsornis africanus*) in order of importance constituted more than 90% of the total sample. Based on stomach analyses harvester termites (*Hodotermes mossambicus*) comprise the dominant food source of all five species concerned. Experimentally a significant and sustainable decrease in termite numbers and activities in disturbed grass areas could be accomplished by administering bait treated with Gaucho, a systemic insecticide. By using a combination of pitfall traps, mouse traps and sweep nets over a period of 15 months, prey abundance in undisturbed long grass (average 57 cm) and grass which were kept permanently short (average 22 cm) by regular mowing was compared. Terrestrial invertebrates, mainly insects, contributed more than 90% of the overall sample with twice as many individuals in short grass than in long grass. The available food source for the predominantly insectivorous avifauna at the Bloemfontein airport can thus be effectively reduced by implementing a so-called long grass policy in combination with seasonal chemical control of harvester termites, thereby limiting bird numbers and also bird strikes.