

December 1993

EVOLUTIONARY PERSPECTIVES ON ANIMALIVORY

Patricia W. Freeman

University of Nebraska-Lincoln, pfreeman1@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/museummammalogy>



Part of the [Zoology Commons](#)

Freeman, Patricia W., "EVOLUTIONARY PERSPECTIVES ON ANIMALIVORY" (1993). *Mammalogy Papers: University of Nebraska State Museum*. 24.

<http://digitalcommons.unl.edu/museummammalogy/24>

This Article is brought to you for free and open access by the Museum, University of Nebraska State at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Mammalogy Papers: University of Nebraska State Museum by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

From:

Abstracts: Annual Meeting of the American Society of
Zoologists

December 26–30, 1993

Published in American Zoologist 33:5 (1993), p. 102A.

Copyright © 1993 by The American Society of Zoologists

Craniofacial Adaptations of Faunivorous Mammals

410

EVOLUTIONARY PERSPECTIVES ON ANIMALIVORY.
P.W. Freeman, University of Nebraska
State Museum, Lincoln.

Derived from insectivorous ancestors, microchiropteran bats have diversified into a remarkable array of feeding habits. Cranial and dental characteristics mirror this diversity. Insectivorous predators of hard-shelled prey have thicker jaws, well-developed cranial crests, fewer but larger teeth, longer canines and abbreviated M3s than species that take soft items. Carnivorous bats have more elongate skulls, larger brain volumes and larger pinnae, lengthened metastylar ridges, and larger protoconids compared to insectivores. Animalivorous species have large ectoloph areas compared to frugivorous and nectarivorous species, but animalivores and frugivores have large tooth areas relative to palatal area compared to nectarivores. Frugivores sacrifice ectoloph for enlarged crushing surfaces on the upper molars while nectarivores have similar proportions as animalivores but have diminutive teeth, elongated rostra, and highly thegosed upper canines. Omnivores have a more equal allocation to more kinds of teeth on the toothrow than do frugivores or animalivores.