

4-29-2005

# Measurement of the Ratio of Inclusive Cross Sections $\sigma(pp \rightarrow Z + b \text{ jet}) / \sigma(pp \rightarrow Z + \text{jet})$ at $\sqrt{s} = 1.96 \text{ TeV}$

V.M. Abazov

*Joint Institute for Nuclear Research, Dubna, Russia*

Gregory R. Snow

*University of Nebraska-Lincoln, gsnow1@unl.edu*

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicssnow>



Part of the [Physics Commons](#)

---

Abazov, V. M.; Snow, Gregory R.; and Collaboration, D0, "Measurement of the Ratio of Inclusive Cross Sections  $\sigma(pp \rightarrow Z + b \text{ jet}) / \sigma(pp \rightarrow Z + \text{jet})$  at  $\sqrt{s} = 1.96 \text{ TeV}$ " (2005). *Gregory Snow Publications*. 24.  
<http://digitalcommons.unl.edu/physicssnow/24>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Gregory Snow Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Measurement of the Ratio of Inclusive Cross Sections $\sigma(p\bar{p} \rightarrow Z + b \text{ jet})/\sigma(p\bar{p} \rightarrow Z + \text{jet})$ at $\sqrt{s} = 1.96 \text{ TeV}$

V. M. Abazov,<sup>33</sup> B. Abbott,<sup>70</sup> M. Abolins,<sup>61</sup> B. S. Acharya,<sup>27</sup> M. Adams,<sup>48</sup> T. Adams,<sup>46</sup> M. Agelou,<sup>17</sup> J.-L. Agram,<sup>18</sup> S. H. Ahn,<sup>29</sup> M. Ahsan,<sup>55</sup> G. D. Alexeev,<sup>33</sup> G. Alkhalaf,<sup>37</sup> A. Alton,<sup>60</sup> G. Alverson,<sup>59</sup> G. A. Alves,<sup>2</sup> M. Anastasoia,<sup>32</sup> S. Anderson,<sup>42</sup> B. Andrieu,<sup>16</sup> Y. Arnoud,<sup>13</sup> A. Askew,<sup>74</sup> B. Åsman,<sup>38</sup> O. Atramentov,<sup>53</sup> C. Autermann,<sup>20</sup> C. Avila,<sup>7</sup> F. Badaud,<sup>12</sup> A. Baden,<sup>57</sup> B. Baldin,<sup>47</sup> P. W. Balm,<sup>31</sup> S. Banerjee,<sup>27</sup> E. Barberis,<sup>59</sup> P. Bargassa,<sup>74</sup> P. Baringer,<sup>54</sup> C. Barnes,<sup>40</sup> J. Barreto,<sup>2</sup> J. F. Bartlett,<sup>47</sup> U. Bassler,<sup>16</sup> D. Bauer,<sup>51</sup> A. Bean,<sup>54</sup> S. Beauceron,<sup>16</sup> M. Begel,<sup>66</sup> A. Bellavance,<sup>63</sup> S. B. Beri,<sup>26</sup> G. Bernardi,<sup>16</sup> R. Bernhard,<sup>47,\*</sup> I. Bertram,<sup>39</sup> M. Besançon,<sup>17</sup> R. Beuselinck,<sup>40</sup> V. A. Bezzubov,<sup>36</sup> P. C. Bhat,<sup>47</sup> V. Bhatnagar,<sup>26</sup> M. Binder,<sup>24</sup> K. M. Black,<sup>58</sup> I. Blackler,<sup>40</sup> G. Blazey,<sup>49</sup> F. Blekman,<sup>31</sup> S. Blessing,<sup>46</sup> D. Bloch,<sup>18</sup> U. Blumenschein,<sup>22</sup> A. Boehnlein,<sup>47</sup> O. Boeriu,<sup>52</sup> T. A. Bolton,<sup>55</sup> F. Borchering,<sup>47</sup> G. Borissov,<sup>39</sup> K. Bos,<sup>31</sup> T. Bose,<sup>65</sup> A. Brandt,<sup>72</sup> R. Brock,<sup>61</sup> G. Brooijmans,<sup>65</sup> A. Bross,<sup>47</sup> N. J. Buchanan,<sup>46</sup> D. Buchholz,<sup>50</sup> M. Buehler,<sup>48</sup> V. Buescher,<sup>22</sup> S. Burdin,<sup>47</sup> T. H. Burnett,<sup>76</sup> E. Busato,<sup>16</sup> J. M. Butler,<sup>58</sup> J. Bystricky,<sup>17</sup> W. Carvalho,<sup>3</sup> B. C. K. Casey,<sup>71</sup> N. M. Cason,<sup>52</sup> H. Castilla-Valdez,<sup>30</sup> S. Chakrabarti,<sup>27</sup> D. Chakraborty,<sup>49</sup> K. M. Chan,<sup>66</sup> A. Chandra,<sup>27</sup> D. Chapin,<sup>71</sup> F. Charles,<sup>18</sup> E. Cheu,<sup>42</sup> L. Chevalier,<sup>17</sup> D. K. Cho,<sup>66</sup> S. Choi,<sup>45</sup> T. Christiansen,<sup>24</sup> L. Christofek,<sup>54</sup> D. Claes,<sup>63</sup> B. Clément,<sup>18</sup> C. Clément,<sup>38</sup> Y. Coadou,<sup>5</sup> M. Cooke,<sup>74</sup> W. E. Cooper,<sup>47</sup> D. Coppage,<sup>54</sup> M. Corcoran,<sup>74</sup> J. Coss,<sup>19</sup> A. Cothenet,<sup>14</sup> M.-C. Cousinou,<sup>14</sup> S. Crépe-Renaudin,<sup>13</sup> M. Cristetiu,<sup>45</sup> M. A. C. Cummings,<sup>49</sup> D. Cutts,<sup>71</sup> H. da Motta,<sup>2</sup> B. Davies,<sup>39</sup> G. Davies,<sup>40</sup> G. A. Davis,<sup>50</sup> K. De,<sup>72</sup> P. de Jong,<sup>31</sup> S. J. de Jong,<sup>32</sup> E. De La Cruz-Burelo,<sup>30</sup> C. De Oliveira Martins,<sup>3</sup> S. Dean,<sup>41</sup> F. Déliot,<sup>17</sup> P. A. Delsart,<sup>19</sup> M. Demarteau,<sup>47</sup> R. Demina,<sup>66</sup> P. Demine,<sup>17</sup> D. Denisov,<sup>47</sup> S. P. Denisov,<sup>36</sup> S. Desai,<sup>67</sup> H. T. Diehl,<sup>47</sup> M. Diesburg,<sup>47</sup> M. Doidge,<sup>39</sup> H. Dong,<sup>67</sup> S. Doulas,<sup>59</sup> L. Duflot,<sup>15</sup> S. R. Dugad,<sup>27</sup> A. Duperrin,<sup>14</sup> J. Dyer,<sup>61</sup> A. Dyshkant,<sup>49</sup> M. Eads,<sup>49</sup> D. Edmunds,<sup>61</sup> T. Edwards,<sup>41</sup> J. Ellison,<sup>45</sup> J. Elmsheuser,<sup>24</sup> J. T. Eltzroth,<sup>72</sup> V. D. Elvira,<sup>47</sup> S. Eno,<sup>57</sup> P. Ermolov,<sup>35</sup> O. V. Eroshin,<sup>36</sup> J. Estrada,<sup>47</sup> D. Evans,<sup>40</sup> H. Evans,<sup>65</sup> A. Evdokimov,<sup>34</sup> V. N. Evdokimov,<sup>36</sup> J. Fast,<sup>47</sup> S. N. Fatakia,<sup>58</sup> L. Felgioni,<sup>58</sup> T. Ferbel,<sup>66</sup> F. Fiedler,<sup>24</sup> F. Filthaut,<sup>32</sup> W. Fisher,<sup>64</sup> H. E. Fisk,<sup>47</sup> M. Fortner,<sup>49</sup> H. Fox,<sup>22</sup> W. Freeman,<sup>47</sup> S. Fu,<sup>47</sup> S. Fuess,<sup>47</sup> T. Gadfort,<sup>76</sup> C. F. Galea,<sup>32</sup> E. Gallas,<sup>47</sup> E. Galyaev,<sup>52</sup> C. Garcia,<sup>66</sup> A. Garcia-Bellido,<sup>76</sup> J. Gardner,<sup>54</sup> V. Gavrilov,<sup>34</sup> P. Gay,<sup>12</sup> D. Gelé,<sup>18</sup> R. Gelhaus,<sup>45</sup> K. Genser,<sup>47</sup> C. E. Gerber,<sup>48</sup> Y. Gershtein,<sup>71</sup> G. Ginther,<sup>66</sup> T. Golling,<sup>21</sup> B. Gómez,<sup>7</sup> K. Gounder,<sup>47</sup> A. Goussiou,<sup>52</sup> P. D. Grannis,<sup>67</sup> S. Greder,<sup>18</sup> H. Greenlee,<sup>47</sup> Z. D. Greenwood,<sup>56</sup> E. M. Gregores,<sup>4</sup> Ph. Gris,<sup>12</sup> J.-F. Grivaz,<sup>15</sup> L. Groer,<sup>65</sup> S. Grünendahl,<sup>47</sup> M. W. Grünewald,<sup>28</sup> S. N. Gurzhiev,<sup>36</sup> G. Gutierrez,<sup>47</sup> P. Gutierrez,<sup>70</sup> A. Haas,<sup>65</sup> N. J. Hadley,<sup>57</sup> S. Hagopian,<sup>46</sup> I. Hall,<sup>70</sup> R. E. Hall,<sup>44</sup> C. Han,<sup>60</sup> L. Han,<sup>41</sup> K. Hanagaki,<sup>47</sup> K. Harder,<sup>55</sup> R. Harrington,<sup>59</sup> J. M. Hauptman,<sup>53</sup> R. Hauser,<sup>61</sup> J. Hays,<sup>50</sup> T. Hebbeker,<sup>20</sup> D. Hedin,<sup>49</sup> J. M. Heinmiller,<sup>48</sup> A. P. Heinson,<sup>45</sup> U. Heintz,<sup>58</sup> C. Hensel,<sup>54</sup> G. Hesketh,<sup>59</sup> M. D. Hildreth,<sup>52</sup> R. Hirosky,<sup>75</sup> J. D. Hobbs,<sup>67</sup> B. Hoeneisen,<sup>11</sup> M. Hohlfeld,<sup>23</sup> S. J. Hong,<sup>29</sup> R. Hooper,<sup>71</sup> P. Houben,<sup>31</sup> Y. Hu,<sup>67</sup> J. Huang,<sup>51</sup> I. Iashvili,<sup>45</sup> R. Illingworth,<sup>47</sup> A. S. Ito,<sup>47</sup> S. Jabeen,<sup>54</sup> M. Jaffré,<sup>15</sup> S. Jain,<sup>70</sup> V. Jain,<sup>68</sup> K. Jakobs,<sup>22</sup> A. Jenkins,<sup>40</sup> R. Jesik,<sup>40</sup> K. Johns,<sup>42</sup> M. Johnson,<sup>47</sup> A. Jonckheere,<sup>47</sup> P. Jonsson,<sup>40</sup> H. Jöstlein,<sup>47</sup> A. Juste,<sup>47</sup> M. M. Kado,<sup>43</sup> D. Käfer,<sup>20</sup> W. Kahl,<sup>55</sup> S. Kahn,<sup>68</sup> E. Kajfasz,<sup>14</sup> A. M. Kalinin,<sup>33</sup> J. Kalk,<sup>61</sup> D. Karmanov,<sup>35</sup> J. Kasper,<sup>58</sup> D. Kau,<sup>46</sup> R. Kehoe,<sup>73</sup> S. Kermiche,<sup>14</sup> S. Kesisoglou,<sup>71</sup> A. Khanov,<sup>66</sup> A. Kharchilava,<sup>52</sup> Y. M. Kharzheev,<sup>33</sup> K. H. Kim,<sup>29</sup> B. Klima,<sup>47</sup> M. Klute,<sup>21</sup> J. M. Kohli,<sup>26</sup> M. Kopal,<sup>70</sup> V. M. Korablev,<sup>36</sup> J. Kotcher,<sup>68</sup> B. Kothari,<sup>65</sup> A. Koubarovsky,<sup>35</sup> A. V. Kozelov,<sup>36</sup> J. Kozminski,<sup>61</sup> S. Krzywdzinski,<sup>47</sup> S. Kulshov,<sup>34</sup> Y. Kulik,<sup>47</sup> S. Kunori,<sup>57</sup> A. Kupco,<sup>17</sup> T. Kurča,<sup>19</sup> S. Lager,<sup>38</sup> N. Lahrichi,<sup>17</sup> G. Landsberg,<sup>71</sup> J. Lazoflores,<sup>46</sup> A.-C. Le Bihan,<sup>18</sup> P. Lebrun,<sup>19</sup> S. W. Lee,<sup>29</sup> W. M. Lee,<sup>46</sup> A. Leflat,<sup>35</sup> F. Lehner,<sup>47,\*</sup> C. Leonidopoulos,<sup>65</sup> P. Lewis,<sup>40</sup> J. Li,<sup>72</sup> Q. Z. Li,<sup>47</sup> J. G. R. Lima,<sup>49</sup> D. Lincoln,<sup>47</sup> S. L. Linn,<sup>46</sup> J. Linnemann,<sup>61</sup> V. V. Lipaev,<sup>36</sup> R. Lipton,<sup>47</sup> L. Lobo,<sup>40</sup> A. Lobodenko,<sup>37</sup> M. Lokajicek,<sup>10</sup> A. Lounis,<sup>18</sup> H. J. Lubatti,<sup>76</sup> L. Lueking,<sup>47</sup> M. Lynker,<sup>52</sup> A. L. Lyon,<sup>47</sup> A. K. A. Maciel,<sup>49</sup> R. J. Madaras,<sup>43</sup> P. Mättig,<sup>25</sup> A. Magerkurth,<sup>60</sup> A.-M. Magnan,<sup>13</sup> N. Makovec,<sup>15</sup> P. K. Mal,<sup>27</sup> S. Malik,<sup>56</sup> V. L. Malyshev,<sup>33</sup> H. S. Mao,<sup>6</sup> Y. Maravin,<sup>47</sup> M. Martens,<sup>47</sup> S. E. K. Mattingly,<sup>71</sup> A. A. Mayorov,<sup>36</sup> R. McCarthy,<sup>67</sup> R. McCroskey,<sup>42</sup> D. Meder,<sup>23</sup> H. L. Melanson,<sup>47</sup> A. Melnitchouk,<sup>62</sup> M. Merkin,<sup>35</sup> K. W. Merritt,<sup>47</sup> A. Meyer,<sup>20</sup> H. Miettinen,<sup>74</sup> D. Mihalcea,<sup>49</sup> J. Mitrevski,<sup>65</sup> N. Mokhov,<sup>47</sup> J. Molina,<sup>3</sup> N. K. Mondal,<sup>27</sup> H. E. Montgomery,<sup>47</sup> R. W. Moore,<sup>50</sup> G. S. Muanza,<sup>19</sup> M. Mulders,<sup>47</sup> Y. D. Mutaf,<sup>67</sup> E. Nagy,<sup>14</sup> M. Narain,<sup>58</sup> N. A. Naumann,<sup>32</sup> H. A. Neal,<sup>60</sup> J. P. Negret,<sup>7</sup> S. Nelson,<sup>46</sup> P. Neustroev,<sup>37</sup> C. Noeding,<sup>22</sup> A. Nomerotski,<sup>47</sup> S. F. Novaes,<sup>4</sup> T. Nunnemann,<sup>24</sup> E. Nurse,<sup>41</sup> V. O'Dell,<sup>47</sup> D. C. O'Neil,<sup>5</sup> V. Oguri,<sup>3</sup> N. Oliveira,<sup>3</sup> N. Oshima,<sup>47</sup> G. J. Otero y Garzón,<sup>48</sup> P. Padley,<sup>74</sup> N. Parashar,<sup>56</sup> J. Park,<sup>29</sup> S. K. Park,<sup>29</sup> J. Parsons,<sup>65</sup> R. Partridge,<sup>71</sup> N. Parua,<sup>67</sup> A. Patwa,<sup>68</sup> P. M. Perea,<sup>45</sup> E. Perez,<sup>17</sup> O. Peters,<sup>31</sup> P. Pétrouff,<sup>15</sup> M. Petteni,<sup>40</sup> L. Phaf,<sup>31</sup> R. Piegaia,<sup>1</sup> P. L. M. Podesta-Lerma,<sup>30</sup> V. M. Podstavkov,<sup>47</sup> Y. Pogorelov,<sup>52</sup> B. G. Pope,<sup>61</sup> W. L. Prado da Silva,<sup>3</sup> H. B. Prosper,<sup>46</sup> S. Protopopescu,<sup>68</sup> M. B. Przybycien,<sup>50,†</sup> J. Qian,<sup>60</sup>

A. Quadt,<sup>21</sup> B. Quinn,<sup>62</sup> K. J. Rani,<sup>27</sup> P. A. Rapidis,<sup>47</sup> P. N. Ratoff,<sup>39</sup> N. W. Reay,<sup>55</sup> S. Reucroft,<sup>59</sup> M. Rijssenbeek,<sup>67</sup> I. Ripp-Baudot,<sup>18</sup> F. Rizatdinova,<sup>55</sup> C. Royon,<sup>17</sup> P. Rubinov,<sup>47</sup> R. Ruchti,<sup>52</sup> G. Sajot,<sup>13</sup> A. Sánchez-Hernández,<sup>30</sup> M. P. Sanders,<sup>41</sup> A. Santoro,<sup>3</sup> G. Savage,<sup>47</sup> L. Sawyer,<sup>56</sup> T. Scanlon,<sup>40</sup> R. D. Schamberger,<sup>67</sup> H. Schellman,<sup>50</sup> P. Schieferdecker,<sup>24</sup> C. Schmitt,<sup>25</sup> A. A. Schukin,<sup>36</sup> A. Schwartzman,<sup>64</sup> R. Schwienhorst,<sup>61</sup> S. Sengupta,<sup>46</sup> H. Severini,<sup>70</sup> E. Shabalina,<sup>48</sup> M. Shamim,<sup>55</sup> V. Shary,<sup>17</sup> W. D. Shephard,<sup>52</sup> D. Shpakov,<sup>59</sup> R. A. Sidwell,<sup>55</sup> V. Simak,<sup>9</sup> V. Sirotenko,<sup>47</sup> P. Skubic,<sup>70</sup> P. Slattery,<sup>66</sup> R. P. Smith,<sup>47</sup> K. Smolek,<sup>9</sup> G. R. Snow,<sup>63</sup> J. Snow,<sup>69</sup> S. Snyder,<sup>68</sup> S. Söldner-Rembold,<sup>41</sup> X. Song,<sup>49</sup> Y. Song,<sup>72</sup> L. Sonnenschein,<sup>58</sup> A. Sopczak,<sup>39</sup> M. Sosebee,<sup>72</sup> K. Soustruznik,<sup>8</sup> M. Souza,<sup>2</sup> B. Spurlock,<sup>72</sup> N. R. Stanton,<sup>55</sup> J. Stark,<sup>13</sup> J. Steele,<sup>56</sup> G. Steinbrück,<sup>65</sup> K. Stevenson,<sup>51</sup> V. Stolin,<sup>34</sup> A. Stone,<sup>48</sup> D. A. Stoyanova,<sup>36</sup> J. Strandberg,<sup>38</sup> M. A. Strang,<sup>72</sup> M. Strauss,<sup>70</sup> R. Ströhmer,<sup>24</sup> M. Strovink,<sup>43</sup> L. Stutte,<sup>47</sup> S. Sumowidagdo,<sup>46</sup> A. Sznajder,<sup>3</sup> M. Talby,<sup>14</sup> P. Tamburello,<sup>42</sup> W. Taylor,<sup>5</sup> P. Telford,<sup>41</sup> J. Temple,<sup>42</sup> S. Tentindo-Repond,<sup>46</sup> E. Thomas,<sup>14</sup> B. Thooris,<sup>17</sup> M. Tomoto,<sup>47</sup> T. Toole,<sup>57</sup> J. Torborg,<sup>52</sup> S. Towers,<sup>67</sup> T. Trefzger,<sup>23</sup> S. Trincaz-Duvoid,<sup>16</sup> B. Tuchming,<sup>17</sup> C. Tully,<sup>64</sup> A. S. Turcot,<sup>68</sup> P. M. Tuts,<sup>65</sup> L. Uvarov,<sup>37</sup> S. Uvarov,<sup>37</sup> S. Uzunyan,<sup>49</sup> B. Vachon,<sup>5</sup> R. Van Kooten,<sup>51</sup> W. M. van Leeuwen,<sup>31</sup> N. Varelas,<sup>48</sup> E. W. Varnes,<sup>42</sup> I. A. Vasilyev,<sup>36</sup> M. Vaupel,<sup>25</sup> P. Verdier,<sup>15</sup> L. S. Vertogradov,<sup>33</sup> M. Verzocchi,<sup>57</sup> F. Villeneuve-Seguiet,<sup>40</sup> J.-R. Vlimant,<sup>16</sup> E. Von Toerne,<sup>55</sup> M. Vreeswijk,<sup>31</sup> T. Vu Anh,<sup>15</sup> H. D. Wahl,<sup>46</sup> R. Walker,<sup>40</sup> L. Wang,<sup>57</sup> Z.-M. Wang,<sup>67</sup> J. Warchol,<sup>52</sup> M. Warsinsky,<sup>21</sup> G. Watts,<sup>76</sup> M. Wayne,<sup>52</sup> M. Weber,<sup>47</sup> H. Weerts,<sup>61</sup> M. Wegner,<sup>20</sup> N. Wermes,<sup>21</sup> A. White,<sup>72</sup> V. White,<sup>47</sup> D. Whiteson,<sup>43</sup> D. Wicke,<sup>47</sup> D. A. Wijngaarden,<sup>32</sup> G. W. Wilson,<sup>54</sup> S. J. Wimpenny,<sup>45</sup> J. Wittlin,<sup>58</sup> M. Wobisch,<sup>47</sup> J. Womersley,<sup>47</sup> D. R. Wood,<sup>59</sup> T. R. Wyatt,<sup>41</sup> Q. Xu,<sup>6</sup> N. Xuan,<sup>52</sup> R. Yamada,<sup>47</sup> M. Yan,<sup>57</sup> T. Yasuda,<sup>47</sup> Y. A. Yatsunenko,<sup>33</sup> Y. Yen,<sup>25</sup> K. Yip,<sup>68</sup> S. W. Youn,<sup>50</sup> J. Yu,<sup>72</sup> A. Yurkewicz,<sup>61</sup> A. Zabi,<sup>15</sup> A. Zatserklyaniy,<sup>49</sup> M. Zdrzil,<sup>67</sup> C. Zeitnitz,<sup>23</sup> D. Zhang,<sup>47</sup> X. Zhang,<sup>70</sup> T. Zhao,<sup>76</sup> Z. Zhao,<sup>60</sup> B. Zhou,<sup>60</sup> J. Zhu,<sup>57</sup> M. Zielinski,<sup>66</sup> D. Zieminska,<sup>51</sup> A. Zieminski,<sup>51</sup> R. Zitoun,<sup>67</sup> V. Zutshi,<sup>49</sup> E. G. Zverev,<sup>35</sup> and A. Zylberstein<sup>17</sup>

(D0 Collaboration)

<sup>1</sup>Universidad de Buenos Aires, Buenos Aires, Argentina<sup>2</sup>LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil<sup>3</sup>Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil<sup>4</sup>Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil<sup>5</sup>Simon Fraser University, Burnaby, Canada, University of Alberta, Edmonton, Canada, McGill University, Montreal, Canada, and York University, Toronto, Canada<sup>6</sup>Institute of High Energy Physics, Beijing, People's Republic of China<sup>7</sup>Universidad de los Andes, Bogotá, Colombia<sup>8</sup>Charles University, Center for Particle Physics, Prague, Czech Republic<sup>9</sup>Czech Technical University, Prague, Czech Republic<sup>10</sup>Institute of Physics, Academy of Sciences, Center for Particle Physics, Prague, Czech Republic<sup>11</sup>Universidad San Francisco de Quito, Quito, Ecuador<sup>12</sup>Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France<sup>13</sup>Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble I, Grenoble, France<sup>14</sup>CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France<sup>15</sup>Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS, Orsay, France<sup>16</sup>LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France<sup>17</sup>DAPNIA/Service de Physique des Particules, CEA, Saclay, France<sup>18</sup>IReS, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France, and Université de Haute Alsace, Mulhouse, France<sup>19</sup>Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France<sup>20</sup>III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany<sup>21</sup>Physikalisches Institut, Universität Bonn, Bonn, Germany<sup>22</sup>Physikalisches Institut, Universität Freiburg, Freiburg, Germany<sup>23</sup>Institut für Physik, Universität Mainz, Mainz, Germany<sup>24</sup>Ludwig-Maximilians-Universität München, München, Germany<sup>25</sup>Fachbereich Physik, University of Wuppertal, Wuppertal, Germany<sup>26</sup>Panjab University, Chandigarh, India<sup>27</sup>Tata Institute of Fundamental Research, Mumbai, India<sup>28</sup>University College Dublin, Dublin, Ireland<sup>29</sup>Korea Detector Laboratory, Korea University, Seoul, Korea<sup>30</sup>CINVESTAV, Mexico City, Mexico<sup>31</sup>FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands<sup>32</sup>University of Nijmegen/NIKHEF, Nijmegen, The Netherlands

- <sup>33</sup>Joint Institute for Nuclear Research, Dubna, Russia  
<sup>34</sup>Institute for Theoretical and Experimental Physics, Moscow, Russia  
<sup>35</sup>Moscow State University, Moscow, Russia  
<sup>36</sup>Institute for High Energy Physics, Protvino, Russia  
<sup>37</sup>Petersburg Nuclear Physics Institute, St. Petersburg, Russia  
<sup>38</sup>Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden  
<sup>39</sup>Lancaster University, Lancaster, United Kingdom  
<sup>40</sup>Imperial College, London, United Kingdom  
<sup>41</sup>University of Manchester, Manchester, United Kingdom  
<sup>42</sup>University of Arizona, Tucson, Arizona 85721, USA  
<sup>43</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA  
<sup>44</sup>California State University, Fresno, California 93740, USA  
<sup>45</sup>University of California, Riverside, California 92521, USA  
<sup>46</sup>Florida State University, Tallahassee, Florida 32306, USA  
<sup>47</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA  
<sup>48</sup>University of Illinois at Chicago, Chicago, Illinois 60607, USA  
<sup>49</sup>Northern Illinois University, DeKalb, Illinois 60115, USA  
<sup>50</sup>Northwestern University, Evanston, Illinois 60208, USA  
<sup>51</sup>Indiana University, Bloomington, Indiana 47405, USA  
<sup>52</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA  
<sup>53</sup>Iowa State University, Ames, Iowa 50011, USA  
<sup>54</sup>University of Kansas, Lawrence, Kansas 66045, USA  
<sup>55</sup>Kansas State University, Manhattan, Kansas 66506, USA  
<sup>56</sup>Louisiana Tech University, Ruston, Louisiana 71272, USA  
<sup>57</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>58</sup>Boston University, Boston, Massachusetts 02215, USA  
<sup>59</sup>Northeastern University, Boston, Massachusetts 02115, USA  
<sup>60</sup>University of Michigan, Ann Arbor, Michigan 48109, USA  
<sup>61</sup>Michigan State University, East Lansing, Michigan 48824, USA  
<sup>62</sup>University of Mississippi, University, Mississippi 38677, USA  
<sup>63</sup>University of Nebraska, Lincoln, Nebraska 68588, USA  
<sup>64</sup>Princeton University, Princeton, New Jersey 08544, USA  
<sup>65</sup>Columbia University, New York, New York 10027, USA  
<sup>66</sup>University of Rochester, Rochester, New York 14627, USA  
<sup>67</sup>State University of New York, Stony Brook, New York 11794, USA  
<sup>68</sup>Brookhaven National Laboratory, Upton, New York 11973, USA  
<sup>69</sup>Langston University, Langston, Oklahoma 73050, USA  
<sup>70</sup>University of Oklahoma, Norman, Oklahoma 73019, USA  
<sup>71</sup>Brown University, Providence, Rhode Island 02912, USA  
<sup>72</sup>University of Texas, Arlington, Texas 76019, USA  
<sup>73</sup>Southern Methodist University, Dallas, Texas 75275, USA  
<sup>74</sup>Rice University, Houston, Texas 77005, USA  
<sup>75</sup>University of Virginia, Charlottesville, Virginia 22901, USA  
<sup>76</sup>University of Washington, Seattle, Washington 98195, USA

(Received 27 October 2004; published 27 April 2005)

Using the data collected with the D0 detector at  $\sqrt{s} = 1.96$  TeV, for integrated luminosities of about  $180 \text{ pb}^{-1}$ , we have measured the ratio of inclusive cross sections for  $p\bar{p} \rightarrow Z + b$  jet to  $p\bar{p} \rightarrow Z + \text{jet}$  production. The inclusive  $Z + b$ -jet reaction is an important background to searches for the Higgs boson in associated  $ZH$  production at the Fermilab Tevatron collider. Our measurement is the first of its kind, and relies on the  $Z \rightarrow e^+e^-$  and  $Z \rightarrow \mu^+\mu^-$  modes. The combined measurement of the ratio yields  $0.021 \pm 0.005$  for hadronic jets with transverse momenta  $p_T > 20 \text{ GeV}/c$  and pseudorapidities  $|\eta| < 2.5$ , consistent with next-to-leading-order predictions of the standard model.

DOI: 10.1103/PhysRevLett.94.161801

PACS numbers: 14.70.Hp, 13.85.Ni, 13.85.Qk, 14.65.Fy

Inclusive  $Z + b$ -jet production is expected to be a major background to Higgs-boson production in the  $p\bar{p} \rightarrow ZH$  channel, with subsequent Higgs-boson decays into  $b\bar{b}$ . The

parton-level subprocesses expected to contribute to the  $Z + b$ -jet final state are  $bg \rightarrow Zb$  (where  $g$  stands for a gluon) and  $q\bar{q} \rightarrow Zg$ , with  $g \rightarrow b\bar{b}$  [1]. The process

$bg \rightarrow Zb$ , where the initial  $b$  is from the sea of the proton parton distribution, is predicted to account for approximately two-thirds of the total inclusive cross section  $\sigma(p\bar{p} \rightarrow Z + b \text{ jet})$  at  $\sqrt{s} = 1.96$  TeV. The  $b$ -quark density of the proton influences the production rates of single top quarks and the final state  $hb$ , with  $h$  representing a supersymmetric Higgs boson. Consequently, the measurement of  $Z + b$ -jet production is an important step in constraining the  $b$ -quark density of protons.

In this Letter, we describe a measurement of the ratio of production cross sections of inclusive  $Z + b$  jets to  $Z +$  jets. The measurement of the ratio benefits from cancellations of many systematic uncertainties, such as the 6.5% uncertainty in the luminosity, and therefore allows a more precise comparison with theory.

We search for  $Z$  bosons in association with hadronic jets in about  $180 \text{ pb}^{-1}$  of data collected at the D0 experiment between August 2002 and September 2003. The D0 detector at the Fermilab Tevatron collider is a general-purpose detector comprising a magnetic central-tracking, pre-shower, calorimeter, and muon systems [2]. The central-tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker, both located within a 2 T superconducting solenoidal magnet. The design was optimized for tracking and vertexing capabilities at pseudorapidities  $|\eta| < 3$ , where  $\eta = -\ln[\tan(\theta/2)]$  and  $\theta$  is the polar angle with respect to the proton beam direction ( $z$ ). Particle energies are measured in three liquid-argon/uranium calorimeters with coverage to  $|\eta| < 4.2$  [3]. The muon detection system is outside the calorimetry and consists of a layer of tracking detectors and scintillation trigger counters before 1.8 T iron toroid magnets, followed by two similar layers after the toroids. The trigger and data acquisition systems are designed to accommodate high luminosities.

The dielectron sample is selected by requiring two clusters of energy in the electromagnetic (EM) layers at the trigger level. In the offline selection, two EM clusters are each required to have transverse momentum  $p_T > 15 \text{ GeV}/c$  and  $|\eta| < 2.5$ . In addition, the shower development in the calorimeter and isolation from hadronic activity must be consistent with that expected of an electron, and at least one of the EM clusters is required to have an associated track to maximize the possibility of having a  $Z$  boson in the event. The electron candidates with matching tracks are required to have a ratio of measured energy in the calorimeter to momentum measured with the tracking system consistent with that expected of an electron. The  $Z$  candidates are selected by requiring a dielectron mass ( $m_{ee}$ ) of  $80 \text{ GeV}/c^2 < m_{ee} < 100 \text{ GeV}/c^2$ . The  $Z +$  jet sample is then selected by requiring the presence of at least one reconstructed hadronic jet with  $p_T > 20 \text{ GeV}/c$  and  $|\eta| < 2.5$ .

Jets are reconstructed from calorimeter clusters using a cone algorithm of cone size  $\Delta\mathcal{R} = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} =$

$0.5$  in pseudorapidity and azimuth ( $\phi$ ). A hadronic jet is called “taggable” if it is associated with a cluster of tracks (track jet) within  $\Delta\mathcal{R} < 0.5$ . The taggability requirement reduces background from noise in the calorimeter. Track jets are found by applying a cone track clustering algorithm of size  $\Delta\mathcal{R} = 0.5$ , with a seed track of  $p_T > 1.0 \text{ GeV}/c$ , to tracks of  $p_T > 0.5 \text{ GeV}/c$  that are close to the primary interaction vertex (whose determination is discussed below). A track jet can consist of two or more tracks.

Applying the taggability criterion to 2661 jets in inclusive 2219  $Z(ee) +$  jet candidate events in the mass  $Z$  window yields 1658 events. Based on sidebands to the  $Z$  mass window,  $121 \pm 4$  events are estimated to be from background sources. The main background is from multijet production where two jets mimic EM objects, with one of the objects having an overlapping track that passes the track-matching criteria. The taggability per jet is  $(75 \pm 1)\%$ , after background subtraction.

The dimuon sample is defined by the detection of at least one muon candidate at the trigger level. In the offline selection, two isolated muons are required to be of opposite charge, and have  $p_T > 15 \text{ GeV}/c$  and  $|\eta| < 2$ , with trajectories in the muon spectrometer matched to tracks in the central-tracking detector. Muon isolation is based on the transverse component of the muon momentum relative to the combined momenta of the muon and the closest calorimeter jet in  $(\eta, \phi)$  space, and requires  $p_{T\text{rel}} > 10 \text{ GeV}/c$ . The  $Z$  candidates are selected by requiring a dimuon mass of  $65 \text{ GeV}/c^2 < m_{\mu\mu} < 115 \text{ GeV}/c^2$ . The  $Z$  mass window is larger than in the dielectron channel due to worse momentum resolution for high  $p_T$  muons. The criteria for reconstructed hadronic jets are the same as in the dielectron channel. A total of 1406 events out of 1754 inclusive  $Z(\mu\mu) +$  jet candidate events remain after the requirement that there be at least one taggable jet. The main background in this channel is from  $b\bar{b}$  production, where both  $b$  jets contain muons that satisfy the isolation criterion (referred to as  $b\bar{b}$  background). The isolation efficiencies of muons from  $Z$  and  $b\bar{b}$  are expected to be different, since, for the latter, a hadronic jet would be expected to be close to the muon. By performing fits to dimuon mass spectra, where the background contributes to the continuum, samples with different numbers of isolated muons are analyzed to measure the isolation efficiencies and background rates. From such analyses, we estimate that the background contribution to the final sample with two isolated muons is  $17.5 \pm 4.1$  events.

Figure 1 shows distributions in transverse momentum of taggable jets for both channels (points with error bars), compared to a  $Z +$  jet Monte Carlo (MC) calculation generated with ALPGEN [4], using PYTHIA [5] for parton showering and hadronization. Also shown is a background estimation based on data obtained from samples that are in the sideband for the dielectron channel, or fail the isolation criterion for the dimuon channel. The background distri-




FIG. 1 (color online). The  $p_T$  distribution of taggable jets in dielectron and dimuon channels compared to  $Z + \text{jet}$  ALPGEN with PYTHIA showering and full detector simulation (open histogram), and background (multijet for  $ee$  channel and  $b\bar{b}$  background for  $\mu\mu$  channel) derived from data. The error bars on the data points are statistical. The prediction is normalized to the data, as described in the text.

bution is normalized to the number of background events estimated in the selected sample. The simulated signal is then normalized so that the total agrees with the measurement in Fig. 1. Within the uncertainty of the jet energy scale (JES), indicated by the darker shading about the expectation, the shape of the distribution is well described by the simulation.

The  $b$  quarks fragment into  $B$  hadrons, which are identified by displaced secondary vertices that are separated from the primary vertex. In the high luminosity environment of the Fermilab Tevatron collider, there can be more than one interaction per beam crossing, one of which is likely to have triggered the recorded event. It is possible to distinguish the main hard-interaction vertex that produces the  $Z$  boson from any additional soft interactions because the vertices are normally well separated along  $z$ . Primary interaction vertices are reconstructed in two passes. In the first pass, all tracks present in an event are used to find seed vertices through an iterative method, where tracks that contribute to a fit to a common vertex with a  $\chi^2/\text{d.o.f.}$  greater than some chosen threshold are removed. The fit is repeated until a stable set of seeds is obtained. The seed vertices are then used in a second pass to fit all tracks within a certain distance of closest approach to any seed. This improves the position resolution on the vertex, since the fit is less affected by poorly reconstructed tracks. The

$p_T$  distribution of the associated tracks is then used to calculate a probability for the vertex to be consistent with that of a soft interaction. The vertex that has the smallest probability is selected as the primary interaction vertex (PV).

A  $b$ -jet tagging algorithm for secondary vertices (SV) is used to identify heavy-quark jets in the analysis. Tracks that are displaced from the PV in the transverse plane are used as seeds to find secondary vertices. First, a fixed-cone jet algorithm of  $\Delta R = 0.5$  is used to cluster the tracks to form track jets. Tracks are required to have hits in at least two layers of the SMT,  $p_T > 0.5 \text{ GeV}/c$ , and be within 0.40 cm in  $z$  relative to the PV. Tracks identified as arising from  $K_S^0$  and  $\Lambda$  decays or photon conversions are not considered. Any pair of tracks within a track jet with an impact parameter relative to the hard-interaction vertex (distance of closest approach,  $d_{ca}$ , of a track to a vertex in the plane transverse to the  $z$  direction) divided by its uncertainty ( $\sigma_{d_{ca}}$ ) of  $d_{ca}/\sigma_{d_{ca}} > 3$  is used as a seed for secondary vertices. Additional tracks are attached iteratively to the seed vertices if their  $\chi^2$  contribution to the vertex fit is consistent with originating from the vertex. A jet is considered  $b$  tagged when it is taggable and has at least one secondary vertex, with a decay length in a plane transverse to the beam line ( $L_{xy}$ ) divided by its uncertainty  $L_{xy}/\sigma_{xy} > 7$ , associated with it. A secondary vertex is associated to a jet if the opening angle between the direction of the calorimeter-based jet axis and the momentum vector of the SV is  $\Delta R < 0.5$ .

The  $b$ -tagging efficiency ( $\epsilon_b$ ) and the light-flavor tagging rate ( $\epsilon_\ell$ ) of the  $b$ -tagging algorithm are parametrized as functions of jet  $p_T$  and  $\eta$ . The parametrization of  $\epsilon_b$  is derived from a different data sample using events with jets containing muons (muonic jets), which are dominated by  $b$  jets, but also have contributions from light quark jets, gluon jets, and charm jets. The  $b$ -tagging efficiency is extracted from the heavy-flavor component in this muonic jet sample. The light-flavor tagging rate is also derived from data, after compensating for effects of displaced vertices that do not originate from heavy-flavor decay ( $K_S^0$ ,  $\Lambda$ , and photon conversions). Different types of samples are used to determine  $\epsilon_\ell$  and  $\epsilon_b$ , and the spreads are taken as systematic uncertainties.

A comparison of inclusive  $Z + \text{jet}$  events, generated with the ALPGEN leading-order matrix element and PYTHIA for showering, with inclusive  $Z + b$  events generated with PYTHIA, shows good agreement for jet  $p_T$  and  $\eta$  distributions. We therefore use the shapes of  $p_T$  and  $\eta$  derived from the  $Z + \text{jet}$  data sample to estimate the expected  $b$ -tagging efficiency and the light-flavor tagging (“mistag”) rate. The average  $b$ -tagging efficiency and the mistag rate per jet, averaged over  $p_T$  and  $\eta$ , are found to be  $(32.8 \pm 1.3)\%$  and  $(0.25 \pm 0.02)\%$ , respectively, for the dielectron channel. Corresponding values for the dimuon channel are  $(33.1 \pm 1.1)\%$  and  $(0.24 \pm 0.02)\%$ . To

obtain the event mistag rate, we take into consideration jet multiplicity and measure the event mistag rate of 0.28% (0.27%) for the dielectron (dimuon) channel.

Since  $\epsilon_b$  is derived from events with a muon embedded in a jet, whereas most of the  $b$ -tagged jets do not contain such muons, the difference in  $b$ -tagging efficiencies for hadronic  $b$  jets and muonic  $b$  jets is derived from MC calculations, and the ratio is used to correct  $\epsilon_b$ . We cannot at this point derive the charm tagging efficiency ( $\epsilon_c$ ) from data, so we rely on PYTHIA MC calculations to compare  $Z \rightarrow b\bar{b}$  and  $Z \rightarrow c\bar{c}$  samples. We assume that  $(\epsilon_c/\epsilon_b)_{\text{data}} = (\epsilon_c/\epsilon_b)_{\text{MC}} = 0.266 \pm 0.003$ .

The jet taggability for light jets,  $t_\ell$ , is measured using data to be  $(75 \pm 1)\%$ , while that for  $b$  jets,  $t_b$ , is obtained from MC calculations, and scaled such that  $(t_b)_{\text{data}} = (t_\ell)_{\text{data}} \times (t_b/t_\ell)_{\text{MC}}$ . The result is  $(t_b)_{\text{data}} = (79.2 \pm 1.3)\%$  for the dielectron channel and  $(80.7 \pm 1.1)\%$  for the dimuon channel. We assume that the taggability of charm jets is same as  $t_b$ .

After applying  $b$  tagging, 27  $Z(\rightarrow ee) + b$ -jet candidate events are left, with an expected background from the Drell-Yan  $ee$  continuum and multijet background of  $4.2 \pm 1.4$  events based on the sideband subtraction method. In the dimuon channel, 22 events are observed with  $5.0 \pm 1.1$  events from  $b\bar{b}$  background.

After subtracting the background contributions, two equations, one before and the other after the requirement of  $b$  tagging, determine the contributions from different flavors in the remaining events:

$$N_{\text{before } b\text{-tag}} = t'_b N_b + t'_c N_c + t'_\ell N_\ell \quad (1)$$

$$N_{b\text{-tagged}} = \bar{\epsilon}_b t'_b N_b + \bar{\epsilon}_c t'_c N_c + \bar{\epsilon}_\ell t'_\ell N_\ell, \quad (2)$$

where  $N_b$ ,  $N_c$ , and  $N_\ell$  are the number of events with  $b$ ,  $c$ , and light jets, respectively;  $t'_i$  are the taggabilities per event for different jet types; and the  $\bar{\epsilon}_i$  are the corresponding mean event-tagging efficiencies. In considering event taggability  $t'_b$  ( $t'_c$ ) and tagging efficiency  $\epsilon_b$  ( $\epsilon_c$ ), we take into account the enhancement due to the presence of two heavy-flavor jets in  $Z + b\bar{b}$  ( $Z + c\bar{c}$ ). The event taggability is 1.02 times the jet taggability, and  $b$  and  $c$  event-tagging efficiency is 1.06 times the jet tagging efficiency. Equations (1) and (2) have three unknowns. We take the theoretical prediction of  $N_c = 1.69N_b$  [1] to provide a solution to Eqs. (1) and (2) for  $N_b$ ,  $N_c$ , and  $N_\ell$ .

The ratio  $\sigma(p\bar{p} \rightarrow Z + b \text{ jet})/\sigma(p\bar{p} \rightarrow Z + \text{jet}) = N_b/(N_b + N_c + N_\ell)$  is  $0.023 \pm 0.007$  for the dielectron channel and  $0.019 \pm 0.005$  for the dimuon channel, where the errors are purely statistical. The combined ratio, using the statistical weighting of the number of observed  $Z + \text{jet}$  candidates, is  $0.021 \pm 0.004$ . The shape of the  $p_T$  spectrum for  $b$ -tagged jets and the significance of decay lengths of secondary vertices are compared to the sum of background and  $Z + b$  MC calculations in Fig. 2. The contribution of each component is given by the solution to Eqs. (1) and (2).




FIG. 2 (color online). (a) The  $p_T$  spectrum for  $b$ -tagged jets. (b) Distribution in the transverse plane of decay-length significance of secondary vertices in the transverse plane, without the requirement of decay-length significance. All error bars are statistical.

The distribution of the decay-length significance for secondary vertices shows clear evidence for a heavy-flavor component in the  $b$ -tagged candidate events.

Sources of systematic uncertainty in the ratio include the following: (i) The jet energy scale. The JES is varied within its uncertainty and the differences observed between light-flavored jets and  $b$  jets are included in the uncertainty. (ii) Different methods of estimating background. The background is varied by its measured uncertainty, and the ratio is recalculated. (iii) Events with two heavy quarks. A  $b\bar{b}$  or  $c\bar{c}$  pair from gluon splitting can be present either inside a jet [ $Z + (Q\bar{Q})$ ] or form two separate jets [ $Z + Q\bar{Q}$ ]. For the former, the increase in tagging probability and uncertainty in theoretical cross section and, for the latter, theoretical uncertainty [1] contribute to the sources of error. (iv) Mistag rate for light jets, which depends on the type of jet sample. Using events collected from hadronic jet triggers, the light-jet tagging efficiency is measured to be 0.23%, and for a sample of events with an enhanced EM fraction and small imbalance in overall  $p_T$ , this is 0.26%. (v) Uncertainty in tagging efficiency for  $b$  and  $c$  jets is obtained by varying the efficiency by  $\pm 1$  standard deviation, assuming complete correlation in the ratio of extracted cross sections. Also, for  $c$  jets, there is additional uncertainty from the  $\epsilon_c/\epsilon_b$  ratio obtained from MC calculations. (vi) A small difference observed in  $t_b/t_\ell$  for different MC samples of  $Z + b \text{ jet}/Z + \text{light jet}$ , and  $Z \rightarrow b\bar{b}/Z \rightarrow q\bar{q}$  is taken into account. (vii) Differences in tagging efficiency between hadronic jets and those containing muons. The  $b$ -tagging efficiency is measured in data using muonic jets. The tagging efficiency for hadronic jets is estimated to be 86% of that of muonic jets, as derived from  $Z \rightarrow b\bar{b}$  MC calculations. The same ratio in  $Z + b\bar{b}$  MC calculations is measured to be 84%, and the difference of 2% is taken as a systematic uncertainty. (viii) Different  $p_T$  dependence in jet reconstruction for light,  $b$ , and  $c$  jets, measured using MC samples, is taken as a systematic uncertainty. (ix) Uncertainty from theory

TABLE I. Systematic uncertainties for the combined ratio of cross sections, showing the impact of  $\pm 1$  standard deviation changes in contributions.

Source	Upward (%)	Downward (%)
Jet energy scale	5.8	6.9
Background estimate	5.7	5.2
$Z + (Q\bar{Q})$ and $Z + Q\bar{Q}$	1.7	5.4
Mistag rate	3.4	3.2
$b/c$ tagging efficiency	3.3	2.7
Taggability	1.8	1.8
Correction for hadronic jet	1.7	1.9
Jet reconstruction efficiency	1.7	1.9
$\sigma(Z + c)/\sigma(Z + b)$	2.8	2.8
Total (added in quadrature)	10.4	11.8

for the ratio  $\sigma(Z + c \text{ jet})/\sigma(Z + b \text{ jet}) = N_c/N_b$  is estimated as 9.5% [1].

The effects of systematic uncertainties on the combined measurement are listed in Table I. All these uncertainties are assumed to be completely correlated for the two channels, except for that due to background estimation. Folding these uncertainties together, yields a ratio of  $0.021 \pm 0.004(\text{stat})_{-0.003}^{+0.002}(\text{syst})$ . This is in good agreement with the next-to-leading-order (NLO) prediction of  $0.018 \pm 0.004$  [1,6] using the CTEQ6 parton distributions [7].

In summary, we have presented the first inclusive measurement of  $b$ -jet production in association with  $Z$  bosons at the Tevatron collider. This is a background for the standard-model Higgs boson in the  $ZH$  production channel. This measurement is the first direct experimental probe into the  $b$ -quark density and is in agreement with the NLO calculations using CTEQ6 parton densities. In the future, with reduced experimental and theoretical uncertainties, the measurement will provide additional constraint on the  $b$ -quark density of protons.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the Department of

Energy and the National Science Foundation (USA), Commissariat à l'Energie Atomique and CNRS/Institut National de Physique Nucléaire et de Physique des Particules (France), Ministry of Education and Science, Agency for Atomic Energy, and RF President Grants Program (Russia), CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil), Department of Atomic Energy and Department Science and Technology (India), Colciencias (Colombia), CONACyT (Mexico), KRF (Korea), CONICET and UBACyT (Argentina), the Foundation for Fundamental Research on Matter (The Netherlands), PPARC (United Kingdom), Ministry of Education (Czech Republic), Natural Sciences and Engineering Research Council and WestGrid Project (Canada), BMBF and DFG (Germany), A.P. Sloan Foundation, Research Corporation, Texas Advanced Research Program, and the Alexander von Humboldt Foundation.

---

\*Visiting from the University of Zurich, Zurich, Switzerland.

†Visiting from the Institute of Nuclear Physics, Krakow, Poland.

- [1] J.M. Campbell, R.K. Ellis, F. Maltoni, and S. Willenbrock, Phys. Rev. D **69**, 074021 (2004).
- [2] D0 Collaboration, V. Abazov *et al.* (to be published); T. LeCompte and H. T. Diehl, Annu. Rev. Nucl. Part. Sci. **50**, 71 (2000).
- [3] S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
- [4] M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. Polosa, J. High Energy Phys. 07 (2003) 001.
- [5] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [6] J.M. Campbell and S. Willenbrock (private communication).
- [7] J. Pumplin *et al.*, J. High Energy Phys. 07 (2002) 012.