

2009

Diseases of Broadleaf Trees

Mark Harrell

Nebraska Forest Service, University of Nebraska

Rachel Allison

Nebraska Forest Service, University of Nebraska

Laurie Stepanek

Nebraska Forest Service, University of Nebraska

Follow this and additional works at: <http://digitalcommons.unl.edu/nebforestpubs>



Part of the [Forest Sciences Commons](#)

Harrell, Mark; Allison, Rachel; and Stepanek, Laurie, "Diseases of Broadleaf Trees" (2009). *Publications, etc. -- Nebraska Forest Service*. 28.

<http://digitalcommons.unl.edu/nebforestpubs/28>

This Article is brought to you for free and open access by the Nebraska Forest Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications, etc. -- Nebraska Forest Service by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Verticillium wilt



Identification

- On maples and other broadleaf trees.
- Branches die.
- Wood has dark green to dark brown streaks or mottled areas.

Control

- Chemical controls are not effective.
- Remove the tree or parts of the tree as they die.
- DO NOT use chips made from branches or trees killed by Verticillium wilt because of the high risk of spreading the fungal pathogen through the chips.
- Plant trees known to be resistant.
- Improve tree health by mulching and watering about 1 inch per week. Avoid overwatering.

Bur oak decline



Identification

- On bur oak.
- In areas that were once native forests but are now parks, residential areas or other developed landscapes or near roads or field edges.
- Branches generally die back from the ends.
- Trees become more susceptible to insects and diseases and may die.

Control

- Improve tree health by mulching with wood or bark chips and watering about 1 inch per week. Avoid overwatering.

Cankers



Identification

- On cottonwood, honeylocust, willow, elm and many other broadleaf trees.
- Dead areas of bark appear, often around pruning cuts or where branches are attached.

Control

- Chemical controls are not effective.
- Improve tree health by mulching and watering.
- Avoid winter pruning or other injuries that weaken the tree or create wounds through which canker fungi can enter.

*** Trade names are examples of available products. No endorsement is implied. Always follow pesticide label instructions.**

Mark Harrell, Rachel Allison, and Laurie Stepanek
Nebraska Forest Service, University of Nebraska
More information: www.nfs.unl.edu

Photo credits: Dutch elm disease #1: Joseph O'Brien, USDA-Forest Service, www.forestryimages.org
Ash rust: Iowa St. Univ. Plant Disease Clinic




The University of Nebraska-Lincoln does not discriminate based on gender, age, disability, race, color, religion, marital status, veteran's status, national or ethnic origin, or sexual orientation.



Diseases of Broadleaf Trees

Anthracnose



Identification

- On sycamore, ash, maple, oak and walnut.
- Worse in years with wet springs.
- Leaf spots and blotches appear in the spring (later for walnut), often following leaf veins.
- Leaves drop early.
- Dead shoots may appear on sycamore and oak.

Control

- For ash, maple and walnut, a control is usually not needed. Trees typically recover.
- For sycamore and oak, a control may be desired if shoot death is severe.
- Spray foliage at budbreak with chlorothalonil (Daconil, Fung-onil), thiophanate-methyl (3336, Fungo) or copper fungicide (Camelot, Bordeaux mixture)* and repeat 2-3 times every 7-14 days.
- Improve tree health by mulching and watering about 1 inch per week. Avoid overwatering.

Apple scab



Identification

- On crabapple and apple.
- Dark spots appear in leaves, often with feathery margins and along leaf veins.
- Leaves turn yellow and drop from the tree.

Control

- Spray with chlorothalonil (Daconil, Fung-onil, Ortho Garden Disease Control), thiophanate-methyl (3336, Fungo), myclobutanil (Immunox, Eagle), mancozeb (Dithane, Fore, Protect), or propiconazole (Banner Maxx, Infuse)* at 7 to 14-day intervals from prebloom (April) through rainy periods.
- Some chemicals cannot be used on trees grown for fruit production.

Cedar-apple rust



Identification

- On crabapple, apple, juniper and redcedar.
- Orange spots appear on the upper leaf surface.
- Raised spots appear on the lower leaf surface.
- Ball-shaped growths (galls) or orange gelatinous masses appear on juniper and redcedar.

Control

- Spray with chlorothalonil (Daconil, Fung-onil, Ortho Garden Disease Control), thiophanate-methyl (3336, Fungo), myclobutanil (Immunox, Eagle), mancozeb (Dithane, Fore, Protect), or propiconazole (Banner Maxx, Infuse)* as flower buds break, at petal drop and 3 or 4 additional times at 7 to 10-day intervals.
- Some chemicals cannot be used on trees grown for fruit production.

Ash rust

Identification

- On all common ash species
- Swollen areas with many circular orange spots on leaves, petioles and green twigs



Control

- Spray with myclobutanil (Immunox, Eagle) at budbreak and repeat 2 to 3 times at 10 to 14-day intervals.

Fire blight

Identification

- Mostly on apple, crabapple, pear, and mountain-ash.
- Shoots and leaves droop and turn dark.



Control

- Prune out dead branches 8-12 inches down from diseased tissue.
- Sterilize pruning tools after each cut with a 70% alcohol solution.
- Spray with streptomycin (Agri-Mycin)* at pink stage (3 to 4 days before blossoms open—usually mid April) and every 5 to 7 days until petal drop.

Dutch elm disease

Identification

- On American elm.
- Leaves turn yellow, then brown.
- Branches die, then the whole tree dies.
- Brown to black streaks appear in the wood.



Control

- Trunk inject with thiabendazole (Arbotect) or propiconazole (Alamo)* before or as symptoms begin appearing (if 5% or less of the crown is affected).

Oak wilt



Identification

- Mostly on red and bur oak.
- Red oaks often die within 2 to 6 weeks.
- Bur oaks decline and may die, especially if stressed by changes in site conditions.
- Leaves turn brown.
- Branches die, then whole tree dies.
- Brown to black streaks appear in the wood.

Control

- For red oak, trunk inject with propiconazole (Alamo)* as soon as symptoms begin appearing or before.
- For bur oak, improve tree health by mulching with wood or bark chips, watering about 1 inch per week, avoiding overwatering and by trunk injecting with propiconazole (Alamo),* if advised, to quicken the recovery.

Tubakia leaf spot

Identification

- Mostly on bur oak.
- Brown blotches on leaves, often along veins.
- Affected leaves may drop from the tree.
- Young shoots may die.
- Symptoms are more extensive in lower branches than in higher branches.



Control

- Control is rarely needed.
- If needed, spray with mancozeb (Dithane, Fore) or propiconazole (Banner Maxx, Infuse)* at budbreak (April) and repeat 2 times at 10 to 14-day intervals.