

2003

Spatial and Temporal Spread of Bovine Tuberculosis in Wild White-Tailed Deer in Michigan

Graham J. Hickling

Stephen M. Schmitt

Daniel J. O'Brien

Follow this and additional works at: <http://digitalcommons.unl.edu/michbovinetb>



Part of the [Veterinary Medicine Commons](#)

Hickling, Graham J.; Schmitt, Stephen M.; and O'Brien, Daniel J., "Spatial and Temporal Spread of Bovine Tuberculosis in Wild White-Tailed Deer in Michigan" (2003). *Michigan Bovine Tuberculosis Bibliography and Database*. 64.
<http://digitalcommons.unl.edu/michbovinetb/64>

This Article is brought to you for free and open access by the Wildlife Disease and Zoonotics at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Michigan Bovine Tuberculosis Bibliography and Database by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

HICKLING, Graham J., Stephen M. SCHMITT and Daniel J. O'BRIEN

Department of Fisheries and Wildlife, Michigan State University, East Lansing, Michigan 48824-1222 (GJH); Michigan Department of Natural Resources, Rose Lake Wildlife Disease Laboratory, 8562 East Stoll Road, East Lansing, Michigan 48823-9421 (SMS, DJO).

SPATIAL AND TEMPORAL SPREAD OF BOVINE TUBERCULOSIS IN WILD WHITE-TAILED DEER IN MICHIGAN

In 1975, a wild white-tailed deer infected with bovine tuberculosis was shot in the northeastern Lower Peninsula, Michigan. The shooting of a second infected deer in the same area in 1994 triggered ongoing disease surveillance in the region. By 2002, bovine tuberculosis had been confirmed in 12 Michigan counties: from 449 deer; two elk; 41 non-cervid wildlife; one captive cervid facility and 28 cattle herds. We analyzed geographic spread of disease since the surveillance began and investigated factors influencing the prevalence of disease within the infected area. These analyses reveal that 78 percent of tuberculous deer came from within a 1560 km² 'core' area, within which the prevalence of apparent disease averaged 2.5 percent. Prevalence declined dramatically outside of the core and was an order of magnitude lower 30km from its boundary. This prevalence gradient was highly significant ($P < 0.0001$) and did not alter over the 6year surveillance period ($P = 0.98$). Within the core, deer density and supplemental feeding by hunters were positively and independently correlated with tuberculosis prevalence in deer. Together, these two factors explained 55 percent of the variation in prevalence. We conclude that bovine tuberculosis was already well established in the deer population in 1994, that the infected area has not expanded significantly since that time, and that deer over-abundance and food supplementation have both contributed to ongoing transmission of disease. Managers are currently enforcing prohibitions on deer feeding in the core and are working to lower deer numbers there through increased hunting pressure.