

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

2002 Bird Strike Committee-USA/Canada, 4th  
Annual Meeting, Sacramento, CA

Bird Strike Committee Proceedings

---

10-24-2002

## Evaluation of Electrobraided Fencing as a Deer Barrier

Thomas W. Seamans

USDA/APHIS/WS National Wildlife Research Center, thomas.w.seamans@aphis.usda.gov

Zachary J. Patton

USDA/APHIS/WS National Wildlife Research Center

Kurt .. VerCauteren

USDA/APHIS/WS National Wildlife Research Center, kurt.c.vercauteren@aphis.usda.gov

Follow this and additional works at: <https://digitalcommons.unl.edu/birdstrike2002>



Part of the [Environmental Health and Protection Commons](#)

---

Seamans, Thomas W.; Patton, Zachary J.; and VerCauteren, Kurt .., "Evaluation of Electrobraided Fencing as a Deer Barrier" (2002).  
2002 Bird Strike Committee-USA/Canada, 4th Annual Meeting, Sacramento, CA. 43.  
<https://digitalcommons.unl.edu/birdstrike2002/43>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2002 Bird Strike Committee-USA/Canada, 4th Annual Meeting, Sacramento, CA by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

### **Evaluation of Electrobraided Fencing as a Deer Barrier**

*Thomas W. Seamans, and Zachary J. Patton, USDA, National Wildlife Research Center, Ohio Field Station, 6100 Columbus Avenue, Sandusky, OH 44870 USA*

*Kurt C. VerCauteren, USDA, National Wildlife Research Center, 4101 LaPorte Avenue, Fort Collins, CO 80521 USA*

Increasing white-tailed deer (*Odocoileus virginianus*) populations in North America have led to direct threats to public safety as well as agricultural losses. Fencing is often used to keep deer from causing damage at both airports and agricultural areas. Tall, chain-link fences have been used successfully but are often prohibitively expensive. Electric fences have potential to offer a less expensive alternative. We tested a new electric fence design marketed under the name ElectroBraid. This fence, comprised of 0.6-cm polyester rope with copper wire woven into it, is carried on frangible, fiberglass posts set at 15-m intervals. From January to March 2002 we conducted both 1- and 2-choice tests on free-ranging deer in northern Ohio. We measured deer intrusions and corn consumption at 10 pairs of fenced sites with and without electricity. Mean deer intrusions at treated sites in both 1- and 2-choice tests were < 1/day while control site intrusions were 84-86/day. Mean corn consumption by all wildlife (e.g., deer, raccoons [*Procyon lotor*], fox squirrels [*Sciurus niger*]) differed between treated (< 2 kg/day) and control sites (15 kg/day). Based upon the results of this test and the cost of ElectroBraid we conclude that this fence, under the conditions of this 5-week test, was an effective and economical deer barrier.