

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Investigations of the Ichthyofauna of Nicaraguan  
Lakes

Papers in the Biological Sciences

---

1976

## Hemoglobin Stability in Bull Sharks

Jack D. Burke

*Medical College of Virginia*

Follow this and additional works at: <http://digitalcommons.unl.edu/ichthynicar>



Part of the [Aquaculture and Fisheries Commons](#)

---

Burke, Jack D., "Hemoglobin Stability in Bull Sharks" (1976). *Investigations of the Ichthyofauna of Nicaraguan Lakes*. 44.  
<http://digitalcommons.unl.edu/ichthynicar/44>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Investigations of the Ichthyofauna of Nicaraguan Lakes by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Reprinted from THE AMERICAN JOURNAL OF ANATOMY  
Vol. 139, No. 3, March 1974 © The Wistar Institute Press 1974

## RAPID COMMUNICATION

## ZIPGRAM

### HEMOGLOBIN STABILITY IN BULL SHARKS (1)

JACK D. BURKE  
Department of Anatomy, Medical College of Virginia,  
Health Sciences Division, Virginia Commonwealth  
University, Richmond, Virginia 23298

**ABSTRACT** No significant differences in oxyhemoglobin affinity, or electrophoretic patterns of hemoglobin were found in 18 bull sharks collected in selected regions of Lake Nicaragua, the Rio San Juan, and the Caribbean Sea. The half-saturation of hemoglobin with oxygen ( $P_{50}$ ) was 11 and 17 mm Hg at pH 7.4 and 6.8, respectively (25° C, 3% hemoglobin solution, potassium phosphate buffer, 0.3 ionic strength). Electrophoresis resolved the hemoglobin into a minor and a major band. Planimetry of densitometric recordings showed that the major band constituted 54% of the total hemoglobin; the minor band, 46%. On the basis of these hemoglobin studies, no subspeciation of bull sharks in Lake Nicaragua was identified, although marine bull sharks have free access to the lake and have been there, at least, since 1535; the synonymy of Carcharhinus nicaraguensis with C. leucas was confirmed.

Lake Nicaragua is the largest lake in Central America. It is approximately 100 miles long, 45 miles wide, and its deepest point is 250 feet. The lake is drained into the Caribbean Sea by the 110 mile-long Rio San Juan which contains eight major rapids. In 1535 de Oviedo recorded sharks present in Lake Nicaragua. Gill and Bransford (1877) described the shark in Lake Nicaragua as being a landlocked, distinct species, Eulamia nicaraguensis. Jordan (1887) renamed the lake shark Carcharhinus nicaraguensis. Beginning with Gill (1893), all of the succeeding reports, check-lists, and catalogues appearing until 1961 referred to the lake shark by one or the other of these two scientific names (Burke, in press). After a morphological study of four preserved lake sharks (Bigelow and Schroeder, '48), and one specimen and a jaw, C. nicaraguensis was placed in synonymy with Carcharhinus leucas, the marine bull shark (Bigelow and Schroeder, '61). The

synonymy received support from Thorson et al. ('66) whose measurements on bull sharks from Lake Nicaragua and the Rio San Juan compared favorably with similar data obtained from body measurements made on pelagic bull sharks (Schwartz, '60; Clark and von Schmidt, '65). In relation to synonymy, it was of interest to investigate hemoglobin, a stable genetic trait, in bull sharks taken from both fresh and salt water. If no variation occurs in bull shark hemoglobin (1) synonymy, based on morphometric data, is supported, and (2) non-variable hemoglobin would indicate no drift toward subspeciation and reproductive isolation (Sick, '61; Gorman and Dessauer, '65; Mourant, '70).

**MATERIALS AND METHODS** Eighteen mature male and female bull sharks were captured: eight in selected regions of the lake, four in the river, four in the river estuaries, and two in the sea. After immobilizing a bull shark by head blows, blood samples were removed from the heart (Burke, '62) using vacutainers, (1mg/ml EDTA) and 1½ inch, 20-gauge needles. The vacutainers, containing blood samples, were packaged in ice, and air-mailed to MCV where hemoglobin solutions were prepared (Dementi and Burke, '72) for oxyhemoglobin affinity curves (Burke, '66) and starch-gel electrophoresis (Smithies, '59). A portion of the hemoglobin solution used in determining an oxyhemoglobin affinity curve was electrophoresed using borate buffer at pH 8.7. At times it was possible to prepare hemoglobin solutions in Nicaragua, and ship them in containers with dry ice. Hemoglobin analyses were performed within 48-72 hr from the time of blood collection.

**RESULTS** The oxygen pressure at which 50% of the hemoglobin of a fresh water bull shark is saturated ( $P_{50}$ ) is 11 mm Hg at pH 7.4; at pH 6.8, the  $P_{50}$  is 17 mm Hg; for a marine bull shark the respective values vary less than one mm Hg at half-saturation (fig. 1). Duplicate analyses at both pH 7.4 and pH 6.8 were made on bull sharks from the lake, river estuaries, and sea. There was

no significant difference between the  $P_{50}$  determined for the lake bull shark group and the  $P_{50}$  of each of the other groups. There was no sexual dimorphism.

Upon electrophoresis, hemoglobin from each bull shark was resolved into two bands identified by benzidine stain. A minor band migrated 16% faster than a major band. Planimetry of densitometer recordings (at 560  $m\mu$ ) of the two bands revealed that the major band constituted 54% of the total hemoglobin and the minor one, 46%.

DISCUSSION Polymorphic hemoglobins are characteristic of fishes (Buhler and Shanks, '59; Chandrasekhar, '59; Manwell et al., '63; Burke, '65; Wilkins and Iles, '66; Riggs, '70). Hemoglobin is such a stable genetic character that it has been used to identify subspecific populations which are isolated, reproductively, within a species (Sick, '61; Gorman and Dessauer, '65; Mourant, '70). The constancy of the  $P_{50}$  and the stable hemoglobin pattern found in each bull shark indicates the homogeneity of bull shark hemoglobin and supports the synonymy of Carcharhinus nicaraguensis with C. leucas. Carcharhinus species do migrate into lakes and rivers from the ocean around the world (Burke, in press), but there is no direct evidence that C. leucas breeds in fresh water (Jensen, '72). Although both anadromous and catadromous migration of C. leucas occurs between Lake Nicaragua and the Caribbean Sea via the Rio San Juan (Thorson, '71), the non-variability of hemoglobin in C. leucas reported here indicates no drift toward subspeciation, and supports the vigorous contention of Herre ('55) that "sharks do not breed in freshwater, but ----return to the sea to breed".

ACKNOWLEDGMENTS I thank Mrs. Jewell Burke and Dr. F. M. Bush for laboratory aid; Sra. Melida C. de Sansón-Román, Dr. José Castillo, Dr. Guillermo Lugo, Sr. Manuel Pilarte, and Sr. Alejandro Argüello for transportation aid in Nicaragua; Dr. E. S. Kline, Dr. Peter Jezyk, Dr. Robert Ellis, Sr. Aldo Zepeda,

Sr. César Arburola, and Messrs. Kurt Koenig, Tom Emerson, Gary Fleming, and Tom Beatty for assistance in collecting sharks and blood samples.


#### LITERATURE CITED

- Bigelow, H. B., and W. C. Schroeder 1948 Sharks. In: Fishes of the Western North Atlantic. Mem. Sears Fd. Marine Res., Yale Univ., New Haven, Connecticut 1: 59-576.
- Bigelow, H. B., and W. C. Schroeder 1961 Carcharhinus nicaraguensis, a synonym of the bull shark, C. leucas. Copeia, 1961: 359.
- Buhler, D. R., and W. E. Shanks 1959 Multiple hemoglobins in fishes. Science, 129: 899-900.
- Burke, J. D. 1962 A simple technique for immobilizing fish to remove blood. Copeia, 1962: 852-854.
- Burke, J. D. 1965 Oxygen affinities and electrophoretic patterns of hemoglobins in trout and bass from Virginia. Med. Coll. Va. Quarterly, 1: 16-21.
- Burke, J. D. 1966 A simple rapid method for determining oxyhemoglobin affinity: Illustration using blood from the Rhesus monkey. Med. Coll. Va. Quarterly, 2: 219-221.
- Burke, J. D. The freshwater shark in Nicaragua. Natl. Geogr. Scient. Repts., in press.
- Chandrasekhar, N. 1959 Multiple haemoglobins in fish. Nature, 184: 1652-1653.
- Clark, E., and K. von Schmidt 1965 Sharks of the Central Gulf Coast of Florida. Bull. Marine Sci., 15: 13-83.
- Dementi, P. L., and J. D. Burke 1972 Oxyhemoglobin affinity in the armadillo. Am. J. Anat., 134: 509-514.
- de Oviedo, G. F. 1535 Historia general de las Indias. Sevilla, Spain, Part I, 1535. Chapters 1-13, Book 42, Histoire du Nicaragua. Translation: H. Ternaux-Compans. A Bertrand, Paris. 1840.
- Herre, A. W. C. T. 1955 Sharks in Fresh water. Science, 122: 417.
- Gill, T. 1893 Sharks in fresh water. Science (Old Series), 22: 165.
- Gill, T., and J. F. Bransford 1877 Synopsis of the fishes of Lake Nicaragua. Proc. Acad. Nat. Sci., 1877: 175-191.
- Gorman, G. C., and H. C. Dessauer 1965 Hemoglobin and transferrin electrophoresis and relationships of island populations of Anolis lizards. Science, 150: 1454-1455.
- Jensen, N. H. 1972 The reproduction and development of the bull shark, Carcharhinus leucas, in the Lake Nicaragua - Rio San Juan system. Dissert. Abst. Inter., 33: 2861-B.
- Jordan, D. S. 1887 A preliminary list of the fishes of the West Indies. Proc. U.S. National Museum, 9: 554-608.
- Manwell, C., C. A. Baker and W. Childress 1963 The genetics of hemoglobin in hybrids. Comp. Biochem. Physiol., 10: 103-120.

---

#### FIGURE LEGEND

- 1 Oxyhemoglobin affinity curves at  $25 \pm 1^{\circ}$  C for Carcharhinus leucas from both fresh and salt water. At pH 7.4,  $P_{50}$  is 11 mm Hg; at pH 6.8  $P_{50}$  is 17 mm Hg.



- Mourant, A. E. 1970 Fish stocks: Biochemical and serological identification. *Science*, 167: 1760-1761.
- Riggs, A. 1970 Properties of fish hemoglobins. In: *Fish Physiology*, W. S. Hoar and D. J. Randall, eds., Academic Press, Vol. IV., Chapt. 6, pp. 209-252.
- Schwartz, F. J. 1960 Additional comments on adult bull sharks Carcharhinus leucas (Müller and Henle) from Chesapeake Bay, Maryland. *Chesapeake Sci.*, 1: 68-71.
- Sick, K. 1961 Hemoglobin polymorphism in fishes. *Nature*, 192: 894-896.
- Smithies, O. 1959 Zone electrophoresis in starch gels and its application to studies of serum proteins. *Adv. Protein Chem.*, 14: 65-113.
- Thorson, T. B. 1971 Movement of bull sharks, Carcharhinus leucas, between Caribbean Sea and Lake Nicaragua demonstrated by tagging. *Copeia*, 1971: 336-338.
- Thorson, T. B., D. E. Watson and C. M. Cowan 1966 The status of the freshwater shark of Lake Nicaragua. *Copeia*, 1966: 385-402.
- Wilkins, N. P., and T. D. Iles 1966 Haemoglobin polymorphism and its ontogeny in herring (Clupea harengus) and sprat (Sprattus sprattus). *Comp. Biochem. Physiol.*, 17: 1141-1158.

#### REFERENCES

- 1 This work was supported initially by the National Geographic Society, Grant No. 826, and aided subsequently by the A. D. Williams Research Committee (MCV), the MCV Foundation (VCU), and the Virginia Academy of Science.