

1-1-1986

Bacteroids Are Stable during Dark-Induced Senescence of Soybean Root Nodules

Gautam Sarath

University of Nebraska-Lincoln, Gautam.sarath@ars.usda.gov

Nancy E. Pfeiffer

University of Nebraska-Lincoln

C.S. Sodhi

USAID/GUYANA, Agency for International Development, Washington, DC

Fred W. Wagner

University of Nebraska-Lincoln, fwagner2@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaarsfacpub>

 Part of the [Agricultural Science Commons](#)

Sarath, Gautam; Pfeiffer, Nancy E.; Sodhi, C.S.; and Wagner, Fred W., "Bacteroids Are Stable during Dark-Induced Senescence of Soybean Root Nodules" (1986). *Publications from USDA-ARS / UNL Faculty*. 49.
<http://digitalcommons.unl.edu/usdaarsfacpub/49>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Bacteroids Are Stable during Dark-Induced Senescence of Soybean Root Nodules^{1,2}

Gautam Sarath, Nancy E. Pfeiffer³, C. S. Sodhi⁴ and Fred W. Wagner

Department of Agricultural Biochemistry, University of Nebraska-Lincoln, Lincoln, Nebraska 68583-0718

Physiological and biochemical markers of metabolic competence were assayed in bacteroids isolated from root nodules of control, dark-stressed, and recovered plants of *Glycine max* Merr. cv 'Woodworth.' Nitrogenase-dependent acetylene reduction by the whole plant decreased to 8% of control rates after 4 days of dark stress and could not be detected in plants dark stressed for 8 days. However, in bacteroids isolated anaerobically, almost 50% of initial acetylene reduction activity remained after 4 days of dark stress but was totally lost after 8 days of dark stress. Bacteroid acetylene reduction activity recovered faster than whole plant acetylene reduction activity when plants were dark stressed for 8 days and returned to a normal light regimen. Significant changes were not measured in bacteroid respiration, protein content, sodium dodecyl sulfate-polyacrylamide gel electrophoresis protein profiles, or in bacteroid proteolytic activity throughout the experiment. Immunoblots of bacteroid extracts revealed the presence of nitrogenase component II in control, 4-day dark-stressed, and 8-day dark-stressed plants that were allowed to recover under a normal light regimen, but not in 8-day dark-stressed plants. Our data indicate that dark stress does not greatly affect bacteroid metabolism or induce bacteroid senescence.

³ Present address: Department of Chemistry, University of Nebraska-Lincoln, Lincoln, NE 68588-0304.

⁴ Present address: USAID/GUYANA, Agency for International Development, Washington, DC 20523.

Link to full text:

<http://www.plantphysiol.org/cgi/content/abstract/82/2/346>