

September 2001

Membrane glycoprotein gp130 of *Dictyostelium discoideum* is lipid-linked and its fate altered in the presence of tunicamycin

Betsy L. Barent
University of Nebraska - Lincoln

Catherine P. Chia
University of Nebraska - Lincoln, cchia1@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscimicro>



Part of the [Microbiology Commons](#)

Barent, Betsy L. and Chia, Catherine P., "Membrane glycoprotein gp130 of *Dictyostelium discoideum* is lipid-linked and its fate altered in the presence of tunicamycin" (2001). *Papers in Microbiology*. 65.
<http://digitalcommons.unl.edu/bioscimicro/65>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Barent, B. L. and C. P. Chia 2001.

Membrane glycoprotein gp130 of *Dictyostelium discoideum* is lipid-linked and its fate altered in the presence of tunicamycin. Session # 1110.

Published in Abstracts: 41st American Society for Cell Biology Annual Meeting, in *Molecular Biology of the Cell* **12** (supplement): p. 204a. Copyright © 2001 American Society for Cell Biology. Used by permission.

Membrane glycoprotein gp130 of *Dictyostelium discoideum* is lipid-linked and its fate altered in the presence of tunicamycin

Betsy L. Barent, Catherine P. Chia, School of Biological Sciences, University of Nebraska

We are studying the structure and role in phagocytosis of gp130, a glycoprotein located on the extracellular surface of *D. discoideum* amoebae. Predictions from the protein sequence of gp130 deduced from the cDNA indicate its attachment to the membrane via a glycosylphosphatidylinositol (GPI)-anchor. This was confirmed when radiolabeled palmitic acid was incorporated into gp130 of axenically grown cells, and when nitrous acid deamination released gp130 from purified membranes. The GPI-anchor of gp130 resisted cleavage by bacterial and mammalian phosphoinositol-specific phospholipases. However, in the presence of protease inhibitors, we detected *in vitro* a time-dependent cleavage of gp130 that was effectively inhibited by ZnCl₂. This activity and its inhibition were analogous to the release documented for two known GPI-linked proteins, gp80 of *D. discoideum* and VSG of *Trypanosoma brucei*. Cells treated with tunicamycin (0.5 µg/ml; TM), an inhibitor of N-linked glycosylation, produced an immunoreactive 85 kDa version of gp130 that was apparently unanchored and soluble rather than membrane-associated. The 85 kDa species did not bind Concanavalin A, and thus presumed to be devoid of (N-linked) carbohydrate. Also detected in TM-treated cells but absent in untreated, control cells were small quantities of an 81 kDa species that was either a second unglycosylated form of gp130 or a proteolytic product. TM-treated cells had two-fold less mature gp130 than control cells, as judged by densitometric analyses of immunoblots of cell lysates and sucrose gradient-purified plasma membranes. The totaled signals from the gp130-immunoreactive species of TM-treated cells were comparable to the gp130 signal of control cells. We conclude that the TM-inhibited glycosylation interfered with the attachment of the GPI-anchor to gp130. Consequently, a soluble and unglycosylated form of gp130 accumulated, presumably in the lumen of the ER/Golgi network.