

1982

A Solution To A Problem Of J. R. Ringrose

David R. Larson
University of Nebraska-Lincoln

Follow this and additional works at: <http://digitalcommons.unl.edu/mathfacpub>

Larson, David R., "A Solution To A Problem Of J. R. Ringrose" (1982). *Faculty Publications, Department of Mathematics*. 96.
<http://digitalcommons.unl.edu/mathfacpub/96>

This Article is brought to you for free and open access by the Mathematics, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Mathematics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

A SOLUTION TO A PROBLEM OF J. R. RINGROSE¹

BY DAVID R. LARSON

We announce a solution to a multiplicity problem for nests posed by J. R. Ringrose approximately twenty years ago. This also answers a question posed by R. V. Kadison and I. M. Singer, and independently by I. Gohberg and M. Krein concerning the invariant subspace lattice of a compact operator. The key to the proof is a result concerning compact perturbations of nest algebras which was recently obtained by Niels Andersen in his doctoral dissertation. The complete proof of the general result as well as of a number of related results will appear elsewhere. A proof for the special case which answers Ringrose's original question is included herein.

Let H be infinite dimensional separable Hilbert space. A *nest* N is a family of closed subspaces of H linearly ordered by inclusion. N is *complete* if it contains $\{0\}$ and H and contains the intersection and the join (closed linear span) of each subfamily. The corresponding *nest algebra* $\text{alg } N$ is the algebra of all operators in $L(H)$ which leave every member of N invariant. The *core* C_N is the von Neumann algebra generated by the projections on the members of N , and the *diagonal* \mathcal{D}_N is the von Neumann algebra $(\text{alg } N) \cap (\text{alg } N)^*$. N is *continuous* if no member of N has an immediate predecessor or immediate successor. Equivalently, N is continuous if the core C_N is a nonatomic von Neumann algebra. N has *multiplicity one* (is multiplicity free) if \mathcal{D}_N is abelian, or equivalently, if C_N is a m.a.s.a.

J. R. Ringrose posed the following question: Let N be a multiplicity free nest and $T: H \rightarrow H$ a bounded invertible operator. Is the image nest $TN = \{TN: N \in N\}$ necessarily multiplicity free? Note that $T(\text{alg } N)T^{-1} = \text{alg}(TN)$, so it is natural to say that TN is the similarity transform of N . Is multiplicity preserved under similarity? We show that the answer is no. It should be noted that a negative answer was conjectured in recent years by several mathematicians including J. Ringrose and W. Arveson.

The following key result is due to N. Andersen [1]. Let LC denote the compact operators in $L(H)$.

Received by the editors December 15, 1981.

1980 *Mathematics Subject Classification*. Primary 46L15, 47A15; Secondary 47B15.

¹ Supported in part by a grant from the NSF.

THEOREM (ANDERSEN). *If H is separable, and if N, M are arbitrary continuous nests in H , then there exists a unitary operator U such that $\text{alg } N + LC = U(\text{alg } M + LC)U^* = \text{alg}(UM) + LC$.*

Next, we answer Ringrose’s question.

THEOREM 1. *Let N be a continuous nest of multiplicity 1. Then there exists a positive invertible operator $T \in L(H)$ such that $TN = \{TN : N \in \mathcal{N}\}$ fails to have multiplicity 1.*

PROOF. By Andersen’s theorem there exists a continuous nest M not of multiplicity 1 such that $\text{alg } M + LC = \text{alg } N + LC$. Since for any algebra A we have $A + LC/LC \approx A/A \cap LC$, the algebras $\text{alg } M/\text{alg } M \cap LC$ and $\text{alg } N/\text{alg } N \cap LC$ are algebraically isomorphic. The diagonal $\mathcal{D}_M = \text{alg } M \cap (\text{alg } M)^*$ is a non-abelian von Neumann algebra so contains a nonzero partial isometry v with orthogonal initial and final spaces. Let $\tilde{S} = v + v^* - vv^* - v^*v + I$ and $\tilde{P} = vv^*$. Then $\tilde{S}^2 = I$ and $\tilde{P}\tilde{S}\tilde{P} = 0$. Since M is continuous \mathcal{D}_M contains no compacts, so \tilde{P} has infinite rank. Thus via the algebraic isomorphism between quotients it follows that $\text{alg } N/\text{alg } N \cap LC$ contains elements \hat{P}, \hat{S} with $\hat{P}^2 = \hat{P} \neq 0, \hat{S}^2 = I, \hat{P}\hat{S}\hat{P} = 0$.

Let A and B be elements of $\text{alg } N$ whose images in the quotient are \hat{P} and \hat{S} respectively. Then $A^2 - A, B^2 - I$ and ABA are contained in $\text{alg } N \cap LC$, and this is contained in the Jacobson radical \mathcal{R}_N of $\text{alg } N$ since N is continuous. So B is invertible in $\text{alg } N$. Also, a well-known result [cf. 9, Theorem 2.3.9] states that an element of a Banach algebra which is idempotent modulo the radical is equal modulo the radical to an idempotent. So there exists an idempotent $P \in \text{alg } N$ with $A - P \in \mathcal{R}_N$. We have $P \neq 0$ since otherwise A would be in \mathcal{R}_N and hence \hat{P} above would be a quasinilpotent idempotent, hence 0.

We have $PBP \in \mathcal{R}_N$. Set $B_1 = B - PBP$. Then B_1 is also invertible in $\text{alg } N$, and $PB_1P = 0$. Now set

$$S = B_1P + PB_1^{-1}(I - P) - B_1PB_1^{-1}(I - P) + I - P.$$

We have $PSP = 0$, and it can be verified that $S^2 = I$. (Let α denote the sum of the first two terms, β the sum of the remaining terms, and compute $\beta^2 = \beta, \alpha\beta = \beta\alpha = 0, \alpha^2 = I - \beta$.)

Let $R = I - 2P$. Then $R^2 = I, S^2 = I, RS \neq SR$. Let G be the group generated by R, S . We have $SRS = I - 2SPS$, and $PSP = 0$, so $PSRS = P = SRSP$. Hence R commutes with SRS since P does. It easily follows that

$$G = \{I, S, R, RS, SR, SRS, RSR, SRSR\}.$$

So G is a finite noncommutative group contained in $\text{alg } N$. Set $T = (\sum_{g \in G} g^*g)^{1/2}$. Then $TGT^{-1} = \{TgT^{-1} : g \in G\}$ is a noncommutative group of

unitaries contained in the diagonal of $\text{alg}(TN)$, and thus TN fails to have multiplicity 1. \square

Theorem 1 serves to answer an open question concerning invariant subspace lattices of compact operators due to Kadison and Singer [5] and to Gohberg and Krein [4]. An operator is said to be *hyperintransitive* if its lattice of invariant subspaces contains a multiplicity one nest.

COROLLARY 2. *There exists a nonhyperintransitive compact operator.*

PROOF. Let V be the Volterra operator. Then $\text{Lat } V$ is a continuous multiplicity one nest. Let $N = \text{Lat } V$ and let T be an invertible operator such that TN does not have multiplicity one. Since $\text{Lat}(TVT^{-1}) = TN$ and since TN is a maximal nest the similarity TVT^{-1} is not hyperintransitive. \square

REMARK. It was known for a number of years that a negative resolution to the Ringrose problem would yield Corollary 2. I believe that this connection was first observed by J. Erdos, and it was first shown to me by W. Arveson.

We strengthen Theorem 1 as follows.

THEOREM 3. *Let N be a continuous nest of multiplicity one. Then given $\epsilon > 0$ there exists a positive invertible operator $T \in L(H)$ with $T - I$ compact and $\|T - I\| < \epsilon$ such that $TN = \{TN : N \in N\}$ fails to have multiplicity one.*

A nest has *purely atomic core* if its core is generated by its minimal projections. The following shows that similarity transforms can fail to act “absolutely continuously” on nests.

THEOREM 4. *If N is a complete uncountable nest with purely atomic core there exists a positive invertible operator T such that TN does not have purely atomic core.*

A nest N is said to have the *factorization property* if every invertible positive operator T factors $T = A^*A$ for $A \in (\text{alg } N) \cap (\text{alg } N)^{-1}$. Arveson [2] proved that nests of the “simplest type” have the factorization property. We generalize this to countable complete nests, and then show that these are the only ones with this property.

THEOREM 5. *A complete nest has the factorization property if and only if it is countable.*

In contrast, if we drop the requirement that A^{-1} also be in $\text{alg } N$ we obtain.

THEOREM 6. *Let N be an arbitrary nest. Then every invertible positive operator T factors $T = A^*A$ for $A \in \text{alg } N$, A invertible in $L(H)$.*

The following answers a question of J. Erdos [3].

THEOREM 7. *Let \mathcal{N} be a continuous nest. Then the commutator ideal of $\text{alg } \mathcal{N}$ is not proper.*

REFERENCES

1. N. Andersen, *Compact perturbations of reflexive algebras*, J. Funct. Anal. **38** (1980), 366–400.
2. W. Arveson, *Interpolation problems in nest algebras*, J. Funct. Anal. **20** (1975), 208–233.
3. J. Erdos, *Non-selfadjoint operator algebras*, Proc. Roy. Irish Acad. Sect. A **81A** (1981), 127–145.
4. I. Gohberg and M. Krein, *Theory and application of Volterra operators in Hilbert space*, Transl. Math. Mono., vol. 24, Amer. Math. Soc., Providence, R.I., 1970.
5. R. Kadison and I. Singer, *Triangular operator algebras*, Amer. J. Math. **82** (1960), 227–259.
6. D. Larson, *On certain reflexive operator algebras*, Ph. D. thesis, Berkeley, 1976.
7. J. Ringrose, *On some algebras of operators*, Proc. London Math. Soc. **15** (1965), 61–83.
8. ———, *Algebraic isomorphisms between ordered bases*, Amer. J. Math. **83** (1961), 463–478.
9. C. Rickart, *General theory of Banach algebras*, Van Nostrand, New York, 1960.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF
NEBRASKA AT LINCOLN, LINCOLN, NEBRASKA 68588-0323