

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Harold W. Manter
Laboratory of Parasitology

Parasitology, Harold W. Manter Laboratory of

2005

Eimeria Species (Apicomplexa: Eimeriidae) from Arctic Ground Squirrels (*Spermophilus parryii*) and Red Squirrels (*Tamiasciurus hudsonicus*) in Alaska and in Siberia, Russia

Robert S. Seville

University of Wyoming, sseville@uwyo.edu

Clint E. Oliver

University of Wyoming

Andrew J. Lynch

University of New Mexico

Michelle C. Bryant

University of Wyoming

Donald W. Duszynski

University of New Mexico, eimeria@unm.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/parasitologyfacpubs>



Part of the [Parasitology Commons](#)

Seville, Robert S.; Oliver, Clint E.; Lynch, Andrew J.; Bryant, Michelle C.; and Duszynski, Donald W., "Eimeria Species (Apicomplexa: Eimeriidae) from Arctic Ground Squirrels (*Spermophilus parryii*) and Red Squirrels (*Tamiasciurus hudsonicus*) in Alaska and in Siberia, Russia" (2005). *Faculty Publications from the Harold W. Manter Laboratory of Parasitology*. 146.
<http://digitalcommons.unl.edu/parasitologyfacpubs/146>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Harold W. Manter Laboratory of Parasitology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

EIMERIA SPECIES (APICOMPLEXA: EIMERIIDAE) FROM ARCTIC GROUND SQUIRRELS (*SPERMOPHILUS PARRYII*) AND RED SQUIRRELS (*TAMIASCIURUS HUDSONICUS*) IN ALASKA AND IN SIBERIA, RUSSIA

Robert S. Seville, Clint E. Oliver, Andrew J. Lynch*, Michelle C. Bryant, and Donald W. Duszynski*

Department of Zoology and Physiology, University of Wyoming/Casper College Center, Casper, Wyoming 82601. e-mail: sseville@uwyo.edu

ABSTRACT: Fecal samples from arctic ground squirrels (*Spermophilus parryii*) collected in Alaska (n = 90) and Russia (n = 46) and from red squirrels (*Tamiasciurus hudsonicus*) in Alaska (n = 35) were examined for the presence of *Eimeria* spp. (Apicomplexa: Eimeriidae). Four species were recovered from arctic ground squirrels, including *Eimeria callospermophili* (prevalence = 18%), *Eimeria cynomysis* (23.5%), *Eimeria lateralis* (19%), and *Eimeria morainensis* (77%). A single species, *Eimeria tamiasciuri* (91%), was recovered from red squirrels. Eimerians recovered from arctic ground squirrels represent new host records, and the single species from red squirrels is a new geographic record. Alaskan arctic ground squirrel prevalence was higher for *E. callospermophili* (Alaska = 22% vs. Russia = 9%), *E. cynomysis* (34% vs. 2%), and *E. lateralis* (27% vs. 4%), but not *E. morainensis* (78% vs. 76%).

Arctic ground squirrels (*Spermophilus parryii* (Richardson, 1825)) occur in northwest Canada, Alaska, United States, and northeast Russia. They are sciurid rodents that include 5 genera and ~82 species adapted to a predominately fossorial–terrestrial existence (Wilson and Reeder, 1993). Wilber et al. (1998) reviewed all published literature on the coccidia known from marmotine rodents and recognized and provided descriptions for 26 valid *Eimeria* species.

Red squirrels (*Tamiasciurus hudsonicus* Erxleben, 1777) range from Alaska through the northeastern United States south to South Carolina. Some sciurids in the Tribe Tamiasciurini (3 genera and ~11 species) are adapted primarily to an arboreal existence (Wilson and Reeder, 1993). To date, only 2 species of coccidia—*Eimeria tamiasciuri* Levine, Ivens & Kruidenier, 1957, and *Eimeria toddi* Dorney, 1962—have been reported from red squirrels in the continental United States (Levine and Ivens, 1990).

In collaboration with the Beringian Coevolution project, which is systematically inventorying the mammals and associated parasites and pathogens of the Beringian region, we received feces from arctic ground squirrels collected from 2 unnamed locations and 5 national preserves in Alaska and 12 locations in northeastern Russia and red squirrels collected from 4 national preserves in Alaska. Here, we report results of our survey of eimerian species in arctic ground squirrels and red squirrels from these 2 regions and taxonomic details for the species recovered.

MATERIALS AND METHODS

Arctic ground squirrels were collected by shooting from 2 locations and 5 national preserves in Alaska (Bering Land Bridge National Preserve, Cape Krusenstern National Monument, Noatak National Preserve, Wrangle–St. Elias National Park [W-SENP], and Yukon-Charley Rivers National Preserve [Y-CRNP]), United States, and 12 locations in Magadanskaya oblast and Chukotka, Russia, during the summers of 2000, 2001, and 2002. Red squirrels were collected in Alaska from Kobuk Valley National Park, W-SENP, and Y-CRNP in 2001. Details regarding mammal collection are provided in Edingsaas et al. (2004). Fresh fecal samples were obtained from the colon of each host in the field and stored in vials containing 2% (w/v) aqueous potassium dichromate at ambient temperature to allow oocyst sporulation until ex-

amination at the University of Wyoming/Casper College Center and the University of New Mexico. Oocysts were isolated by flotation in saturated sucrose solution (specific gravity = 1.2) and identified with oil immersion lenses on a compound microscope with bright field and Nomarski differential interference contrast microscopy. Standardized abbreviations for oocyst and sporocyst characters, with a single exception (sporozoites = SZ), are those recommended by Wilber et al. (1998), including oocyst characters length (L) and width (W)—their ranges and ratio (L/W)—micropyle (M), residuum (OR), and polar granules (PG) and sporocyst characters length (L), width (W)—their ranges and ratio (L/W)—stieda body (SB), substieda body (SSB), parastieda body (PSB), residuum (SR), refractile bodies (RB), and nucleus (N). Measurements of standard morphologic parameters were made with an ocular micrometer and are reported as an average (μm) followed by the range in parentheses. Photomicrographs were produced, and photovouchers of sporulated oocysts were accessioned into the U.S. National Parasite Collection (USNPC), Beltsville, Maryland.

RESULTS

Fecal samples from 136 arctic ground squirrels and 35 red squirrels were examined, and 118 of the former and 32 of the latter were positive for the presence of *Eimeria* (87 and 91%, respectively). Four species were identified from arctic ground squirrels, including *Eimeria callospermophili* (prevalence = 18%), *Eimeria cynomysis* (23.5%), *Eimeria lateralis* (19%), and *Eimeria morainensis* (77%). Only a single species, *E. tamiasciuri* (91%), was identified from red squirrels. Table I presents summary data on prevalences for each species in each geographic region. Arctic ground squirrel species richness (number of eimerian species per animal/number hosts examined) was 1.4, with 67 animals (49.2%) having single-species infections and 35 (25.7%) with 2 species, 14 (10.3%) with 3 species, and 2 (1.5%) with 4 species infecting simultaneously.

DESCRIPTION

Eimeria callospermophili Henry, 1932

Oocyst shape spheroid to subspheroid; wall ~1–2 thick, with 2 layers: outer layer smooth and approximately two thirds of total thickness; L × W (n = 13) 23.5 × 20.9 (19–25.5 × 16.5–26); L/W 1.1 (1–1.2); M absent; OR present; OR characteristics compact, homogeneous mass, granular or smooth, 3–6 × 3–6; PG usually 1, but up to 4 present. Sporocysts (n = 10) lemon-shaped; L × W 10.5 × 8.2 (9–14 × 7–9); L/W 1.3 (1–1.6); SB present and nipplelike; SSB, PSB both absent; SR present as few dispersed granules.

Received 10 September 2004; revised 10 November 2004; accepted 6 December 2004.

* Department of Biology, The University of New Mexico, Albuquerque, New Mexico 87131.

TABLE I. Country of origin, locations, geographic coordinates, sample sizes, and eimerian species present (number of animals infected) for samples collected from arctic ground squirrels (*Spermophilus parryii*).

Country	Location	Geographic coordinates	n	Species present (no. infected)*			
Russia	1	64.25°N, 172.32°W	5		EICY (1)		EIMO (2)
	2	64.25°N, 172.45°W	5	EICA (1)			EIMO (4)
	3	64.32°N, 172.45°W	4			EILA (1)	EIMO (4)
	4	64.39°N, 172.32°W	9	EICA (3)		EILA (1)	EIMO (7)
	5	64.52°N, 172.40°W	3				EIMO (3)
	6	65.13°N, 172.20°W	9				EIMO (5)
	7	60.79°N, 151.73°E	1				EIMO (1)
	8	60.83°N, 151.70°E	1				EIMO (1)
	9	61.85°N, 147.64°E	2				EIMO (2)
	10	63.35°N, 158.58°E	1				
	11	64.57°N, 177.32°E	3				EIMO (3)
	12	64.81°N, 177.55°E	4				EIMO (3)
Total			46	EICA (4)	EICY (1)	EILA (2)	EIMO (35)
United States	1	64.72°N, 163.98°W	7	EICA (5)	EICY (4)	EILA (3)	EIMO (5)
	2	64.90°N, 165.11°W	1				
	Bering Land Bridge National Preserve	65.38°N, 163.23°W	9	EICA (2)	EICY (4)	EILA (1)	EIMO (9)
		65.85°N, 164.70°W	8	EICA (2)	EICY (5)	EILA (3)	EIMO (7)
		66.39°N, 164.49°W	12	EICA (1)	EICY (5)	EILA (5)	EIMO (11)
	Cape Krusenstern National Monument	67.52°N, 163.60°W	4	EICA (4)		EILA (2)	EIMO (4)
		67.52°N, 163.58°W	3	EICA (1)	EICY (2)	EILA (2)	EIMO (3)
	Noatak National Preserve	67.47°N, 162.22°W	5		EICY (3)		EIMO (3)
		68.34°N, 158.75°W	3		EICY (2)	EILA (2)	EIMO (2)
	Yukon-Charley Rivers National Preserve	65.05°N, 140.97°W	14	EICA (4)	EICY (1)		EIMO (11)
	Wrangle-St. Elias National Park	60.99°N, 142.03°W	6		EICY (2)	EILA (1)	EIMO (3)
		61.00°N, 142.75°W	4				EIMO (3)
		61.07°N, 143.90°W	2	EICA (1)		EILA (2)	EIMO (2)
		61.32°N, 144.24°W	5		EICY (2)	EILA (1)	EIMO (5)
		61.79°N, 141.21°W	1				EIMO (1)
	61.83°N, 141.83°W	4		EICY (1)	EILA (2)	EIMO (1)	
	62.03°N, 141.13°W	2					
Total			90	EICA (20)	EICY (31)	EILA (24)	EIMO (70)

* Abbreviation: EICA, *Eimeria callospermophili*; EICY, *Eimeria cynomys*; EILA, *Eimeria lateralis*; EIMO, *Eimeria morainensis*.

Taxonomic summary

Synonyms: Seven (see Wilber et al., 1998).

Type host: *Spermophilus lateralis* (Say, 1823), golden-mantled ground squirrel.

Other hosts: Twenty-five species in the genera *Cynomys* (3 species), *Marmota* (5), and *Spermophilus* (17) (see Wilber et al., 1998); *S. parryii* (this study).

Type locality: North America: Placer County, California, United States.

Geographic distribution: Asia: Kazakhstan, Buriatia. North America: Alberta, Canada; Mexico; Colorado, Idaho, Illinois, Iowa, Montana, New York, New Mexico, Pennsylvania, Texas, Utah, and Wyoming, United States. In 2000–2003, collected from *S. parryii* in Alaska and in Magadanskaya oblast and Chukotka, Russia (this study).

Prevalence: One of three (33%) in type host; 76 of 123 (63%) *Cynomys gunnisoni* from Utah (Thomas and Stanton, 1994); 43 of 61 (71%) *Cynomys leucurus* and 25 of 39 (64%) *Cynomys ludovicianus* from Wyoming (Seville, 1997); 2 of 2 (100%) *Spermophilus beecheyi* from California (Henry, 1932); 211 of 1,007 (21%) and 36 of 69 (52%) *Spermophilus elegans* from Wyoming (Shults et al., 1990); 36 of 100 (36%) *Spermophilus richardsonii* from Alberta, Canada (Seville and Stanton, 1993); 5–80% *Spermophilus townsendii* in Idaho (Wilber et al., 1994); 6 of 81 (7%) *Spermophilus undulatus* from Kazakhstan (Matschoulsky, 1949); 4 of 7 (57%) *Spermophilus variegatus* from Utah (Thomas and Stanton, 1994); and 20 of 90 (22%) and 4 of 46 (9%) *S.*

parryii from Alaska and Magadanskaya oblast and Chukotka, Russia, respectively (this study).

Sporulation: From 2.5 to 7 days (Crouch and Becker, 1931; Henry, 1932; Todd and Hammond, 1968a).

Prepatent period: Five to six days (Todd and Hammond, 1968a).

Patent period: Nine days (Todd and Hammond, 1968a).

Site of infection: Endogenous stages most numerous in the jejunum and upper ileum. The epithelium of the entire villus infected, but the endogenous stages were usually concentrated at the tips and along the distal one third. There was no observed pathogenicity (Todd and Hammond, 1968a).

Materials deposited: Photovouchers of sporulated oocysts from *S. parryii* deposited in USNPC 95305 (Russia) and 95306 (Alaska).

Remarks

The description of *E. callospermophili* from arctic ground squirrels from Alaska and Russia is similar to the composite description developed by reviewing all published descriptions of this species from all marmotine hosts by Wilber et al. (1998). In addition, the arctic ground squirrel specimens had 2-layered oocyst walls, with the outer layer two thirds of the total thickness; were slightly larger ($L \times W = 23.5 \times 20.9$ vs. 19.2×16.0); but had similar subspheroid sporocysts ($L \times W = 10.5 \times 8.2$ vs. 10.2×8.5), with SR consisting of few dispersed granules.

When oocysts from Alaska ($n = 8$) and Russia ($n = 5$) were com-

pared, few differences were noted. Oocyst L × W was similar (Alaska = 23.7 × 20.8 vs. Russia = 23.2 × 21.0), but Russian sporocysts were larger (L × W = 11.3 × 9.0 vs. 10.1 × 7.8).

Eimeria cynomysis Andrews, 1928

Oocyst shape ovoid; wall ~ 1.5 thick, with 2 layers: outer layer yellow/brown, rough; L × W (n = 28) 35.9 × 28.1 (34–39.5 × 25.6–32.5); L/W 1.3 (1.4–1.2); M present, 5.7–10 wide; MC absent; OR absent; 0–1 PG present. Sporocysts (n = 18) ovoid; L × W 18.2 × 10.1 (14–21 × 8–12); L/W 1.9 (1.6–2.1); SB present; SSB and PSB both absent; SR present as small granular mass or occasionally as large, membrane-bound granular mass located centrally between the sporozoites or occasionally below the SB.

Taxonomic summary

Synonyms: Two (see Wilber et al., 1998).

Type host: *Cynomys ludovicianus* (Ord, 1815), the black-tailed prairie dog.

Other hosts: Thirteen in *Cynomys* (2) and *Spermophilus* (11) (see Wilber et al., 1998); *S. parryii* (this study).

Type locality: North America: Ohio, United States (animals from a supply house, origin unknown, but analysis done in Ohio).

Geographic distribution: Europe: Czech Republic. North America: Alberta, Canada; California, Colorado, Idaho, Iowa, Ohio, Utah, and Wyoming, United States. In 2000–2003, collected from *S. parryii* in Alaska and in Magadanskaya oblast and Chukotka, Russia (this study).

Prevalence: Two of two (100%) in type host; 12 of 123 (10%) *C. gunnisoni* from Utah (Thomas and Stanton, 1994); 3 of 18 (17%); Shults et al., 1990) and 1 of 61 (2%); Seville, 1997) *C. leucurus* from Wyoming; 9 of 26 (35%) *Spermophilus columbianus* from Alberta, Canada (Hilton and Mahrt, 1971); 121 of 1,007 (12%); Shults et al., 1990) and 11% (number infected and sample size not reported; Seville et al., 1996) *S. elegans* from Wyoming; 2 of 8 (25%) *Spermophilus franklinii* from Alberta, Canada (Hilton and Mahrt, 1971); 2 of 3 (66%) *S. lateralis* from California (Henry, 1932); 6 of 100 (6%); Seville and Stanton, 1993) and 13 of 121 (11%); Hilton and Mahrt, 1971) *S. richardsonii* from Alberta, Canada; 35 of 788 (4%) *S. townsendii* from Idaho (Wilber et al., 1994); 6 of 18 (19%) from Iowa (Kietzmann and Kietzmann, 1987) and 2 of 56 (4%) *Spermophilus tridecemlineatus* from Wyoming (Seville et al., 1992); 2 of 7 (29%) *S. variegatus* from Utah (Thomas and Stanton, 1994); and 31 of 90 (34%) from Alaska and 1 of 46 (2%) *S. parryii* from Magadanskaya oblast and Chukotka, Russia (this study).

Sporulation: Eight to eleven days (Henry, 1932; Hall and Knippling, 1935; Todd et al., 1968).

Prepatent period: Ten to eleven days (Todd et al., 1968).

Patent period: Five to twenty-one days (Todd et al., 1968).

Site of infection: Todd et al. (1968) observed endogenous stages in the tips of the villi in the ileum and jejunum, but not in the duodenum. Little cell damage was observed, but infections did cause pathogenicity. Oocysts used for description recovered from feces.

Materials deposited: Phototype of sporulated oocyst in USNPC 87250. Photovouchers of sporulated oocysts from *S. parryii* deposited in USNPC 95307 (Alaska) and 95308 (Russia).

Remarks

The description of oocysts of *E. cynomysis* from arctic ground squirrels is consistent with the composite description developed by Wilber et al. (1998). Oocysts from arctic ground squirrels were longer and narrower (L × W = 35.9 × 28.1 vs. 35.4 × 30.0), making them more ellipsoidal (L/W 1.3 vs. 1.2); the micropyle width was more variable (6–10 vs. 5–6); the sporocysts were larger (L × W = 18.2 × 10.1 vs. no average reported, range 14–20 × 8–11.5 vs. 13–17 × 8–12); and the SR was either a small or large membrane-bound granular mass rather than coarsely granular in large amounts.

When oocysts from Alaska (n = 8) and Russia (n = 20) were compared, only minor differences were observed. Oocyst L × W was similar (Alaska = 36.4 × 28.0 vs. Russia = 35.7 × 28.2); Alaska sporocysts were smaller (L × W = 17.3 × 9.3 vs. 18.9 × 10.6); and oocysts from Alaska had a wider micropyle than those from Russia (MW = 7.2 vs. 8.8).

Eimeria lateralis Levine, Ivens, and Kruidenier, 1957

Oocyst shape ellipsoid to ovoid; wall ~ 2 thick, with 2 layers: outer layer rough, yellow/brown, and approximately two thirds of total thickness; L × W (n = 42) 31.7 × 24.7 (27–37.5 × 20–28); L/W 1.3 (1.1–1.5); M absent; OR absent; PG 0–4 present. Sporocysts (n = 43) ovoidal, L × W 15.8 × 10.0 (12–18 × 9–11); L/W 1.6 (1.4–1.8); SB present, nipplelike; SSB and PSB both absent; SR present as numerous, large granules dispersed throughout sporocyst or along edge of sporocyst.

Taxonomic summary

Synonyms: Five (see Wilber et al., 1998).

Type host: *Spermophilus lateralis*.

Other hosts: Eighteen in the genera *Cynomys* (3), *Marmota* (3), *Spermophilus* (10), and *Tamias* (2) (see Wilber et al., 1998); *S. parryii* (this study).

Type locality: North America: Point Imperial, Arizona, on the North Rim of the Grand Canyon, Grand Canyon National Park, United States.

Geographic distribution: Asia: Kazakhstan. North America: Alberta, Canada; Baja California, Mexico; Arizona, California, Colorado, Idaho, Illinois, Massachusetts, Montana, New York, Pennsylvania, Utah, Wisconsin, and Wyoming, United States. In 2000–2003, collected from *S. parryii* in Alaska and in Magadanskaya oblast and Chukotka, Russia (this study).

Prevalence: One of one (100%) in type host; 4 of 123 (3%) *C. gunnisoni* from Utah (Thomas and Stanton, 1994); 4 of 17 (24%) *C. leucurus* from Wyoming (Seville and Williams, 1989); 4 of 86 (5%) *C. ludovicianus* from Colorado (Vetterling, 1964); 4 of 25 (16%) *Marmota monax* (Dorney, 1965) from Pennsylvania; 21 of 38 (55%) *M. monax* from Illinois (McQuiston and Wright, 1984); 39 of 232 (17%) *M. monax* from New York (Fleming et al., 1979); 2 of 36 (6%) *Spermophilus beldingi* from California (Veluvolu and Levine, 1984); 2 of 36 *S. columbianus* from Alberta, Canada (Hilton and Mahrt, 1971); 168 of 1,007 (17%) *S. elegans* from Wyoming (Shults et al., 1990); 3 of 69 (4%) *S. elegans* from Wyoming (Shults et al., 1990); 18 of 121 (15%) *S. richardsonii* from Alberta, Canada (Hilton and Mahrt, 1971); 6 of 100 (6%) *S. richardsonii* from Alberta, Canada (Seville and Stanton, 1993); 1–5% *S. townsendii* from Idaho (Wilber et al., 1994); 9 of 56 (16%) *S. tridecemlineatus* from Wyoming (Seville et al., 1992); 3 of 96 (3%) *Tamias obscurus* from Baja California, Mexico (Hill and Duszynski, 1986); 12 of 84 (14%) *Tamias striatus* from Wisconsin (Dorney, 1962); 6 of 75 (8%) *T. striatus* from Massachusetts (Duncan, 1968); 21 of 189 (11%) *T. striatus* from Wisconsin (Dorney 1963, 1966); 3 of 41 (9%) *T. striatus* from Pennsylvania (Seville and Patrick, 2001); and 24 of 90 (27%) from Alaska and 2 of 46 (4%) *S. parryii* from Magadanskaya oblast and Chukotka, Russia (this study).

Sporulation: Two to four weeks at room temperature (Dorney, 1962); 50% of oocysts sporulated after 49 days in the laboratory (Dorney, 1965); 7–10 days in the laboratory (Vetterling, 1964); 10–14 days at room temperature (Duncan, 1968).

Prepatent period: Five days in *Spermophilus armatus* (Todd and Hammond, 1968b).

Patent period: Six and a half days in *S. armatus* (Todd and Hammond, 1968b).

Site of infection: Dorney (1962) observed endogenous stages primarily in the middle of the small intestine of *M. monax* (Dorney, 1962). Todd and Hammond (1968b) observed endogenous stages in the epithelial cells of the jejunum and ileum of experimentally infected *S. armatus*.

Materials deposited: Phototype of sporulated oocyst in USNPC 87251. Symbiotype in University of Illinois Natural History Museum 10300. Photovouchers of sporulated oocysts from *S. parryii* deposited in USNPC 95309 (Alaska) and 95310 (Russia).

Remarks

The description of *E. lateralis* from arctic ground squirrels is similar to the composite description by Wilber et al. (1998). Overall, the oocysts were smaller (L × W = 31.7 × 24.7 vs. 35.1 × 26.8) and more ellipsoid (L/W = 1.6 vs. 1.3). The sporocysts were similar in size (L × W = 16 × 10) and had a prominent SB, and the SR was present as numerous large, dispersed granules.

When oocysts from Alaska (n = 4) and Russia (n = 39) were com-

pared, oocysts were similar in all respects, including $L \times W$ (Alaska = 32.6×24.8 vs. Russia = 31.7×24.7); L/W (1.3 vs. 1.2), and sporocyst size ($L \times W = 15.5 \times 10.1$ vs. 15.8×10.0).

Eimeria morainensis Torbett, Marquardt, and Carey, 1982

Oocyst shape subspheroid; wall 1–1.5, with 2 layers of equal thickness; $L \times W$ ($n = 47$) 23.3×19.5 (19.5 – 28×14 – 22.5); L/W 1.2 (1–1.4); M absent; OR absent; PB 0–3 present, often bilobed. Sporocysts ellipsoidal; $L \times W$ ($n = 37$) 11.7×7.4 (7.9 – 14.0×5.5 – 9.0); L/W 1.6 (1.3–2); SB present and usually prominent, dark and buttonlike; SR present as compact, occasionally membrane-bound granular mass, dispersed granules, or both.

Taxonomic summary

Type host: *Spermophilus lateralis*.

Other hosts: Nine in the genera *Cynomys* (3), *Marmota* (1), *Spermophilus* (5) (see Wilber et al., 1998); *S. parryii* (this study).

Type locality: North America: Moraine Park in Rocky Mountain National Park, Colorado (T5N, R73W, S29), United States.

Geographic distribution: North America: Alberta, Canada; Colorado, Idaho, Utah, and Wyoming, United States. In 2000–2003, collected from *S. parryii* in Alaska and in Magadanskaya oblast and Chukotka, Russia (this study).

Prevalence: One of thirty-five (3%) in type host; 4 of 18 (22%) *C. leucurus* from Wyoming (Shults et al., 1990); 435 of 1,007 (43%) *S. elegans* from Wyoming (Shults et al., 1990); 36 of 69 (52%) *S. elegans* from Colorado (Torbett et al., 1982); 5–18% *S. townsendii* from Idaho (Wilber et al., 1994); 6 of 7 (86%) *S. variegatus* from Utah (Thomas and Stanton, 1994); and 70 of 90 (78%) from Alaska and 35 of 46 (76%) *S. parryii* from Magadanskaya oblast and Chukotka, Russia (this study).

Sporulation: Six to seven days (Torbett et al., 1982).

Prepatent period: Eight to nine days (Torbett et al., 1982).

Patent period: Nine days (Torbett et al., 1982).

Site of infection: Unknown. Oocysts recovered from feces.

Materials deposited: Phototype of sporulated oocyst in USNPC 82933. Photovouchers of sporulated oocysts from *S. parryii* deposited in USNPC 95311–95313 (Russia).

Remarks

The description of *E. morainensis* from arctic ground squirrels is similar to the composite description by Wilber et al. (1998). Oocysts were larger ($L \times W = 23.3 \times 19.5$ vs. 20.3×19.8), had shorter but wider sporocysts ($L \times W = 11.7 \times 7.4$ vs. 12.1×6.9), and possessed a buttonlike SB and a membrane-bound granular SR, or a row of globules along side the sporocyst wall, or both.

When oocysts from Alaska ($n = 30$) and Russia ($n = 17$) were compared, oocysts were similar in all respects. Russian oocysts were larger ($L \times W = 23.4 \times 20.0$ vs. 23.2×19.2), but sporocyst size was similar ($L \times W = 11.9 \times 7.4$ vs. 11.8×7.4).

Eimeria tamiasciuri Levine, Ivens, and Kruidenier, 1957

Oocyst shape elongate ellipsoid; $L \times W$ 29×16 ; M absent; OR absent; PB present. Sporocysts ellipsoidal; $L \times W$ 13×7 ; SB cone-shaped.

Taxonomic summary

Type host: *Microtus montanus* (Peale, 1848), the montane vole.

Other hosts: *Sciurus aberti* Woodhouse, 1853; *Sciurus griseus* Ord, 1818; *Tamias dorsalis* Baird, 1855; *Tamias merriami* J.A. Allen, 1889; *T. obscurus* J.A. Allen, 1889; *Tamiasciurus mearnsi* (Townsend, 1897); *T. hudsonicus*.

Type locality: North America: north rim of the Grand Canyon, near the entrance to Grand Canyon National Park, Arizona, United States.

Geographic distribution: North America: Baja California, Mexico; California, New Mexico, and Wisconsin, United States. In 2000–2003, collected from *T. hudsonicus* in Alaska (this study).

Prevalence: One of five (20%) in type host; 3 of 9 (33%) *S. aberti* from Arizona (Hill and Duszynski, 1986); 4 of 11 (36%) *S. aberti* from New Mexico (Patrick and Wilson, 1995); 1 of 1 (100%) *S. griseus* from

California (Hill and Duszynski, 1986); 1 of 12 (8%) *T. dorsalis* from Arizona (Hill and Duszynski, 1986); 1 of 50 (2%) *T. merriami* from California (Hill and Duszynski, 1986); 1 of 96 (1%) *T. obscurus* from Baja California Norte, Mexico (Hill and Duszynski, 1986); 3 of 5 (60%) *T. mearnsi* from Baja California Norte, Mexico (Hill and Duszynski, 1986); 2 of 2 (100%) *T. hudsonicus* from New Hampshire (Bullock, 1959); 49 of 50 (98%) *T. hudsonicus* from Wisconsin (Dorney, 1962); 149 of 152 (98%) *T. hudsonicus* from Wisconsin (Dorney, 1966); 12 of 13 (92%) *T. hudsonicus* from Ontario, Canada (Soon and Dorney, 1969); 166 of 171 (97%) *T. hudsonicus* from Alberta, Canada (Mahrt and Chai, 1970); 88 of 89 (99%) *T. hudsonicus* from Alberta (Mahrt and Chai, 1972); 3 of 12 (25%) *T. hudsonicus* from New Mexico (Patrick and Wilson, 1995); 1 of 1 (100%) *T. h. mogollonensis* from Arizona (Hill and Duszynski, 1986); and 32 of 35 (91%) *T. hudsonicus* from Alaska (this study: 28/31 [90%] Yukon Charley; 2/2 [100%] Wrangle St. Elias; 2/2 [100%] Kobuk).

Sporulation: Three days (Soon and Dorney, 1969).

Prepatent period: Unknown.

Patent period: Unknown.

Site of infection: Unknown. Oocysts recovered from feces.

Materials deposited: Phototype of sporulated oocyst in USNPC 87256. No photovoucher from this host produced.

Remarks

The description of *E. tamiasciuri* from red squirrels from Alaska is similar to the original description by Levine et al. (1957) from the same host species in Arizona. Oocysts and sporocysts were smaller (oocyst $L \times W = 29 \times 16$ vs. 33×19 ; sporocyst $L \times W = 13 \times 7$ vs. 16×8), but there were no differences in overall morphology, including the elongate ellipsoid oocyst shape and cone-shaped SB. The recovery of *E. tamiasciuri* from *T. hudsonicus* establishes a new geographic record for the parasite.

DISCUSSION

Previously, Wilber et al. (1998) suggested that because members of the rodent tribe Marmotini arose fairly recently and are likely to harbor closely related parasites and that few cross-transmission or molecular studies have been conducted, the morphologic species concept was the only practical approach to naming and identifying *Eimeria* spp. in this rodent group. Comparison of the morphologies of the different species we observed in samples from arctic ground squirrels from both continents and with species reported from other marmotine hosts from other regions strongly supports conspecificity both across the different marmotine host species and between the eimerians recovered from arctic ground squirrels in Russia and in Alaska. However, because Alaska marmotine populations and their parasites have likely been isolated from Russia since the last Beringian land bridge, which was broken approximately 15,500 yr ago (Pielou, 1991), we suggest that molecular and cross-transmission studies be conducted to verify conspecificity.

Prevalences for all species in arctic ground squirrels except *E. morainensis* were significantly higher for the Alaskan than the Russian populations (Table I). Although we cannot offer a definitive explanation, one plausible reason could be that Alaskan samples were collected from National Parklands, so estimated prevalences represent undisturbed conditions, whereas Russian populations might have been subject to human disturbance via habitat disruption or squirrel population control, although we are unable to confirm this possibility.

There have been few surveys of arctic sciurids for coccidia, and this is the first to examine a large number of samples from a single host species. Arctic marmotine hosts not adequately sampled include *Marmota broweri*, *Marmota caligata*, *Marmota camtschatica*, and *Tamias sibiricus*. Being the first survey

of arctic ground squirrels and red squirrels in Beringia for coccidia, this report constitutes new host and geographic ranges for *E. callospermophili*, *E. cynomysis*, *E. lateralis*, and *E. morainensis* and a new geographic record for *E. tamiasciuri* in red squirrels. *Eimeria cynomysis* and *E. morainensis* are now documented to occur from the central United States to arctic Alaska and eastern Russia, *E. callospermophili* and *E. lateralis* from Mexico to arctic Alaska and eastern Russia, and *E. tamiasciuri* from Alaska to Mexico and the eastern United States.

ACKNOWLEDGMENTS

Fecal samples from arctic ground squirrels and red squirrels were collected as part of the Beringian Coevolution Project (BCP) funded by the NSF (DBI-0196095 to J. A. Cook). We thank all of the biologists of the various BCP field teams for their diligent efforts in providing specimens from remote field camps of Beringia. Angela Hick processed some of the samples. Additional support was provided by the University of Wyoming/Casper College Center, University of Wyoming Department of Zoology and Physiology, and the UW-NIH BRIN (Biomedical Research Infrastructure Network) and UW-NSF EPSCoR programs.

LITERATURE CITED

- BULLOCK, W. L. 1959. The occurrence of *Eimeria tamiasciuri* in New Hampshire. *Journal of Parasitology* **45**: 39–40.
- CROUCH, H. B., AND E. R. BECKER. 1931. Three species of coccidia from the woodchuck, *Marmota monax*. *Iowa State College Journal of Science* **2**: 127–131.
- DORNEY, R. S. 1962. A survey of the coccidia of some Wisconsin Sciuridae with descriptions of three new species. *Journal of Protozoology* **9**: 258–261.
- . 1963. Coccidiosis—Incidence, epizootiology, in two Wisconsin Sciuridae. North American Wildlife Natural Resources Conference **28**: 207–215.
- . 1965. *Eimeria tuscarorensis* n. sp. (Protozoa: Eimeriidae) and redescription of other coccidia of the woodchuck, *Marmota monax*. *Journal of Protozoology* **12**: 423–426.
- . 1966. Quantitative data on four species of *Eimeria* in eastern chipmunks and red squirrels. *Journal of Protozoology* **13**: 549–550.
- DUNCAN, S. 1968. *Eimeria ovata* n. sp. and other coccidia of the eastern chipmunk *Tamias striatus* in Massachusetts. *Journal of Protozoology* **15**: 319–320.
- EDDINGSAAS, A. A., B. K. JACOBSEN, E. P. LESSA, AND J. P. COOK. 2004. Evolutionary history of the arctic ground squirrel (*Spermophilus parryi*) in nearctic Beringia. *Journal of Mammalogy* **85**: 601–610.
- FLEMING, W. J., J. R. GEORGI, AND J. W. CASLICK. 1979. Parasites of the woodchuck (*Marmota monax*) in central New York State. *Proceedings of the Helminthological Society of Washington* **46**: 115–127.
- HALL, P. R., AND E. F. KNIPLING. 1935. *Eimeria franklinii* and *Eimeria eubeckeri*, two new species of coccidia from the Franklin ground squirrel, *Citellus franklinii* Sabine. *Journal of Parasitology* **21**: 128–129.
- HENRY, D. P. 1932. Observations on the coccidia of small mammals in California, with descriptions of seven new species. *University of California Publications in Zoology* **37**: 279–290.
- HILL, T. P., AND D. W. DUSZYNSKI. 1986. Coccidia (Apicomplexa: Eimeriidae) from sciurid rodents (*Eutamias*, *Sciurus*, *Tamiasciurus* spp.) from the Western U.S.A. and northern Mexico and descriptions of two new species. *Journal of Protozoology* **33**: 282–288.
- HILTON, D. F. J., AND J. L. MAHRT. 1971. *Eimeria spermophili* n. sp. and other *Eimeria* spp. (Sporozoa, Eimeriidae) from three species of Alberta *Spermophilus* (Rodentia, Sciuridae). *Canadian Journal of Zoology* **49**: 699–701.
- KIETZMANN, J. G. E., AND E. A. KIETZMANN. 1987. Parasites associated with thirteen-lined ground squirrels (*Spermophilus tridecemlineatus* Mitchell) from Ames, Iowa. *Proceedings of Iowa Academy of Sciences* **94**: 22–23.
- LEVINE, N. D., AND V. IVENS. 1990. *The coccidian parasites of rodents*. CRC Press, Boca Raton, Florida, 228 p.
- , ———, AND F. J. KRUIDENIER. 1957. New species of *Eimeria* from Arizona rodents. *Journal of Protozoology* **4**: 80–88.
- MAHRT, J. L., AND S. J. CHAL. 1970. Prevalence of coccidia in red squirrels (*Tamiasciurus hudsonicus*) in Alberta. *Canadian Journal of Zoology* **48**: 606.
- , AND ———. 1972. Parasites of red squirrels in Alberta, Canada. *Journal of Parasitology* **58**: 639–640.
- MATSCHOULSKY, S. N. 1949. About coccidia in rodents of southern areas of Buryat-Mongol, U.S.S.R. *Trudy Buryet-Mongol'skoi Zooveterinarnogo Instituta, Ulan-Ude* **5**: 40–56.
- MCQUISTON, T. E., AND J. M. WRIGHT. 1984. The prevalence and seasonal distribution of coccidial parasites of woodchucks (*Marmota marmota*). *Journal of Parasitology* **70**: 994–996.
- PATRICK, M. J., AND W. D. WILSON. 1995. Parasites of the Abert's squirrel (*Sciurus aberti*) and red squirrel (*Tamiasciurus hudsonicus*) of New Mexico. *Journal of Parasitology* **81**: 321–324.
- PIELOU, E. C. 1991. *After the Ice Age*. University of Chicago Press, Chicago, Illinois, 366 p.
- SEVILLE, R. S. 1997. *Eimeria* spp. (Apicomplexa: Eimeriidae) from black- and white-tailed prairie dogs (*Cynomys ludovicianus* and *Cynomys leucurus*) in central and southeast Wyoming. *Journal of Parasitology* **83**: 166–168.
- , AND M. J. PATRICK. 2001. *Eimeria* spp. (Apicomplexa: Eimeriidae) from the eastern chipmunk (*Tamias striatus*) in Pennsylvania with a description of one new species. *Journal of Parasitology* **87**: 165–168.
- , AND N. L. STANTON. 1993. Eimerian guilds (Apicomplexa: Eimeriidae) in Richardson's (*Spermophilus richardsonii*) and Wyoming (*Spermophilus elegans*) ground squirrels. *Journal of Parasitology* **79**: 973–975.
- , ———, AND K. GEROW. 1996. Stable parasite guilds: Coccidia in spermophiline rodents. *Oikos* **75**: 365–372.
- , D. M. THOMAS, R. PICKERING, AND N. L. STANTON. 1992. Species of *Eimeria* from the thirteen-lined ground squirrel, *Spermophilus tridecemlineatus* from Wyoming. *Great Basin Naturalist* **52**: 309–312.
- , AND E. S. WILLIAMS. 1989. Endoparasites of the white-tailed prairie dog *Cynomys leucurus* at Meeteetse, Park County, Wyoming USA. *Proceedings of the Helminthological Society of Washington* **56**: 204–206.
- SHULTS, L. M., R. S. SEVILLE, N. L. STANTON, AND G. MENKENS JR. 1990. *Eimeria* sp. (Apicomplexa: Eimeriidae) from Wyoming ground squirrels, *Spermophilus elegans*, and white-tailed prairie dogs, *Cynomys leucurus*, in Wyoming. *Great Basin Naturalist* **50**: 327–331.
- SOON, B. L., AND R. S. DORNEY. 1969. Occurrence of *Eimeria tamiasciuri* in Ontario red squirrels (*Tamiasciurus hudsonicus*). *Canadian Journal of Zoology* **47**: 731–732.
- THOMAS, D. M., AND N. L. STANTON. 1994. Eimerian species (Apicomplexa: Eimeriina) in Gunnison's prairie dogs (*Cynomys gunnisoni zuniensis*) and rock squirrels (*Spermophilus variegatus grammurus*) from southeastern Utah. *Journal of the Helminthological Society of Washington* **61**: 17–21.
- TODD, J. K. S., AND D. M. HAMMOND. 1968a. Life cycle and host specificity of *Eimeria callospermophili* Henry, 1932 from the Uinta ground squirrel *Spermophilus armatus*. *Journal of Protozoology* **15**: 1–8.
- , AND ———. 1968b. Life cycle and host specificity of *Eimeria larimerensis* Vetterling, 1964, from the Uinta ground squirrel *Spermophilus armatus*. *Journal of Protozoology* **15**: 268–275.
- , ———, AND L. C. ANDERSON. 1968. Observations on the life cycle of *Eimeria bilamellata* Henry, 1932 in the Uinta ground squirrel *Spermophilus armatus*. *Journal of Protozoology* **15**: 732–740.
- TORBETT, B. E., W. C. MARQUARDT, AND A. C. CAREY. 1982. A new species of *Eimeria* from the golden mantled ground squirrel, *Spermophilus lateralis*, in northern Colorado. *Journal of Protozoology* **29**: 157–159.
- VELUVOLU, P., AND N. D. LEVINE. 1984. *Eimeria beldingii* n. sp. and other coccidia (Apicomplexa) of the ground squirrel *Spermophilus beldingi*. *Journal of Protozoology* **31**: 357–358.
- VETTERLING, J. M. 1964. Coccidia (*Eimeria*) from the prairie dog, *Cy-*

- nomys ludovicianus ludovicianus*, in northern Colorado. *Journal of Protozoology* **11**: 89–91.
- WILBER, P. G., D. W. DUSZYNSKI, S. J. UPTON, R. S. SEVILLE, AND J. O. CORLISS. 1998. A revision of the taxonomy and nomenclature of the *Eimeria* spp. (Apicomplexa: Eimeriidae) from rodents in the Tribe Marmotini. *Systematic Parasitology* **39**: 113–135.
- , B. HANELT, B. VAN HORNE, AND D. W. DUSZYNSKI. 1994. Two new species and temporal changes in the prevalence of eimerians in a free-living population of Townsend's ground squirrels (*Spermophilus townsendii*) in Idaho. *Journal of Parasitology* **80**: 251–259.
- WILSON, D. E., AND D. M. REEDER. 1993. *Mammal species of the world*, 2nd ed. Smithsonian Institution Press, Washington, D.C., 1207 p.