

2007

# Genetic structure and molecular variability of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) collected in maize and cotton fields in Brazil

S. Martinelli

*Universidade de Sao Paulo*, samuelmartinelli@hotmail.com

P. L. Clark

*University of Nebraska-Lincoln*

M. I. Zucchi

*Instituto Agronômico de Campinas*

M. C. Silva-Filho

*ESALQ-USP*

John E. Foster

*University of Nebraska-Lincoln*, john.foster@unl.edu

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

Martinelli, S.; Clark, P. L.; Zucchi, M. I.; Silva-Filho, M. C.; Foster, John E.; and Omoto, C., "Genetic structure and molecular variability of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) collected in maize and cotton fields in Brazil" (2007). *Faculty Publications: Department of Entomology*. 164.

<http://digitalcommons.unl.edu/entomologyfacpub/164>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

S. Martinelli, P. L. Clark, M. I. Zucchi, M. C. Silva-Filho, John E. Foster, and C. Omoto

# Genetic structure and molecular variability of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) collected in maize and cotton fields in Brazil

S. Martinelli<sup>1\*</sup>, P.L. Clark<sup>2</sup>, M.I. Zucchi<sup>3</sup>, M.C. Silva-Filho<sup>4</sup>,  
J.E. Foster<sup>2</sup> and C. Omoto<sup>1</sup>

<sup>1</sup>Departamento de Entomologia, Fitopatologia e Zoologia Agrícola, Universidade de São Paulo/ESALQ, Av. Pádua Dias 11, 13418-900, Piracicaba, SP, Brazil: <sup>2</sup>Insect Genetics Laboratory, Department of Entomology, University of Nebraska, Lincoln, Nebraska, USA: <sup>3</sup>Instituto Agronômico de Campinas, Campinas, SP, Brazil: <sup>4</sup>Departamento de Genética, ESALQ-USP, Piracicaba, SP, Brazil

## Abstract

The purpose of this research was to evaluate the genetic similarity and structure of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith), populations associated with maize and cotton crops in Brazil using amplified fragment length polymorphisms. Mean genetic similarity among populations was 0.45. The unweighted pair group method with arithmetic mean analysis dendrograms did not separate populations of *S. frugiperda* into clusters related to the host plant in which the insects were collected. No genetic variation was observed among maize and cotton populations of *S. frugiperda*, suggesting that the same populations are injuring both crops in Brazil. This research validates the need for stewardship of crop-protection methods for managing *S. frugiperda* to reduce the incidence of pesticide resistance, due to the spatial and temporal overlapping of maize and cotton crops in some regions in Brazil.

**Keywords:** AFLP markers, fall armyworm, population structure, resistance management

## Introduction

The evolution of pesticide resistance has been identified worldwide as the most serious threat to the development of sustainable integrated pest management practices (Labbe *et al.*, 2005). The development of resistance is an evolutionary (Dobzhansky, 1951) and multi-disciplinary process, which is influenced by several interacting factors, such as the initial resistance allele frequency, inheritance of resistance, relative fitness of the various genotypes, management practices, among other factors (Georghiou & Taylor, 1977a,b).

The fall armyworm, *Spodoptera frugiperda* (J.E. Smith), is a migratory pest, which is endemic to the Western Hemisphere. The high adult dispersal capacity of this pest allows it to quickly spread along its range of host plants (Sparks, 1979; Johnson, 1988). *Spodoptera frugiperda* has a highly polyphagous feeding behaviour, which includes the consumption of different cultivated host plants, such as maize, cotton and rice (Yu *et al.*, 2003). Two morphologically indistinguishable strains of *S. frugiperda*, one associated with maize (maize host strain) and the other associated with rice (rice host strain), have been identified already in the USA (Pashley *et al.*, 1985; McMichael & Prowell, 1994; Levy *et al.*, 2002; Meagher *et al.*, 2004; Meagher & Nagoshi, 2004) and also in Brazil (Busato *et al.*, 2002, 2004).

\*Fax: +55 193433 0562

E-mail: samuelmartinelli@hotmail.com

Table 1. Identification of the populations of *Spodoptera frugiperda* obtained for AFLP analysis.

Population	Country	State or province	Host plant	Longitude	Latitude	Collection
1 Mex	Mexico	San Luis Potosi	Maize	100° 58' W	22° 09' N	08/10/2002
2 BR1	Brazil	Minas Gerais	Maize	48° 09' W	18° 56' S	11/13/2003
3 USA	United States of America	Florida	Maize	82° 33' W	29° 65' N	08/18/2004
4 ARG1	Argentina	Santa Fe	Maize	59° 40' W	29° 09' S	03/10/2004
5 ARG2	Argentina	Santiago del Estero	Maize	63° 53' W	27° 55' S	12/15/2004
6 ARG3	Argentina	Chaco	Maize	60° 27' W	26° 48' S	03/10/2004
7 ARG4	Argentina	Salta	Maize	64° 04' W	25° 18' S	03/20/2004
8 BR2	Brazil	São Paulo	Maize	50° 32' W	20° 16' S	03/08/2004
9 BR3	Brazil	Minas Gerais	Maize	46° 54' W	16° 21' S	03/17/2004
10 BR4	Brazil	Bahia	Maize	44° 59' W	12° 06' S	02/09/2004
11 BR5	Brazil	Goiás	Cotton	50° 55' W	17° 48' S	03/16/2004
12 BR6	Brazil	Mato Grosso	Cotton	54° 17' W	15° 33' S	03/04/2004
13 BR7	Brazil	Rio Grande do Sul	Maize	58° 48' W	29° 41' S	03/03/2004

*Spodoptera frugiperda* is the most destructive and economically important insect pest in maize fields in Brazil (Sena *et al.*, 2003). Insecticide resistant strains of *S. frugiperda* have already been identified by Pitre (1988), Yu (1991) and Diez-Rodriguez & Omoto (2001). In addition, *S. frugiperda* has been observed injuring cotton plants in the Midwest region of Brazil.

The variation of genetic structure of a pest population in space and time and gene flow among its sub-populations are greatly responsible for the rate of resistance evolution (Fuentes-Contreras *et al.*, 2004). The estimation of gene flow derived from the use of molecular markers can serve as an index of dispersal rates (Slatkin, 1985; Roderick, 1996). Molecular markers have been extensively used to evaluate the genetic similarity and to estimate the gene flow among insect populations (Figueroa *et al.*, 2002; Fuentes-Contreras *et al.*, 2004; Sosa-Gomez, 2004; Martinelli *et al.*, 2006). Amplified fragment length polymorphism (AFLP) markers have been successfully used to investigate the genetic similarity and population structure of different insect species (Meldenson & Shaw, 2002; Timm *et al.*, 2005; Zhang *et al.*, 2005), including *S. frugiperda* (McMichael & Prowell, 1994; Busato *et al.*, 2004).

Insect population genetic structure describes how populations are subdivided into finite breeding units (Roush & Daly, 1990) and can be correlated to local selection, patterns of mate selection and gene flow (McKenzie, 1996). Therefore, the comprehension of insect population genetic structure and intraspecific gene flow is essential for designing management practices to delay the evolution of resistance to any control method (Tabashnik, 1991; Caprio & Tabashnik, 1992).

The objective of this research was to estimate the genetic variability and structure of *S. frugiperda* populations associated with maize and cotton crops from different geographic regions in Brazil.

## Materials and methods

### *Insect material*

*Spodoptera frugiperda* adults and larvae were collected from maize and cotton fields at seven locations across Brazil. Additionally, four maize populations collected across Argentina, one population in Mexico and another one in the USA, were included in the analysis as outgroups (table 1).

Table 2. Sequences of specific adapters and preamplification primers.

Specification	Sequence
Adapter <i>Eco</i> R1	5'-CTCGTAGACTGCGTACC-3' 3'-CATCTGACGCATGGTTAA-5'
Preamplification <i>Eco</i> R1 primer	5'-GACTGCGTACCAATTC-3'
Adapter <i>Mse</i> 1	5'-GACGATGAGTCTCTGAG-3' 3'-TACTCAGGACTCAT-5'
Preamplification <i>Mse</i> 1 primer	5'-GATGAGTCTCTGAGTAA-3'

The collected larvae and adults were frozen and stored at  $-80^{\circ}\text{C}$ .

### *DNA extraction and quantification*

*Spodoptera frugiperda* genomic DNA was extracted using a CTAB method (Black & Duteau, 1997). Larvae were prepared for DNA extraction by removing the gut and the head. Larvae and adult thoraces were homogenized in 500  $\mu\text{l}$  extraction buffer (100 mM Tris-HCL pH 8.0, 1.4 M NaCl, 0.02 M EDTA, 2% CTAB, and 0.2% beta-mercaptoethanol). Proteinase K (concentration of 200  $\mu\text{g ml}^{-1}$  extraction buffer) was added to the homogenate for 2 h at  $65^{\circ}\text{C}$ . RNase A (500  $\mu\text{g ml}^{-1}$ ) was added to the homogenate and held for 3 h at  $37^{\circ}\text{C}$ . After RNA and protein were removed from each sample, the homogenate was centrifuged at 13,000 rpm for 5 min at room temperature. The supernatant was extracted with 500  $\mu\text{l}$  of chloroform:isoamyl alcohol (24:1) by centrifugation at 14,000 rpm for 20 min to separate the phases. The aqueous phase was transferred into an autoclaved 1.5 ml microcentrifuge tube and the chloroform:isoamyl step was repeated. DNA was precipitated by adding 400  $\mu\text{l}$  chilled ( $-20^{\circ}\text{C}$ ) isopropanol to the aqueous phase and incubated at  $4^{\circ}\text{C}$  for 8 h. After incubation, the precipitate was centrifuged at 13,000 rpm at  $4^{\circ}\text{C}$  for 30 min. The isopropanol was decanted off; the DNA pellet was rinsed with 500  $\mu\text{l}$  100% ETOH and centrifuged at 13,000 rpm at  $4^{\circ}\text{C}$  for 5 min and repeated a second time. The ETOH was decanted off and the pellet was air dried (24°) for 45 min. Fifty  $\mu\text{l}$  1 $\times$ TE buffer (10 mM Tris-HCL pH 8.0; 0.1 mM EDTA) was poured onto the DNA pellet and stored at  $4^{\circ}\text{C}$  for 8 h. After the DNA was re-suspended in 1 $\times$ TE buffer, each sample was

Table 3. Sequence of the primers used at the selective amplification step at the AFLP protocol.

Primer code	Sequence (5'-3')
EM01	GACTGCGTACCAATTC <u>ACT</u> × GATGAGTCCTGATAA <u>CAC</u>
EM02	GACTGCGTACCAATTC <u>ACT</u> × GATGAGTCCTGAGTAA <u>CAG</u>
EM03	GACTGCGTACCAATTC <u>ACT</u> × GATGAGTCCTGAGTAA <u>CTA</u>
EM04	GACTGCGTACCAATTC <u>ACT</u> × GATGAGTCCTGAGTAA <u>CTC</u>




Fig. 1. Mean genetic similarity pattern obtained for the thirteen *Spodoptera frugiperda* populations. UPGMA clustering criteria based on the Jaccard similarity index through AFLP analysis. Cophenetic correlation = 0.98.

quantified by running a 1% agarose gel with a  $\lambda$  DNA marker ( $22.2 \text{ ng } \mu\text{l}^{-1}$ ). TE buffer (1×) was added to DNA samples until they reached a concentration of  $22.2 \text{ ng } \mu\text{l}^{-1}$  concentration of genomic DNA. The agarose gels were visualized under UV light using Genomics Solutions software. The quantified DNA product was stored at  $-80^\circ\text{C}$  until used for AFLP analysis.

The DNA samples were diluted down to the concentration of  $22.2 \text{ ng } \mu\text{l}^{-1}$  concentration of genomic DNA by adding TE buffer (1×). The agarose gels were visualized under UV light by using the Genomics Solutions software. The quantified DNA product was stored at  $-20^\circ\text{C}$  until used for AFLP analysis.

#### AFLP methods for genetic analysis

A modified AFLP protocol (Vos *et al.*, 1995) was used to assess the genetic variability of *S. frugiperda* populations. It used the DNA extracted from seven individuals per population of *S. frugiperda* in order to keep consistent with the lowest number of insects available in a given population. The AFLP procedure was completed out in three basic steps:

(i) DNA template preparation, (ii) DNA template preamplification, and (iii) selective amplification. The DNA extracts were digested with *EcoRI* and *MseI* restriction enzymes and ligated with specific adapters (table 2). The samples were diluted ten-fold and used as a template for the preamplification and selective amplification. The selective primers selected for this analysis had three additional nucleotides at the 3' end of the preamplification primers and were end-labelled (table 3). The AFLP products were separated in 6.5% denaturing polyacrilamide gels visualized in a Li-Cor Gene Read IR 4200 DNA sequencer (LI-COR, Lincoln, Nebraska). The following electrophoresis conditions were used: 1500 volts, 40 watts, 40 mA of current, at  $45^\circ\text{C}$  for 2.5 h.

#### Data collection and analysis

AFLP bands were visualized using IRD-700 labeled 50–700 bp markers as a reference and scored using SAGA Generation 2 Software version 3.2 (LI-COR, Lincoln, Nebraska). Visibility and sharpness of bands were used as criteria for marker selection. The bands were identified by scoring the presence (1) or absence (0) of the bands among

Table 4. Matrix of genetic similarity obtained through AFLP markers based on the Jaccard index among populations of *Spodoptera frugiperda*.

Population	MEX	BR1	USA	ARG1	ARG2	ARG3	ARG4	BR2	BR3	BR4	BR5	BR6	BR7
MEX	1.00												
BR1	0.22	1.00											
USA	0.35	0.28	1.00										
ARG1	0.14	0.14	0.17	1.00									
ARG2	0.16	0.14	0.17	0.16	1.00								
ARG3	0.45	0.27	0.47	0.15	0.20	1.00							
ARG4	0.44	0.27	0.47	0.16	0.19	0.72	1.00						
BR2	0.36	0.23	0.39	0.15	0.18	0.50	0.49	1.00					
BR3	0.34	0.25	0.39	0.16	0.18	0.52	0.50	0.43	1.00				
BR4	0.37	0.24	0.39	0.15	0.19	0.52	0.53	0.43	0.44	1.00			
BR5	0.41	0.24	0.43	0.16	0.19	0.59	0.58	0.51	0.50	0.49	1.00		
BR6	0.38	0.24	0.41	0.16	0.19	0.56	0.54	0.47	0.46	0.44	0.55	1.00	
BR7	0.39	0.28	0.45	0.17	0.19	0.60	0.58	0.48	0.49	0.48	0.57	0.53	1.00

Table 5. Analysis of molecular variance (AMOVA) to explore geographical and host plant associated subdivision in populations of *Spodoptera frugiperda*.

Source of variation	DF	SSD	% of total variation	$\Phi_{ST}$	$N_m$
(a) Among populations	12	931.92	32.19*		
Within populations	76	1389.23	67.81*	0.3219	0.53
Total	88	2321.15	100.00		
(b) Among Brazil and Argentina	1	152.62	7.01*		
Among populations/regions	10	726.95	28.37*		
Within populations	70	1273.80	64.62*	0.3538	0.78
Total	81	2153.67	100.00		
(c) Among host plants	1	43.60	0.00*		
Among populations/host plants	5	230.95	16.65*	0.1588	1.32
Within populations	40	796.94	84.12*		
Total	46	1071.49	100.00		

\* $P < 0.001$ .

the *S. frugiperda* populations for every selective primer pair combination.

A binary data matrix (1 or 0) was created from the selected markers to estimate the Jaccard similarity index by using the SINGUAL procedure available in the Ntsys-2.1 software (Rohlf, 2000). Dendrograms were constructed using methods described by Sneath & Sokal (1973). A bootstrap procedure was used to evaluate the reliability of the dendrograms using 10,000 pseudo-replicates (BOOT-P software, version 3.1, Coelho, 2001). The goodness of fit of the dendrograms to the data was estimated by using the procedures COPH and MAXCOMP in Ntsys 2.1 (Rohlf, 2000). The correlation of geographical distance to genetic distance was performed using the Mantel test with 999 permutations (Mantel, 1967). The analysis of molecular variance (AMOVA) was performed to access the genetic structure and gene flow among populations of *S. frugiperda*. The AMOVA procedure partitioned the total variation both between and within *S. frugiperda* populations by the use of the squared distances as described by Excoffier *et al.* (1992) and by using the Arlequin program (Schneider *et al.*, 2000).

## Results and discussion

The results showed that four primer pair combinations (table 3) provided unambiguously scorable, clear, consistent, polymorphic and reproducible banding patterns. The mean genetic similarity among the populations obtained from

the Jaccard similarity matrix was 0.45. The unweighted pair group method with arithmetic mean analysis (UPGMA) dendrogram, based on the Jaccard algorithm, did not indicate a molecular classification of the populations of *S. frugiperda* associated with the host plant in which the insects were collected (fig. 1). For example, BR5 collected in cotton plants and BR7 collected in maize were the most similar populations from Brazil (table 4, fig. 1). The similarity matrix showed that ARG3 and ARG4 were the most similar *S. frugiperda* populations (0.72) (table 4). The lowest genetic similarity value was observed between the BR1 and ARG1 populations (0.14). The Mantel test of genetic isolation by distance showed that genetic dissimilarity and geographical distances were not significantly correlated ( $n = 78$ ,  $r = 0.04$ ,  $P = 0.3917$ ).

The overall non-hierarchical AMOVA test (table 5a) revealed that 32% of the total variation could be attributed to the variability among the tested populations. The hierarchical AMOVA (table 5b,c) analysis revealed that the local populations were the major reservoir of genetic variability. Approximately 7% of the total variation was related to the arbitrary subdivision within the Brazilian and Argentine populations (table 5b). In addition, the AMOVA results revealed no molecular variation between the maize and cotton groups of populations of *S. frugiperda* collected in Brazil. This information indicated no significant structuring within the *S. frugiperda* populations associated with maize and cotton crops (table 5c).

The AFLP analysis provided appropriate results to assess the genetic variability and population structure of populations of *S. frugiperda* associated with maize and cotton in Brazil. Moreover, high values of molecular similarity were found between the populations of fall armyworm collected in Brazil and USA (Florida). Similar results were found with the velvetbean caterpillar, *Anticarsia gemmatalis* (Hübner) (Lepidoptera: Noctuidae) (Sosa-Gomez, 2004). The results obtained from the present study and those gathered by Sosa-Gomez (2004) suggest that certain populations of Noctuidae insects in South America may be composed in their majority by non-migratory or permanently resident genotypes. Nevertheless, migration is a complex attribute with great evolutionary flexibility in its own appearance and disappearance. The balance between migratory and resident individuals may even occur in populations of the same species (Alerstam *et al.*, 2003).

None of the dendrograms showed a branch or a major cluster that could separate the populations of *S. frugiperda* into distinctive groups associated with the maize or cotton crop. This result has suggested that different lineages of this pest associated with maize and cotton plants have not been established in Brazil. Moreover, it could not be detected as a significant portion of genetic variability associated with the geographical origin of the populations. However, the detection of few differences among *S. frugiperda* populations may also be due to the limited resolution of the AFLP markers or high gene flow rates. Both processes would result in the lack of an association between genetic differences and geographic distance, and they could hardly be distinguished from each other.

The higher values of variation ( $\Phi_{ST}$ ) are common in species characterized by a closed population structure. The hierarchical AMOVA (table 5b,c) analysis confirmed that the local populations are the major reservoir of genetic variability. Gene flow is the evolutionary force which counteracts the effects of genetic drift and selection, ultimately resulting in homogenization of the genetic background of populations at greater geographical scales (Vandewoestijne *et al.*, 1999). When the homogenization effect doesn't take place, higher values of variation among populations can be detected. The percentage of variation found in the present study was most likely to be due to the non-hierarchical AMOVA analysis taking into account populations from South America (Brazil and Argentina) and North America (Mexico and USA). These populations were collected in areas with different agro-ecosystems. The large geographic area and the different cropping systems that were used for sampling are most likely to be contributing factors to the observed differences in the genetic structure of *S. frugiperda* in this study. The majority of molecular markers studies performed with *S. frugiperda* have been focused on the maize and rice host strains issue (Nagoshi & Meagher, 2003a,b; Prowell *et al.*, 2004). Recently, genetic variation within and between *S. frugiperda* host strains has been assessed by analysing the ITS-1 region (Lewter *et al.*, 2006). The authors also found high values of overall  $\Phi_{ST}$  values which combined with the hierarchical analysis output suggested structuring of *S. frugiperda* populations between states in the USA. According to Lewter *et al.* (2006), this variation was possibly due to the distribution of fall armyworm strains. In order to further understand the genetic structure of an organism across a larger geographical scale, efforts should be made to approach the issue by

combining landscape ecology and population genetics (Manel *et al.*, 2003).

The hierarchical AMOVA (table 5c) revealed a significant value of immigrants ( $Nm=1.32$ ) between maize and cotton. In population genetics studies, the values of  $Nm > 1$  are considered sufficient enough to make gene flow overcome genetic drift (Wright, 1931) and consequently acts in favour of the homogenization of the genetic background of populations. Different papers have addressed the estimation of gene flow and dispersal rates by using molecular markers and their impact on population genetics (Milligan *et al.*, 1994; Bossart & Prowell, 1998a), but there has been a heated debate about the usefulness of these estimates (Bohonak *et al.*, 1998; Bossart & Prowell, 1998b). Indirect estimates of gene flow and dispersal rates have been extensively criticized. Particularly since those estimates rely on a mathematical relationship between the genetic structure and the rate of gene flow, such estimates implicitly assume that the ecological properties of the populations from which the genetic data were taken match the often unrealistic assumptions of the theoretical model upon which that mathematical relationship was based (Whitlock & McCauley, 1999).

Despite the limitations of the traditional approaches for indirect estimation of gene flow, they still provide a first scenario for many cases (Bohonak *et al.*, 1998). The information reported herein provided insight to integrated pest management scientists because *S. frugiperda* is well known to feed on both crops. The AMOVA results also showed a 0% variation in Brazil among those populations of *S. frugiperda*. The results of this study are particularly important when considering the potential for a mixed agro-ecosystem of Bt maize and Bt cotton. This study will help researchers to more clearly define insect resistance management techniques for conventional agricultural practices, as well as for genetically modified crops. The constant genetic mixing of resident and immigrant populations of *S. frugiperda* has the potential to impact insect resistance management programmes (Roush & McKenzie, 1987; Roush & Daly, 1990). Therefore, further studies should be concentrated on the estimation of the gene flow rate between *S. frugiperda* populations associated with different crops and wild hosts over the crop seasons in Brazil.

### Acknowledgements

The authors acknowledge Capes, FAPESP and CNPq for funding this research and the Brazilian Insecticide Resistance Action Committee (IRAC-BR), Dr David Isenhour, Dr Jaime Molina-Ochoa and Claudia Gianni for providing the logistic support for collecting populations.

### References

- Alerstam, T., Henderstrom, A. & Akesson, S. (2003) Long-distance migration: evolution and determinants. *Oikos* **103**, 247–260.
- Black, W.C., & Duteau, N.M. (1997) RAPD-PCR and SSCP analysis for insect population genetic studies. pp. 361–373 in Crampton, J.M., Beard, C.B. & Louis, C. (Eds) *The molecular biology of insect disease vectors: a methods manual*. London, Chapman & Hall.

- Bohonak, A.J., Davies, N., Roderick, G.K. & Villablanca, F.X. (1998) Is population genetic mired on the past? *Trends in Ecology and Evolution* **13**, 360.
- Bossart, J.L. & Prowell, D.P. (1998a) Genetic estimates of population structure and gene flow: limitations, lesson and new directions. *Trends in Ecology and Evolution* **13**, 202–206.
- Bossart, J.L. & Prowell, D.P. (1998b) Reply from J.L. Bossart and D. Pashley Prowell. *Trends in Ecology and Evolution* **13**, 360.
- Busato, G.R., Grutzmacher, A.D., Garcia, M.S., Giolo, F.P. & Martin, A.F. (2002) Consumo e utilização de alimento por *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) originária de diferentes regiões do Rio Grande do Sul, nas culturas do milho e arroz irrigado. *Neotropical Entomology* **31**, 525–529.
- Busato, G.R., Grutzmacher, A.D., Oliveira, A.C., Vieira, E.A., Zimmer, P.D., Kopp, M.M., Bandeira, J.M. & Magalhães, T.R. (2004) Análise da estrutura e diversidade molecular de populações de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) associadas às culturas do milho e arroz no Rio Grande do Sul. *Neotropical Entomology* **33**, 709–716.
- Caprio, M.A. & Tabashnik, B.E. (1992) Gene flow accelerates local adaptation among finite populations: simulating the evolution of insecticide resistance. *Journal of Economic Entomology* **85**, 611–620.
- Coelho, A.S.G. (2001) BOOD-P Avaliação de dendogramas baseados em estimativas de distâncias/similaridades genéticas através do procedimento de bootstrap, Versão 3.0. Departamento de Biologia Geral, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Goiânia, GO.
- Diez-Rodriguez, G.I. & Omoto, C. (2001) Herança da resistência de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) à lambda-cialotrina. *Neotropical Entomology* **30**, 311–316.
- Dobzhansky, T. (1951) *Genetics and the origin of species*. 3rd edn. New York, Columbia University Press.
- Excoffier, L., Smouse, P.E. & Quattro, J.M. (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**, 479–491.
- Figuroa, C.C., Loyayza-Muro, R. & Niemeyer, H.M. (2002) Temporal variation of RAPD-PCR phenotype composition of the grain aphid *Sitobion avenae* (Hemiptera: Aphididae) on wheat: the role of hydroxamic acids. *Bulletin of Entomological Research* **91**, 148–156.
- Fuentes-Contreras, E., Figuroa, C.C., Reyes, M., Briones, L.M. & Niemeyer, H.M. (2004) Genetic diversity and insecticide resistance of *Myzuz persicae* (Hemiptera: Aphididae) populations from tobacco in Chile: evidence for the existence of a single predominant clone. *Bulletin of Entomological Research* **94**, 11–18.
- Georghiou, G.P. & Taylor, C.E. (1977a) Genetic and biological influences in the evolution of insecticide resistance. *Journal of Economic Entomology* **70**, 319–323.
- Georghiou, G.P. & Taylor, C.E. (1977b) Operational influences in the evolution of insecticide resistance. *Journal of Economic Entomology* **70**, 653–658.
- Johnson, S.J. (1988) Migration and life history strategy of fall armyworm, *Spodoptera frugiperda*, in the Western Hemisphere. *Insect Science and Its Application* **8**, 543–549.
- Labbe, P., Lenormand, T. & Raymond, M. (2005) On the world wide spread of an insect resistance gene: a role for local selection. *Journal of Evolutionary Biology*, on line publication 23 May 2005.
- Levy, H.C., Garcia-Maruniak, A. & Maruniak, J.E. (2002) Strain identification of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) insects and cell line: PCR-RFLP of cytochrome oxidase c subunit I gene. *Florida Entomologist* **85**, 186–190.
- Lewter, J.A., Szalanski, A.L., Nagoshi, R.N., Meagher, R.L., Owens, C.B. & Luttrell, R.G. (2006) Genetic variation within and between strains of the fall armyworm, *Spodoptera frugiperda*, (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* **89**, 63–68.
- Manel, S., Schwarz, M.K., Luikart, G. & Taberlet, P. (2003) Landscape genetics: combining landscape ecology and populations genetics. *Trends in Ecology and Evolution* **18**, 189–197.
- Mantel, N. (1967) The detection of disease clustering and a generalized regression approach. *Cancer Research* **27**, 209–220.
- Martinelli, S., Montrazi, R.B., Zucchi, M.I., Silva-Filho, M.C. & Omoto, C. (2006) Molecular variability of *Spodoptera frugiperda* populations associated to maize and to cotton in Brazil. *Journal of Economic Entomology*, **99**, 516–526.
- McKenzie, J.A. (1996) *Ecological and evolutionary aspects of insecticide resistance*. 185 pp. New York, Academic Press.
- McMichael, M. & Prowell, D.P. (1994) Differences in amplified fragment-length polymorphisms in fall armyworm (Lepidoptera: Noctuidae) host strains. *Annals of the Entomological Society of America* **92**, 175–181.
- Meagher, R.N. & Nagoshi, R.L. (2004) Behavior and distribution of the two fall armyworm strains in Florida. *Florida Entomologist* **87**, 440–449.
- Meagher, R.N., Nagoshi, R.L., Stuhl, C. & Mitchell, E.R. (2004) Larval development of fall armyworm (Lepidoptera: Noctuidae) in different cover crops. *Florida Entomologist* **87**, 454–460.
- Meldenson, T. & Shaw, K.L. (2002) Genetic and behavioral component of the cryptic species boundary between *Laupala cerasiana* and *L. kohalensis* (Orthoptera: Gryllidae). *Genetica* **116**, 301–310.
- Milligan, B.G., Leebens-Mack, J. & Strand, A.E. (1994) Conservation genetics beyond the maintenance of marker diversity. *Molecular Ecology* **3**, 423–435.
- Nagoshi, R.N. & Meagher, R. (2003a) Fall armyworm *FR* sequences map to sex chromosomes and their distribution in the wild indicate limitation in the interstrain mating. *Insect Molecular Biology*, **12**, 453–458.
- Nagoshi, R.N. & Meagher, R. (2003b) *FR* tandem-repeat sequence in (Lepidoptera: Noctuidae) host strains. *Annals of the Entomological Society of America*, **96**, 329–335.
- Pashley, D.P., Johnson, S.J. & Sparks, A.N. (1985) Genetic population structure of migratory moths: the fall armyworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* **78**, 756–762.
- Pitre, H.W. (1988) Relationship of fall armyworm (Lepidoptera: Noctuidae) from Florida, Honduras, Jamaica and Mississippi: susceptibility to insecticides with reference to migration. *Florida Entomologist* **71**, 56–61.
- Prowell, D.P., McMichael, M. & Silvain, J.F. (2004) Multilocus genetic analysis of host use, introgression and speciation in host strains of fall armyworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* **97**, 1034–1044.
- Roderick, K.R. (1996) Geographic structure of insect populations: gene flow, phylogeography, and their uses. *Annual Review of Entomology* **41**, 325–352.
- Rohlf, F.J. (2000) NTSYS-pc Numerical taxonomy and multivariate analysis system, version 2.1. Owner's manual.



- Roush, R.T. & Daly, J.C. (1990) The role of population genetics in resistance research and management. pp. 97–152 in Roush, R.T. & Tabashnik, B.E. (Eds) *Pesticide resistance in arthropods*. New York, Chapman and Hall.
- Roush, R.T. & McKenzie, J.A. (1987) Ecological genetics of insecticide and acaricide resistance. *Annual Review of Entomology* **32**, 361–380.
- Schneider, S., Kueffer, J.M., Roessli, D. & Excoffier, L. (2000) ARLEQUIN version 2.0. A software for population genetic data analysis. Genetic and Biometry Laboratory, University of Geneva, Switzerland.
- Sena Jr, D.G., Pinto, F.A.C., Queiroz, D.M. and Viana, P.A. (2003) Fall armyworm damaged maize plant identification using digital images. *Biosystems Engineering* **85**, 449–454.
- Slatkin, M. (1985) Gene flow in natural populations. *Annual Review of Evolution and Systematics* **16**, 393–430.
- Sneath, P.H.A. & Sokal, R.R. (1973) *Numerical taxonomy*. 573 pp. San Francisco, W.H. Freeman and Company.
- Sosa-Gomez, D.R. (2004) Intraspecific variation and population structure of the velvetbean caterpillar, *Anticarsia gemmatilis* Hübner, and 1818 (Insecta: Lepidoptera: Noctuidae). *Genetics and Molecular Biology* **27**, 378–384.
- Sparks, A.N. (1979) A review of the biology of the fall armyworm. *Florida Entomologist* **62**, 82–87.
- Tabashnik, B.E. (1991) Determining the mode of inheritance of pesticide resistance with blackcross experiments. *Journal of Economic Entomology* **38**, 25–326.
- Timm, A.E., Pringle, K.L. & Warnich, L. (2005) Genetic diversity of woolly apple aphid *Eriosoma lanigerum* (Hemiptera: Aphididae) populations in the Western Cape, South Africa. *Bulletin of Entomological Research* **95**, 187–191.
- Vandewoestijne, S., Neve, G. & Baguette, M. (1999) Spatial and temporal population genetic structure of the butterfly *Aglais urticae* L. (Lepidoptera, Nymphalidae). *Molecular Ecology* **8**, 1539–1543.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., Van de Lee, T., Hornes, M., Frijters, A., Peleman, J., Kuiper, M. & Zabeau, M. (1995) AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Research* **23**, 4407–4414.
- Whitlock, M.C. & McCauley, D.E. (1999) Indirect measures of gene flow:  $F_{st} \neq 1/(4Nm + 1)$ . *Heredity* **82**, 117–125.
- Wright, S. (1931) Evolution in Mendelian populations. *Genetics* **16**, 97–157.
- Yu, S.J. (1991) Insecticide resistance in the fall armyworm *Spodoptera frugiperda* (J.E. Smith). *Pesticide Biochemistry and Physiology* **39**, 84–91.
- Yu, S.J., Nguyen, S.N. & Abo-Elghar, G.E. (2003) Biochemical characteristics of insecticide resistance in the fall armyworm *Spodoptera frugiperda* (J.E. Smith). *Pesticide Biochemistry and Physiology* **77**, 1–11.
- Zhang, L.P., Zhang, Y.J., Zhang, W.J., Wu, Q.J., Xu, B.Y. & Xu, D. (2005) Analysis of genetic diversity among different geographical populations and determination of biotypes of *Bemisia tabaci* in China. *Journal of Applied Entomology* **129**, 121–128.

(Accepted 18 October 2006)

© 2007 Cambridge University Press